

01-92502-14

Original Effective Date: 06/15/05

Reviewed: 12/08/23

Revised: 01/01/24

Subject: Vestibular Rehabilitation

THIS MEDICAL COVERAGE GUIDELINE IS NOT AN AUTHORIZATION, CERTIFICATION, EXPLANATION OF BENEFITS, OR A GUARANTEE OF PAYMENT, NOR DOES IT SUBSTITUTE FOR OR CONSTITUTE MEDICAL ADVICE. ALL MEDICAL DECISIONS ARE SOLELY THE RESPONSIBILITY OF THE PATIENT AND PHYSICIAN. BENEFITS ARE DETERMINED BY THE GROUP CONTRACT, MEMBER BENEFIT BOOKLET, AND/OR INDIVIDUAL SUBSCRIBER CERTIFICATE IN EFFECT AT THE TIME SERVICES WERE RENDERED. THIS MEDICAL COVERAGE GUIDELINE APPLIES TO ALL LINES OF BUSINESS UNLESS OTHERWISE NOTED IN THE PROGRAM EXCEPTIONS SECTION.

[Position Statement](#)

[Billing/Coding](#)

[Reimbursement](#)

[Program Exceptions](#)

[Definitions](#)

[Related Guidelines](#)

[Other](#)

[References](#)

[Updates](#)

DESCRIPTION:

Vestibular rehabilitation therapy (VRT) is a form of physical therapy that uses specialized exercises that result in gaze and gait stabilization. Most VRT exercises involve head movement, and head movements are essential in stimulating and retraining the vestibular system. Vestibular rehabilitation therapy is a modality for treatment of disorders of the vestibular or central balance system.

Evaluation for vestibular rehabilitation may include one or more of the following types of vestibular testing:

- Caloric vestibular testing [electronystagmography (ENG) or infrared videonystagmography (VNG)]
- Clinical head shaking test
- Rotational chair testing (computer-driven chair rotations)
- Passive examiner-generated head rotation testing
- Active head rotation (self-generated head turns)
- Hyperventilation induced nystagmus test
- Visual fixation of vestibular nystagmus maneuver
- Dynamic or head shaking acuity testing
- Head impulse or head thrust test
- Optokinetic nystagmus test
- Spontaneous nystagmus test

- Valsalva test for nystagmus
- Vibration induced nystagmus testing (VIN)
- Skull vibration induced nystagmus testing (SVINT)
- Bone conduction vibration
- Subjective visual vertical testing
- Saccades (calibration) test
- Vestibular autorotation (VAT)

Assessment should also include a complete medical history and a detailed history of the balance symptoms including a description of the type of symptoms (e.g., vertigo, imbalance, disequilibrium, pre-syncope sensations, gait ataxia), frequency and duration of symptoms, specific activities or positions that provoke symptoms, presence of visual disturbances, and the individual’s perception of the impact of the symptoms on daily activities.

POSITION STATEMENT:

Vestibular rehabilitation **meets the definition of medical necessity** for the treatment of chronic vertigo when **ALL** of the following criteria are met:

- The individual has a diagnosis of a vestibular disorder (eg, Ménière’s disease, vertigo, benign paroxysmal positioning vertigo) or has had ablative vestibular surgery
- Symptoms of vertigo and imbalance have existed for duration of 8 weeks or more
- The individual has persistent symptoms despite optimal medical management such as vestibular suppressant medication prescribed to reduce symptoms

BILLING/CODING INFORMATION:

The following codes may be used to describe vestibular rehabilitation:

HCPCS Coding:

S9476	Vestibular rehabilitation program, non-physician provider, per diem
-------	---

REIMBURSEMENT INFORMATION:

NOTE: Vestibular rehabilitation services are considered part of the contract benefit for rehabilitative services.

LOINC Codes:

The following information may be required documentation to support medical necessity: physician history and physical, physician progress notes, treatment plan, medication history and operative report (if applicable).

Documentation Table	LOINC Codes	LOINC Time Frame Modifier Code	LOINC Time Frame Modifier Codes Narrative
---------------------	-------------	--------------------------------	---

Physician history and physical	28626-0	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim
Attending physician visit note	18733-6	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.
Treatment plan	18776-5	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.
History of medication use	10160-0	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.
Surgical report	28573-4	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim

PROGRAM EXCEPTIONS:

Federal Employee Program (FEP): Follow FEP guidelines.

State Account Organization (SAO): Follow SAO guidelines.

Medicare Advantage products: No National Coverage Determination (NCD) and/or Local Coverage Determination (LCD) were found at the time of the last guideline reviewed date.

If this Medical Coverage Guideline contains a step therapy requirement, in compliance with Florida law 627.42393, members or providers may request a step therapy protocol exemption to this requirement if based on medical necessity. The process for requesting a protocol exemption can be found at [Coverage Protocol Exemption Request](#)

DEFINITIONS:

Sacculle: the smaller chamber of the membranous labyrinth of the ear.

Vertigo: the sensation of moving around in space (subjective vertigo) or of having objects move about the person (objective vertigo).

Vestibular: of or relating to the vestibule of the inner ear, the vestibular apparatus, the vestibular nerve, or the labyrinthine sense.

RELATED GUIDELINES:

01-92502-12, Computerized Dynamic Posturography

OTHER:

None applicable.

REFERENCES:

1. American Speech-Language-Hearing Association. Balance System Disorders (2020). Accessed at <https://www.asha.org/PRPSpecificTopic.aspx?folderid=8589942134§ion=Assessment>.
2. American Speech-Language-Hearing Association. Role of Audiologist in Vestibular and Balance Rehabilitation: Position Statement (1999).
3. Angeli SI, Hawley R, Gomez O. Systematic approach to benign paroxysmal positional vertigo in the elderly. *Otolaryngol Head Neck Surg.* 2003 May; 128(5): 719-25.
4. Baylor College of Medicine website: Vestibular Rehabilitation. Last modified 02/14/11.
5. Bhattacharyy, N, et al. Clinical practice guideline: Benign paroxysmal positional vertigo. *Otolaryngology–Head and Neck Surgery* (2008) 139, S47-S81.
6. Blue Cross Blue Shield Association. Medical Policy Reference Manual. Canalith Repositioning as a Treatment of Benign Paroxysmal Positional Vertigo (BPPV) 2.01.37, 12/15/00 (archived 07/08/10).
7. Blue Cross Blue Shield Association Evidence Positioning System®. 2.01.104 - Vestibular Function Testing, 03/22.
8. Cohen HS, Kimball KT. Changes in repetitive head movement task after vestibular rehabilitation. *Clin Rehabil.* 2004 Mar; 18(2): 125-31.
9. Cohen HS, Kimball KT. Decreased ataxia and improved balance after vestibular rehabilitation. *Otolaryngol Head Neck Surg.* 2004 Apr; 130(4): 418-25.
10. Cohen HS, Kimball KT. Increased independence and decreased vertigo after vestibular rehabilitation. *Otolaryngol Head Neck Surg.* 2003 Jan; 128 (1): 60-70.
11. Dumas G, et al. The Skull Vibration-Induced Nystagmus Test of Vestibular Function-A Review. *Front Neurol.* 2017 Mar 9;8:41.
12. ECRI Windows on Technology. “Vestibular Rehabilitation and Particle Repositioning for Benign Positional Vertigo” (01/03).
13. García-Muñoz C, Cortés-Vega MD, Heredia-Rizo AM, Martín-Valero R, García-Bernal MI, Casuso-Holgado MJ. Effectiveness of Vestibular Training for Balance and Dizziness Rehabilitation in People with Multiple Sclerosis: A Systematic Review and Meta-Analysis. *J Clin Med.* 2020;9(2):590. Published 2020 Feb 21. doi:10.3390/jcm9020590.
14. Hall CD, Herdman SJ, Whitney SL, et al. Vestibular Rehabilitation for Peripheral Vestibular Hypofunction: An Updated Clinical Practice Guideline From the Academy of Neurologic Physical Therapy of the American Physical Therapy Association. *J Neurol Phys Ther.* 2022 Apr 1;46(2):118-177. doi: 10.1097/NPT.0000000000000382.
15. Hayes Medical Technology Directory, Vestibular Rehabilitation for the Treatment of Vestibular and Balance Disorders, VEST0501.04, (10/99; updated 09/07/05).
16. Huh YE, Kim JS. Evaluation of Dizzy Patients. *J Clin Neurol.* 2013 Oct;9(4):203-13.
17. Iwasaki S, Yamasoba T. Dizziness and Imbalance in the Elderly: Age-related Decline in the Vestibular System. *Aging Dis.* 2014 Feb 9;6(1):38-47.

18. Krebs DE, Gill-Body KM, Parker SW, et al. Vestibular rehabilitation: useful but not universally so. *Otolaryngol Head Neck Surg.* 2003 Feb; 128(2): 240-50.
19. Pavlou M, Lingeswaran A, Davies, RA, Gresty MA, Bronstein AM. Simulator based rehabilitation in refractory dizziness. *J Neurol.* 2005 Aug; 251(8): 983-95.
20. Rine, RM, Braswell J, Fosher D, Joyce K, Kalar K, Shaffer M. Improvement of motor development and postural control following intervention in children with sensorineural hearing loss and vestibular impairment. *Int J Pediatr Otorhinolaryngol.* 2004 Sept; 68(9): 1141-8.
21. Smółka W, Smółka K, Markowski J, Pilch J, Piotrowska-Seweryn A, Zwierzchowska A. The efficacy of vestibular rehabilitation in patients with chronic unilateral vestibular dysfunction. *Int J Occup Med Environ Health.* 2020;33(3):273-282. doi:10.13075/ijomeh.1896.01330.
22. Strupp M, Długaiczek J, Ertl-Wagner BB, Rujescu D, Westhofen M, Dieterich M. Vestibular Disorders. *Dtsch Arztebl Int.* 2020;117(17):300-310. doi:10.3238/arztebl.2020.0300.
23. UpToDate. Benign paroxysmal positional vertigo. 2022. Accessed at uptodate.com.
24. UpToDate. Treatment of vertigo. 2021. Accessed at uptodate.com.
25. Xie S, et al. Vibration-induced nystagmus in patients with unilateral peripheral vestibular disorders. *Indian J Otolaryngol Head Neck Surg.* 2013 Dec;65(4):333-8.
26. Yardley L., Donovan-Hall M, Smith HE, Walsh BM, Mullee M, Bronstein AM. Effectiveness of primary care-based vestibular rehabilitation for chronic dizziness. *ANN Intern Med.* 2004 Oct 19; 141(8): 641-3.
27. Zapanta PE, Meyers AD, et al. Vestibular Rehabilitation (Medscape). July 2020.

COMMITTEE APPROVAL:

This Medical Coverage Guideline (MCG) was approved by the Florida Blue Medical Policy and Coverage Committee on 12/08/23.

GUIDELINE UPDATE INFORMATION:

06/15/05	New Medical Coverage Guideline.
06/15/07	Scheduled review; reformatted guideline; updated references.
06/15/09	Scheduled review of guideline. Update position statement and ICD 9 coding section. Remove reference to canalith repositioning guideline.
10/15/10	Revision; related ICD-10 codes added.
06/15/11	Scheduled review; Position Statement unchanged; references updated; formatting changes.
09/15/11	Revision; formatting changes.
02/15/14	Revision; Program Exceptions section updated.
11/01/15	Revision: ICD-9 Codes deleted.
10/01/16	Revision: Billing/Coding Information section updated.
10/15/18	Revision: Updated description, related guidelines, and references. Reformatted guideline.
09/15/20	Scheduled review. Revised description. Maintained position statement and updated references.
06/15/22	Scheduled review. Revised description, maintained position statement, and updated references.
05/22/23	Update to Program Exceptions section.

01/01/24	Position statements maintained.
----------	---------------------------------