

01-93000-26

Original Effective Date: 01/01/00

Reviewed: 08/23/18

Revised: 09/15/18

Subject: External Counterpulsation (ECP)

THIS MEDICAL COVERAGE GUIDELINE IS NOT AN AUTHORIZATION, CERTIFICATION, EXPLANATION OF BENEFITS, OR A GUARANTEE OF PAYMENT, NOR DOES IT SUBSTITUTE FOR OR CONSTITUTE MEDICAL ADVICE. ALL MEDICAL DECISIONS ARE SOLELY THE RESPONSIBILITY OF THE PATIENT AND PHYSICIAN. BENEFITS ARE DETERMINED BY THE GROUP CONTRACT, MEMBER BENEFIT BOOKLET, AND/OR INDIVIDUAL SUBSCRIBER CERTIFICATE IN EFFECT AT THE TIME SERVICES WERE RENDERED. THIS MEDICAL COVERAGE GUIDELINE APPLIES TO ALL LINES OF BUSINESS UNLESS OTHERWISE NOTED IN THE PROGRAM EXCEPTIONS SECTION.

[Position Statement](#)

[Billing/Coding](#)

[Reimbursement](#)

[Program Exceptions](#)

[Definitions](#)

[Related Guidelines](#)

[Other](#)

[References](#)

[Updates](#)

DESCRIPTION:

External counterpulsation (ECP), also known as enhanced external counterpulsation (EECP), is a noninvasive treatment that uses timed, sequential inflation of pressure cuffs on the calves, thighs and buttocks to augment diastolic pressure, decrease left ventricular afterload, and increase venous return. The mechanism of action is the augmentation of diastolic pressure by displacement of a volume of blood backward into the coronary arteries during diastole when the heart is in a state of relaxation and the resistance in the coronary arteries is at a minimum. The resulting increase in coronary artery perfusion pressure may enhance coronary collateral development or increase flow through existing collaterals. In addition, when the left ventricular contracts, it faces a reduced aortic counterpressure because the counterpulsation has somewhat emptied the aorta. ECP has been primarily investigated as a treatment for chronic stable angina.

Under direct supervision of a physician, a full course of therapy usually consists of 35 one-hour treatments, which may be offered once or twice daily, usually 5 days per week. The multiple components of the procedure include the use of the device itself, finger plethysmography to follow the blood flow, continuous electrocardiograms to trigger inflation and deflation, and the optional use of pulse oximetry to measure oxygen saturation before and after treatment.

Multiple external counterpulsation devices have been cleared for marketing by the Food and Drug Administration (FDA) through the 510(k) process.

POSITION STATEMENT:

External counterpulsation (using FDA-approved external counterpulsation systems) **meets the definition of medical necessity** for members with disabling stable angina (Class III or Class IV based on the

Canadian Cardiovascular Society Functional Classification scale or equivalent*) **AND** who are not amenable to surgical intervention, such as percutaneous transluminal coronary angioplasty (PTCA) or coronary artery bypass graft (CABG) due to any of the following:

- Condition is inoperable; **OR**
- High risk of operative complications or post-operative failure; **OR**
- Coronary anatomy is not readily amenable to such procedures; **OR**
- Co-morbid states that create excessive risk.

NOTE: This procedure must be performed under direct supervision of a physician.

Hydraulic versions of external counterpulsation devices **do not meet the definition of medical necessity**. The standard of practice for external counterpulsation does not include the use of hydraulic ECP devices.

External counterpulsation is considered **experimental or investigational** for all other indications, including but not limited to unstable angina, congestive heart failure, acute myocardial infarction and cardiogenic shock. The evidence is insufficient to determine the effects of the technology on health outcomes.

***Functional Classification Scales:**

Class	Canadian Cardiovascular Society Functional Classification	New York Heart Association (NYHA) Functional Classification
I	Usual physical activity, eg walking or climbing stairs, does not cause angina; angina is evoked by strenuous and/or rapid work or recreation.	Cardiac disease, but no symptoms and no limitation in ordinary physical activity, e.g. no shortness of breath when walking, climbing stairs etc.
II	Slight limitation of ordinary activities, eg after walking 2 blocks, climbing one flight of steps, under normal circumstances, after meals, in the cold, wind, in the morning, or when under emotional stress.	Mild symptoms (mild shortness of breath and/or angina) and slight limitation during ordinary activity.
III	Marked limitation of ordinary activities, eg walking 1-2 blocks or climbing stairs under normal circumstances.	Marked limitation in activity due to symptoms, even during less-than-ordinary activity, e.g. walking short distances (20–100 m). Comfortable only at rest.
IV	Inability to carry out any physical activity without discomfort—angina may be present at rest.	Severe limitations. Experiences symptoms even while at rest. Mostly bedbound patients.

BILLING/CODING INFORMATION:

HCPSC Coding:

G0166	External counterpulsation per treatment session
-------	---

ICD-10 Diagnosis Codes That Support Medical Necessity:

I20.8	Other forms of angina pectoris
I20.9	Angina pectoris, unspecified
I25.118 – I25.119, I25.708 – I25.709, I25.718 – I25.719, I25.728 – I25.729, I25.738 – I25.739, I25.758 – I25.759, I25.768 – I25.769, I25.798 – I25.799	Atherosclerosis

REIMBURSEMENT INFORMATION:

External counterpulsation (G0166) is limited to 35 sessions in a twelve (12) month period.

PROGRAM EXCEPTIONS:

Federal Employee Program (FEP): Follow FEP guidelines.

State Account Organization (SAO): Follow SAO guidelines.

Medicare Advantage products:

The following National Coverage Determination (NCD) was reviewed on the last guideline reviewed date:
External Counterpulsation (ECP) Therapy for Severe Angina (20.20) located at cms.gov.

DEFINITIONS:

None Applicable

RELATED GUIDELINES:

[Transmyocardial Revascularization \(TMR\), 02-33000-19](#)

OTHER:

The following information may be required documentation to support medical necessity: Physician history and physical, attending physician treatment plan, progress or hospital notes, and treatment notes including documentation of Class III or Class IV disabling stable angina.

LOINC Codes:

Documentation Table	LOINC Codes	LOINC Time Frame Modifier Code	LOINC Time Frame Modifier Codes Narrative
Physician history and physical	28626-0	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.
Attending physician visit note	18733-6	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.
Physician Initial Assessment	18736-9	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.
Attending physician progress note	18741-9	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.
Physician hospital discharge summary	11490-0	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.

REFERENCES:

1. Anderson JL, Adams CD, Antman EM, Bridges CR, Califf RM, Casey DE Jr, Chavey WE II, Fesmire FM, Hochman JS, Levin TN, Lincoff AM, Peterson ED, Theroux P, Wenger NK, Wright RS. ACC/AHA 2007 guidelines for the management of patients with unstable angina/non–ST-elevation myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 2002 Guidelines for the Management of Patients With Unstable Angina/Non–ST-Elevation Myocardial Infarction): developed in collaboration with the American College of Emergency Physicians, American College of Physicians, Society for Academic Emergency Medicine, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *J Am Coll Cardiol* 2007; 50:e1–157.
2. Beck DT, Casey DP, et al, Enhanced external counterpulsation reduces indices of central blood pressure and myocardial oxygen demand in patients with left ventricular dysfunction. *Clin Exp Pharmacol Physiol*. 2015 Apr;42(4):315-20.
3. Blue Cross Blue Shield Association Medical Policy. Reference Manual 2.02.06 Enhanced External Counterpulsation, 05/18.
4. Blue Cross Blue Shield Association Technology Evaluation Center Assessment. External Counterpulsation for Treatment of Chronic Stable Angina Pectoris and Chronic Heart Failure. Volume 20, No. 12, 12/05.
5. Centers for Medicare & Medicaid Services (CMS) National Coverage Determination (NCD) for External Counterpulsation (ECP) Therapy for Severe Angina (20.20); accessed at cms.gov.
6. ClinicalTrials.gov, Effect of External Counter Pulsation on Coronary Artery Disease (ECP-CAD), sponsored by Eighth Affiliated Hospital, accessed 07/10/18.

7. ClinicalTrials.gov, A Prospective, Randomized Study of Enhanced External Counterpulsation for Patients With Erectile Dysfunction; sponsored by First Affiliated Hospital, Sun Yat-Sen University; accessed 07/10/18.
8. ClinicalTrials.gov, A Prospective, Randomized Study of Enhanced External Counterpulsation Therapy on Ventricular Function in Patients With Coronary Heart Disease; sponsored by Sun Yat-sen University; accessed 07/10/18.
9. ClinicalTrials.gov, Study of Efficacy of External CounterPulsation on Reducing the Prevalence of Vein Graft Failure and Improving Cardiac Function for Postoperative Coronary Artery Bypass Grafting Patients, sponsored by Mengya Liang; accessed 07/10/18.
10. Fihn SD, Blankenship JC, Alexander KP, et al. 2014 ACC/AHA/AATS/PCNA/SCAI/STS focused update of the guideline for the diagnosis and management of patients with stable ischemic heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines, and the American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Association, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. *J Am Coll Cardiol*. Nov 4 2014;64(18):1929-1949.
11. Gibbons RJ, Abrams J, Chatterjee K, Daley J, Deedwania PC, Douglas JS, Ferguson TB Jr, Fihn SD, Fraker TD Jr, Gardin JM, O'Rourke RA, Pasternak RC, Williams SV. American College of Cardiology; American Heart Association Task Force on practice guidelines (Committee on the Management of Patients With Chronic Stable Angina). ACC/AHA 2002 guideline update for the management of patients with chronic stable angina-summary article: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines (Committee on the Management of Patients with Chronic Stable Angina). *J Am Coll Cardiol*, 2003 Jan; 41(1): 159-68.
12. Michaels AD, Barsness GW, Soran O, Kelsey SF, Kennard ED, Hui JC, Lawson WE; International EECPPatient Registry Investigators. Frequency and efficacy of repeat enhanced external counterpulsation for stable angina pectoris (from the International EECPPatient Registry). *Am J Cardiol*. 2005 Feb 1; 95(3): 394-7.
13. Michaels AD, Raisinghani A, Soran O, de Lame PA, Lemaire ML, Kligfield P, Watson DD, Conti CR, Beller G. The effects of enhanced external counterpulsation on myocardial perfusion in patients with stable angina: a multicenter radionuclide study. *Am Heart J*. 2005 Nov; 150(5): 1066-73.
14. Ontario Ministry of Health and Long-Term Care. Enhanced external counterpulsation (EECP): health technology policy assessment update. Toronto: Medical Advisory Secretariat, Ontario Ministry of Health and Long-Term Care (MAS), 2006: 66.
15. Soran O, Kennard ED, Kfoury AG, Kelsey SF; IEPR Investigators. Two-year clinical outcomes after enhanced external counterpulsation (EECP) therapy in patients with refractory angina pectoris and left ventricular dysfunction (report from The International EECPPatient Registry). *Am J Cardiol*. 2006 Jan 1; 97(1): 17-20.

COMMITTEE APPROVAL:

This Medical Coverage Guideline (MCG) was approved by the BCBSF Medical Policy & Coverage Committee on 08/23/18.

GUIDELINE UPDATE INFORMATION:

01/01/00	Medical Coverage Guideline developed.
06/15/00	Revision of guideline consisting of the addition of code G0166.

08/23/01	Review of guideline with no changes.
07/15/03	Review and revision of guideline consisting of the addition of limitation.
04/15/05	Review and revision of guideline consisting of updated references and addition of limitation.
03/15/06	Review and revision of guideline consisting of updated references.
03/15/07	Review and revision of guideline consisting of updated references.
06/15/07	Reformatted guideline.
03/15/08	Review and revision of guideline consisting of updated references.
02/15/09	Review and revision of guideline consisting of updated references.
12/15/09	Annual review: position statements maintained, references updated.
10/15/10	Revision; related ICD-10 codes added.
07/15/11	Revision; formatting changes.
05/11/14	Revision: Program Exceptions section updated.
10/01/15	Revision; ICD10 coding section updated.
11/01/15	Revision: ICD-9 Codes deleted.
09/15/18	Review; coverage maintained; description, position statements, program exception, and references updated.