

02-31000-03

Original Effective Date: 06/15/25

Reviewed: 05/28/26

Revised: 06/15/26

Next Review: 05/27/27

Subject: Surgical Ablation for Treatment of Chronic Rhinitis

THIS MEDICAL COVERAGE GUIDELINE IS NOT AN AUTHORIZATION, CERTIFICATION, EXPLANATION OF BENEFITS, OR A GUARANTEE OF PAYMENT, NOR DOES IT SUBSTITUTE FOR OR CONSTITUTE MEDICAL ADVICE. ALL MEDICAL DECISIONS ARE SOLELY THE RESPONSIBILITY OF THE PATIENT AND PHYSICIAN. BENEFITS ARE DETERMINED BY THE GROUP CONTRACT, MEMBER BENEFIT BOOKLET, AND/OR INDIVIDUAL SUBSCRIBER CERTIFICATE IN EFFECT AT THE TIME SERVICES WERE RENDERED. THIS MEDICAL COVERAGE GUIDELINE APPLIES TO ALL LINES OF BUSINESS UNLESS OTHERWISE NOTED IN THE PROGRAM EXCEPTIONS SECTION.

[Position Statement](#)

[Billing/Coding](#)

[Reimbursement](#)

[Program Exceptions](#)

[Definitions](#)

[Related Guidelines](#)

[Other](#)

[References](#)

[Updates](#)

DESCRIPTION:

Cryoablation, radiofrequency, and laser ablation of the posterior nasal nerve (PNN) is proposed to improve symptoms of chronic rhinitis by blocking the PNN. These minimally invasive procedures can be performed under endoscopy.

Several devices have been approved by the U.S. Food and Drug Administration for ablation of the posterior nasal nerve (e.g., Clarifix, RhinAer, Neuromark™ System).

Summary and Analysis of Evidence

In an UpToDate article "Chronic nonallergic rhinitis" (Lieberman) "Chronic nonallergic rhinitis (NAR) is not a specific disease but rather a syndrome diagnosed by excluding other specific types of rhinitis. NAR is best defined by the chronic presence of one or more of the four following cardinal symptoms of rhinitis, in the absence of a specific etiology (such as an immunologic, infectious, pharmacologic, structural, hormonal, vasculitic, metabolic, or atrophic cause): sneezing, rhinorrhea, nasal congestion and postnasal drainage. Although there is no consensus about how long symptoms should be present to establish chronicity, some studies have utilized a minimum duration of over one year. The lack of clarity regarding the pathogenesis of this condition has led to an imprecise terminology and a number of synonyms used to refer to this entity. These include, idiopathic rhinitis, nonallergic, noninfectious rhinitis, intrinsic rhinitis and vasomotor rhinitis or nonallergic vasomotor rhinitis. Some experts use the term "nonallergic rhinitis" to refer more generally to any form of rhinitis that is noninfectious and nonallergic. Role of surgery — Several surgical approaches have been used in patients with severe chronic nonallergic rhinitis (NAR). These have been reported as uncontrolled case series. Such interventions may be helpful in patients with difficult symptoms that are refractory to multiple therapies (e.g., corticosteroid nasal spray in combination with azelastine and/or decongestants and/or ipratropium nasal spray). Six to 12 months of medical management should be allowed before surgical options are considered. Studies of efficacy are lacking. A number of other surgical procedures have been tried in the past, including vidian nerve resection, electrocoagulation of anterior ethmoidal nerve, sphenopalatine ganglion block, and others]. None of these techniques have been shown to have long-term benefits, and the potential risks (e.g., persistent pain) have to be considered carefully since they may outweigh any possible benefits. However,

this is an area of ongoing research, and future therapies may emerge. For example, temperature-controlled radiofrequency neurolysis has been successfully employed for ablation of the posterior nasal nerve area to produce a significant reduction in symptoms in a group of patients deemed refractory to medication treatment. Relief persisted for 52 weeks, and there were no significant side effects.”

Gerka et al (2021) reported on a prospective single-arm trial of 24 adult patients at seven locations within a large health maintenance organization. Patients with chronic rhinitis that failed medical therapy were offered an in-office cryoablation of posterior nasal nerve (PNN). Patients completed the Total Nasal Symptom Score (TNSS) questionnaire consisting of 5 items reported based on the previous 12 hours and 2 weeks at the following time points: pre-treatment, 30 days, 90 days and 1-year post-treatment. Following cryoablation of the PNN, the TNSS 12-hour symptom score improved from 6.92 (± 2.9) to 3.17 (± 2.4 , $P < 0.001$) at 30 days, 2.92 (± 1.4 , $P < 0.001$) at 90 days and 3.08 (± 2.6 , $P < 0.001$) at 1 year post treatment. Similar results were noted for the 2 weeks scores improving from 7.75 (± 3.1) to 3.79 (± 2.1 , $P < 0.001$) at 30 days, 3.88 (± 1.9 , $P < 0.001$) at 90 days and 3.76 (± 2.1 , $P < 0.001$) at 1-year post-treatment. 64.7% of respondents stated the procedure decreased or eliminated nasal sprays. The authors note that their study has several limitations. Primary limitations of the study include small sample size and short follow up time. However, this independent non-sponsored evaluation is consistent with prior sponsored studies showing sustained benefit post-procedure for up to a 1 year. The rhinitis type subset analysis is limited by the relatively few pure allergic rhinitis patients. Future studies will be needed to demonstrate longer term improvement and sustained results without use of additional medications. The authors concluded that their independent evaluation of cryoablation of the PNN shows improvement in nasal symptoms over a 1-year period and is consistent with other published data.

Stolovitzky et al (2021) conducted a prospective, multicenter, single-blinded randomized controlled trial with a sham procedure control arm to determine the safety and efficacy of temperature-controlled radiofrequency (RF) neurolysis of the posterior nasal nerve (PNN) area for the treatment of chronic rhinitis. This trial was industry sponsored. Patients with 24-hour reflective Total Nasal Symptom Score (rTNSS) ≥ 6 , including moderate to severe rhinorrhea and mild to severe congestion, were randomized 2:1 to active treatment of the posterior nasal nerve area with a temperature-controlled RF device or a sham procedure, with no RF energy delivery. The stylus was applied bilaterally to nonoverlapping areas of the posterior middle meatus and posterior inferior turbinate in each nostril in the region of the PNN. The primary endpoint was responder rate at 3 months, where a response was defined as $\geq 30\%$ improvement (decrease) in rTNSS from baseline. Patients had a mean baseline rTNSS of 8.3 (95% CI, 7.9-8.7) and 8.2 (95% CI, 7.6-8.8) ($P = .797$) in the active treatment ($n = 77$) and sham control ($n = 39$) arms, respectively. At 3 months, responder rate was significantly higher in the active treatment arm: 67.5% (95% CI, 55.9%-77.8%) vs 41.0% (95% CI, 25.6%-57.9%) ($P = .009$). The active treatment arm had a significantly greater decrease in rTNSS (mean, -3.6 [95% CI, -4.2 to -3.0] vs -2.2 [95% CI, -3.2 to -1.3]) ($P = .013$). Three adverse events related to the device/procedure were reported, and all resolved. The authors concluded that this randomized controlled trial showed temperature-controlled neurolysis of the PNN area is free from significant adverse events and superior to a sham procedure in decreasing the symptom burden of chronic rhinitis. Longer-term follow-up is needed.

In an industry sponsored prospective, multicenter, patient-blinded randomized controlled trial (RCT) by Takashima and colleagues (2024), patients in the index active treatment arm were unblinded at 3 months and followed through 12 months. At 3 months, eligible patients from the sham-control arm of the study were invited to crossover to active treatment. Eligibility criteria included reflective total nasal symptom score ((rTNSS) ≥ 6 , with moderate-severe rhinorrhea and mild-severe congestion. The temperature-controlled radiofrequency (TCRF) stylus was applied bilaterally to nonoverlapping areas in the region of the PNN. Patients in the index active treatment arm ($n = 77$) had a mean baseline rTNSS of 8.3 (95% confidence interval [CI], 7.9-8.7). At 12 months, the responder rate was 80.6% ($n = 67$) (95% CI, 69.1%-89.2%). At 12 months, the mean change in rTNSS was -4.8 (95% CI, -5.5 to -4.1; $p < 0.001$), a 57.8% improvement. The available initial rTNSS-based outcomes in the crossover active treatment arm ($n = 27$) were following the same course as the index treatment arm. No serious adverse events and 8 adverse events related to the device/procedure were reported in the trial to date. There were several limitations noted by the authors. The investigators were not blinded in the initial stages of the trial. Medication use was not controlled and could potentially have had some confounding effect on symptom relief, as measured by the rTNSS. The authors concluded that the results of this RCT after the primary endpoint

and patient unblinding demonstrate that the treatment effect of TCRF neurolysis of the posterior nasal nerve (PNN) area is safe and effective in reducing the symptom burden of chronic rhinitis patients through 12 months post-procedure. To date, eligible patients in the index sham-control arm who elected to cross over to active treatment have continued to exhibit the same course of symptom improvement as the index active treatment arm. The combined active treatment group will be followed through 2 years in this trial to demonstrate the durability of this effect.

Lee and colleagues (2022) reported on an industry sponsored prospective single-arm study of 129 patients with chronic rhinitis at 16 medical centers in the United States and Germany to determine clinical outcomes and quality of life (QoL) following temperature-controlled radiofrequency (TCRF) neurolysis of the posterior nasal nerve (PNN) (RhinAer). The mean 24-h reflective total nasal symptom score (rTNSS) improved from 7.8 (95% CI, 7.5-8.1) at baseline to 3.6 (95% CI, 3.2-4.0) at 3 months and continued to improve to 2.9 (95% CI, 2.5-3.3) at 6 months ($p < .001$ comparing follow-up to baseline and $p = .002$ comparing 3 and 6 months). This represents 53.8% improvement over baseline at 3 months and 62.8% improvement at 6 months. Rhinorrhea, congestion, sneezing, and itching subscores and postnasal drip and cough scores were all significantly improved over baseline at both timepoints. At 3 months, 76.2% (95% CI, 68.1%-82.8%) of patients achieved a minimal clinically important difference of $\geq 30\%$ improvement in rTNSS over baseline and the percentage was higher at 6 months (83.5% [95% CI, 75.8%-89.0%]). At 3 months, 80.3% (95% CI, 72.6%-86.3%) reported a minimal clinically important difference of ≥ 0.4 -point improvement in the mini rhinoconjunctivitis quality of life questionnaire score, and the percentage was higher at 6 months; 87.7% (95% CI, 80.7%-92.4%). There were no serious adverse events with a relationship to the device/procedure reported through 6 months. There were several limitations noted by the authors. The lack of a control arm and limited follow-up to date. The authors concluded that TCRF neurolysis of the PNN was safe and resulted in a significant reduction in rhinitis symptom burden at 3 months that was sustained/improved through 6 months. The majority of patients reported a clinically relevant improvement in QoL at 3- and 6-months post-procedure.

POSITION STATEMENT:

Surgical ablation of the posterior nasal nerve by radiofrequency, cryoablation or laser ablation for treatment of chronic rhinitis and all other indications is considered **experimental or investigational**. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

BILLING/CODING INFORMATION:

CPT Coding:

31242	Nasal/sinus endoscopy, surgical; with destruction by radiofrequency ablation, posterior nasal nerve (investigational)
31243	Nasal/sinus endoscopy, surgical; with destruction by cryoablation, posterior nasal nerve (investigational)

REIMBURSEMENT INFORMATION:

Refer to section entitled [POSITION STATEMENT](#).

PROGRAM EXCEPTIONS:

Federal Employee Program (FEP): Follow FEP guidelines.

State Account Organization (SAO): Follow SAO guidelines.

Medicare Advantage products: No National Coverage Determination (NCD) and/or Local Coverage Determination (LCD) were found at the time of the last guideline reviewed date.

If this Medical Coverage Guideline contains a step therapy requirement, in compliance with Florida law 627.42393, members or providers may request a step therapy protocol exemption to this requirement if based on medical necessity. The process for requesting a protocol exemption can be found at [Coverage Protocol Exemption Request](#)

DEFINITIONS:

No guideline specific definitions apply.

RELATED GUIDELINES:

No guideline specific definitions apply.

OTHER:

None applicable.

REFERENCES:

1. American Academy of Otolaryngology Head and Neck Surgery (AAO HNS) Position Statement: Posterior Nasal Nerve (PNN) Ablation for the Treatment of Chronic Rhinitis, January 17, 2023.
2. American Academy of Allergy, Asthma, and Immunology International Consensus Statement on Allergy and Rhinology stated the following for cryotherapy/radiofrequency ablation of posterior nasal nerve, 2023.
3. American Rhinologic Society (ARS) Position Statement: Posterior Nasal Nerve Ablation, January 2022.
4. Blue Cross Blue Shield Association Evidence Positioning System®. 7.01.168 Cryoablation, Radiofrequency Ablation, and Laser Ablation for Treatment of Chronic Rhinitis, 03/25.
5. Chang MT, Song S, Hwang PH. Cryosurgical ablation for treatment of rhinitis: A prospective multicenter study. *Laryngoscope*. 2020 Aug;130(8):1877-1884.
6. DeI Signore AG, Greene JB, Russell JL, et al. Cryotherapy for treatment of chronic rhinitis: 3-month outcomes of a randomized, sham-controlled trial. *Int Forum Allergy Rhinol*. 2022 Jan;12(1):51-61.
7. Ehmer D, McDuffie CM, McIntyre JB, et al. Long-term Outcomes Following Temperature-Controlled Radiofrequency Neurolysis for the Treatment of Chronic Rhinitis. *Allergy Rhinol (Providence)*. 2022 May 29;13: 21526575221096045.
8. Ehmer D, McDuffie CM, Scurry WC Jr, et al. Temperature-Controlled Radiofrequency Neurolysis for the Treatment of Rhinitis. *Am J Rhinol Allergy*. 2022 Jan;36(1):149-156.
9. Gerka Stuyt JA, Luk L, Keschner D, et al. Evaluation of In-Office Cryoablation of Posterior Nasal Nerves for the Treatment of Rhinitis. *Allergy Rhinol (Providence)*. 2021 Jan 29;12: 2152656720988565.
10. Hwang PH, Lin B, Weiss R, et al. Cryosurgical posterior nasal tissue ablation for the treatment of rhinitis. *Int Forum Allergy Rhinol*. 2017 Oct;7(10):952-956.
11. Krespi YP, Wilson KA, Kizhner V. Laser ablation of posterior nasal nerves for rhinitis. *Am J Otolaryngol*. 2020 May-Jun;41(3):102396. [Abstract]
12. Lee JT, Abbas GM, Charous DD, et al. Two-Year Outcomes After Radiofrequency Neurolysis of Posterior Nasal Nerve in Chronic Rhinitis. *Laryngoscope*. 2024 May;134(5):2077-2084.

13. Lee JT, Abbas GM, Charous DD, et al. Clinical and Quality of Life Outcomes Following Temperature-Controlled Radiofrequency Neurolysis of the Posterior Nasal Nerve (RhinAer) for Treatment of Chronic Rhinitis. *Am J Rhinol Allergy*. 2022 Nov; 36(6):747-754.
14. Lieberman PL. Chronic nonallergic rhinitis. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA. March 2025 (Accessed on April 10, 2025.)
15. Ow RA, O'Malley EM, Han JK, Lam KK, Yen DM. Cryosurgical Ablation for Treatment of Rhinitis: Two-Year Results of a Prospective Multicenter Study. *Laryngoscope*. 2021 Sep;131(9):1952-1957.
16. Stolovitzky JP, Ow RA, Silvers SL, et al. Effect of Radiofrequency Neurolysis on the Symptoms of Chronic Rhinitis: A Randomized Controlled Trial. *OTO Open*. 2021 Sep 10;5(3):2473974X211041124.
17. Takashima M, Stolovitzky JP, Ow RA. et al. Temperature-controlled radiofrequency neurolysis for treatment of chronic rhinitis: 12-month outcomes after treatment in a randomized controlled trial. *Int Forum Allergy Rhinol*. 2023 Feb;13(2):107-115.
18. Takashima M, Stolovitzky JP, Ow RA, et al. Temperature-controlled radiofrequency ablation for the treatment of chronic rhinitis: Two-year outcomes from a prospective multicenter trial. *Int Forum Allergy Rhinol*. 2024 Jul;14(7):1182-1194.
19. Wise SK, Damask C, Roland LT, et al. International consensus statement on allergy and rhinology: Allergic rhinitis - 2023. *Int Forum Allergy Rhinol*. 2023 Apr;13(4):293-859.

COMMITTEE APPROVAL:

This Medical Coverage Guideline (MCG) was approved by the Florida Blue Medical Policy and Coverage Committee on 05/28/26.

GUIDELINE UPDATE INFORMATION:

06/15/25	New Medical Coverage Guideline.
06/15/26	Review; no change to position statement.