

02-61000-23

Original Effective Date: 01/01/01

Reviewed: 12/08/23

Revised: 01/01/24

Subject: Sacral Nerve Neuromodulation/Stimulation

THIS MEDICAL COVERAGE GUIDELINE IS NOT AN AUTHORIZATION, CERTIFICATION, EXPLANATION OF BENEFITS, OR A GUARANTEE OF PAYMENT, NOR DOES IT SUBSTITUTE FOR OR CONSTITUTE MEDICAL ADVICE. ALL MEDICAL DECISIONS ARE SOLELY THE RESPONSIBILITY OF THE PATIENT AND PHYSICIAN. BENEFITS ARE DETERMINED BY THE GROUP CONTRACT, MEMBER BENEFIT BOOKLET, AND/OR INDIVIDUAL SUBSCRIBER CERTIFICATE IN EFFECT AT THE TIME SERVICES WERE RENDERED. THIS MEDICAL COVERAGE GUIDELINE APPLIES TO ALL LINES OF BUSINESS UNLESS OTHERWISE NOTED IN THE PROGRAM EXCEPTIONS SECTION.

| | | | | | |
|------------------------------------|--------------------------------|-------------------------------|------------------------------------|-----------------------------|------------------------------------|
| Position Statement | Billing/Coding | Reimbursement | Program Exceptions | Definitions | Related Guidelines |
| Other | References | Updates | | | |

DESCRIPTION:

Treatment using sacral nerve neuromodulation, also known as indirect sacral nerve stimulation, is one of several alternative modalities for patients with urinary or fecal incontinence (urge incontinence, significant symptoms of urgency-frequency, nonobstructive urinary retention) who have failed behavioral (e.g., prompted voiding) and/or pharmacologic therapies.

The sacral nerve neuromodulation device consists of an implantable pulse generator that delivers controlled electrical impulses. This pulse generator is attached to wire leads that connect to the sacral nerves, most commonly the S3 nerve root. Two external components of the system help control the electrical stimulation. A control magnet, kept by the patient, is used to turn the device on or off. A console programmer is kept by the physician and used to adjust the settings of the pulse generator.

POSITION STATEMENT:

Urinary Incontinence and Non-Obstructive Urinary Retention

A trial period of sacral nerve neuromodulation with either percutaneous nerve stimulation or a temporarily implanted lead **meets the definition of medical necessity** in members who meet **ALL** of the following criteria:

1. There is a diagnosis of at least **ONE** of the following:
 - a. Urge incontinence
 - b. Urgency-frequency syndrome
 - c. Non-obstructive urinary retention

- d. Overactive bladder.
2. There is documented failure or intolerance to at least two conventional conservative therapies (e.g., behavioral training such as bladder training, prompted voiding, or pelvic muscle exercise training, pharmacologic treatment for at least a sufficient duration to fully assess its efficacy, and/or surgical corrective therapy).
3. Incontinence is not related to a neurologic condition **AND**
4. The member is an appropriate surgical candidate.

Permanent implantation of a sacral nerve neuromodulation device **meets the definition of medical necessity** in members who meet **ALL** of the following criteria:

1. **ALL** of the criteria listed above (1-4) are met **AND**
2. A trial stimulation period demonstrates at least 50% improvement in symptoms over a period of at least 48 hours.

Other urinary/voiding applications of sacral nerve neuromodulation are considered **experimental or investigational**, including but not limited to treatment of stress incontinence or urge incontinence due to a neurologic condition (e.g., detrusor hyperreflexia, multiple sclerosis, spinal cord injury, or other types of chronic voiding dysfunction). The evidence is insufficient to determine the effects of the technology on health outcomes.

Fecal Incontinence

A trial period of sacral nerve neuromodulation with either percutaneous nerve stimulation or a temporarily implanted lead **meets the definition of medical necessity** in members who meet **ALL** of the following criteria:

1. There is a diagnosis of chronic fecal incontinence of more than 2 incontinent episodes on average per week for more than 6 months or for more than 12 months after vaginal childbirth.
2. There is documented failure or intolerance to conventional conservative therapy (e.g., dietary modification, the addition of bulking and pharmacologic treatment) for at least a sufficient duration to fully assess its efficacy.
3. The condition is not related to an anorectal malformation (e.g., congenital anorectal malformation; defects of the external anal sphincter over 60 degrees; visible sequelae of pelvic radiation; active anal abscesses and fistulae) or chronic inflammatory bowel disease.
4. The member has not had rectal surgery in the previous 12 months, or in the case of cancer, the member has not had rectal surgery in the past 24 months.
5. Incontinence is not related to a neurologic condition **AND**
6. The member is an appropriate surgical candidate.

Permanent implantation of a sacral nerve neuromodulation device **meets the definition of medical necessity** in members who meet **ALL** of the following criteria:

1. All of the criteria listed above (1-6) above are met **AND**

2. A trial stimulation period demonstrates at least 50% improvement in symptoms over a period of at least 48 hours.

Sacral nerve neuromodulation is considered **experimental or investigational** in the treatment of chronic constipation or chronic pelvic pain. The evidence is insufficient to determine the effects of the technology on health outcomes.

BILLING/CODING INFORMATION:

CPT Coding

| | |
|-------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 64561 | Percutaneous implantation of neurostimulator electrode array; sacral nerve (transforaminal placement) including image guidance, if performed |
| 64581 | Open implantation of neurostimulator electrode array; sacral nerve (transforaminal placement) |
| 64585 | Revision or removal of peripheral neurostimulator electrodes |
| 64590 | Insertion or replacement of peripheral, sacral, or gastric neurostimulator pulse generator or receiver, requiring pocket creation and connection between electrode array and pulse generator or receiver |
| 64595 | Revision or removal of peripheral, sacral, or gastric neurostimulator pulse generator or receiver, with detachable connection to electrode array Revision or removal of implanted peripheral or gastric neurostimulator pulse generator or receiver |
| 95970 | Electronic analysis of implanted neurostimulator pulse generator/transmitter (eg, contact group[s], interleaving, amplitude, pulse width, frequency [Hz], on/off cycling, burst, magnet mode, dose lockout, patient selectable parameters, responsive neurostimulation, detection algorithms, closed loop parameters, and passive parameters) by physician or other qualified health care professional; with brain, cranial nerve, spinal cord, peripheral nerve, or sacral nerve, neurostimulator pulse generator/transmitter, without programming |
| 95971 | Electronic analysis of implanted neurostimulator pulse generator/transmitter (eg, contact group[s], interleaving, amplitude, pulse width, frequency [Hz], on/off cycling, burst, magnet mode, dose lockout, patient selectable parameters, responsive neurostimulation, detection algorithms, closed loop parameters, and passive parameters) by physician or other qualified health care professional; with simple spinal cord or peripheral nerve (eg, sacral nerve) neurostimulator pulse generator/transmitter programming by physician or other qualified health care professional |
| 95972 | Electronic analysis of implanted neurostimulator pulse generator/transmitter (eg, contact group[s], interleaving, amplitude, pulse width, frequency [Hz], on/off cycling, burst, magnet mode, dose lockout, patient selectable parameters, responsive neurostimulation, detection algorithms, closed loop parameters, and passive parameters) by physician or other qualified health care professional; with complex spinal cord or peripheral nerve (eg, sacral |

| | |
|-------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| | nerve) neurostimulator pulse generator/transmitter programming by physician or other qualified health care professional |
| 0786T | Insertion or replacement of percutaneous electrode array, sacral, with integrated neurostimulator, including imaging guidance, when performed |
| 0787T | Revision or removal of neurostimulator electrode array, sacral, with integrated neurostimulator |
| 0788T | Electronic analysis with simple programming of implanted integrated neurostimulation system (eg, electrode array and receiver), including contact group(s), amplitude, pulse width, frequency (Hz), on/off cycling, burst, dose lockout, patient-selectable parameters, responsive neurostimulation, detection algorithms, closed-loop parameters, and passive parameters, when performed by physician or other qualified health care professional, spinal cord or sacral nerve, 1-3 parameters |
| 0789T | Electronic analysis with complex programming of implanted integrated neurostimulation system (eg, electrode array and receiver), including contact group(s), amplitude, pulse width, frequency (Hz), on/off cycling, burst, dose lockout, patient-selectable parameters, responsive neurostimulation, detection algorithms, closed-loop parameters, and passive parameters, when performed by physician or other qualified health care professional, spinal cord or sacral nerve, 4 or more parameters |

HCPCS Coding

| | |
|-------|---------------------------------------------------------------------------------------------------------------------------------------------------|
| A4290 | Sacral nerve stimulator test lead, each |
| E0745 | Neuromuscular stimulator, electronic shock unit |
| L8679 | Implantable neurostimulator pulse generator, any type |
| L8680 | Implantable neurostimulator electrode, each |
| L8681 | Patient programmer (external) for use with implantable programmable neurostimulator pulse generator, replacement only |
| L8682 | Implantable neurostimulator radiofrequency receiver |
| L8683 | Radiofrequency transmitter (external) for use with implantable neurostimulator radiofrequency receiver |
| L8684 | Radiofrequency transmitter (external) for use with implantable sacral root neurostimulator receiver for bowel and bladder management, replacement |
| L8685 | Implantable neurostimulator pulse generator, single array, rechargeable, includes extension |
| L8686 | Implantable neurostimulator pulse generator, single array, non-rechargeable, includes extension |
| L8687 | Implantable neurostimulator pulse generator, dual array, rechargeable, includes extension |
| L8688 | Implantable neurostimulator pulse generator, dual array, non-rechargeable, includes extension |

ICD-10 Diagnosis Codes That Support Medical Necessity:

| | |
|-------------|--------------------|
| N39.41 | Urge incontinence |
| R15.0-R15.9 | Fecal incontinence |
| R33.0-R33.9 | Retention of urine |

| | |
|--------|--------------------------|
| R35.0 | Frequency of micturition |
| R35.81 | Nocturnal polyuria |
| R35.89 | Other polyuria |

LOINC Codes

The following information may be required documentation to support medical necessity: Physician history and physical, treatment plan, treatment notes including documentation of symptoms, behavior or pharmacologic interventions, and prior test stimulation (if applicable).

| Documentation Table | LOINC Codes | LOINC Time Frame Modifier Code | LOINC Time Frame Modifier Codes Narrative |
|------------------------------------------------------------------------------------------------------------------------|-------------|--------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------|
| Physician history and physical | 28626-0 | 18805-2 | Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim |
| Attending physician visit note/treatment notes including documentation of symptoms | 18733-6 | 18805-2 | Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim. |
| Treatment plan | 18776-5 | 18805-2 | Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim. |
| Current, Discharge, or Administered Medications (i.e., pharmacologic interventions) | 34483-8 | 18805-2 | Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim |
| Neuromuscular electrophysiology studies (i.e. electronic analysis of implanted neurostimulator pulse generator system) | 27897-8 | 18805-2 | Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim |

REIMBURSEMENT INFORMATION:

Refer to sections entitled [POSITION STATEMENT](#).

PROGRAM EXCEPTIONS:

Federal Employee Program (FEP): Follow FEP guidelines.

State Account Organization (SAO): Follow SAO guidelines.

Medicare Advantage Products:

The following National Coverage Determination (NCD) was reviewed on the last guideline reviewed date: Sacral Nerve Stimulation for Urinary Incontinence (230.18) located at [cms.gov](https://www.cms.gov).

If this Medical Coverage Guideline contains a step therapy requirement, in compliance with Florida law 627.42393, members or providers may request a step therapy protocol exemption to this requirement if based on medical necessity. The process for requesting a protocol exemption can be found at [Coverage Protocol Exemption Request](#)

DEFINITIONS:

None applicable.

RELATED GUIDELINES:

[Pelvic Floor Stimulation as a Treatment of Urinary Incontinence, 01-97000-06](#)

[Percutaneous Tibial Nerve Stimulation, 02-64000-01](#)

OTHER:

None applicable.

REFERENCES:

1. Abrams P, Blaivas JG, Fowler CJ, The Role of Neuromodulation in the Management of Urinary Urge Incontinence, BJU International, Vol 91, Issue 4, Pages 355-359, July 2009.
2. Altomare DF, Ratto C, Ganio E, et al, Long-Term Outcomes of Sacral Nerve Stimulation for Fecal Incontinence, Dis Colon Rectum, 2009 January; 52(1): 11-7.
3. American Urological Association (AUA) Guideline. Diagnosis and Treatment of Interstitial Cystitis/Bladder Pain Syndrome, 2011, accessed at auanet.org.
4. American Urological Association (AUA). Guideline on Diagnosis and Treatment of Overactive Bladder, 2014; accessed at auanet.org.
5. American Urological Association (AUA). Position Statement Regarding Sacral Nerve Stimulation for Urinary Incontinence. October 18, 2000, accessed at auanet.org.
6. Blue Cross Blue Shield Association Evidence Positioning System®; 7.01.69 Sacral Nerve Neuromodulation/Stimulation, 05/22.
7. Blue Cross Blue Shield Association Technology Evaluation Center (TEC) Sacral Nerve Stimulation Urge Incontinence. TEC Assessments 1998, Tab 18.
8. Blue Cross Blue Shield Association. Technology Evaluation Center (TEC). Sacral Nerve Stimulation Urinary Urgency/Frequency. TEC Assessments 2000, Tab 7.
9. Centers for Medicare & Medicaid Services (CMS), National Coverage Determination (NCD) for Sacral Nerve Stimulation for Urinary Incontinence (230.18), accessed at [cms.gov](https://www.cms.gov).
10. Coolen RL, Groen J, Blok B. Electrical Stimulation in the Treatment of Bladder Dysfunction: Technology Update. Med Devices (Auckl). 2019 Sep 11;12:337-345. Doi: 10.2147/MDER.S179898. eCollection 2019.

11. Diagnosis and Treatment of Non-Neurogenic Overactive Bladder (OAB) in Adults: an AUA/SUFU Guideline (2019); accessed at auanet.org.
12. El-Gazzaz G, Zutshi M, et al, Sacral Neuromodulation for the Treatment of Fecal Incontinence and Urinary Incontinence in Female Patients: Long-Term Follow-up, *International Journal of Colorectal Disease*, Vol 24, Number 12, December 2009.
13. Groenendijk IM, Groen J, et al. Acute Effect of Sacral Neuromodulation for Treatment of Detrusor Overactivity on Urodynamic Parameters. *Neurourol Urodyn*. 2020 Feb;39(2):695-701. Doi: 10.1002/nau.24252. Epub 2019 Dec 5. PMID: 31804759.
14. Groenendijk PM, et al, Urodynamic Evaluation of Sacral Neuromodulation for Urge Urinary Incontinence, *BJU Int*. 2008 February; 101(3): 325-9.
15. Herbison GP, Arnold EP, Sacral Neuromodulation with Implanted Devices for Urinary Storage and Voiding Dysfunction in Adults, National Institutes of Health, April 15, 2009.
16. Kenefick NJ. Sacral nerve neuromodulation for the treatment of lower bowel motility disorders. *Ann R Coll Surg Engl*. 2006 Nov; 88(7): 617-23.
17. Leroi AM, Parc Y, Lehur PA, Mion F, Barth X, Rullier E, Bresler L, Portier G, Michot F; Study Group. Efficacy of sacral nerve stimulation for fecal incontinence: results of a multicenter double-blind crossover study. *Ann Surg*. 2005 Nov; 242(5): 662-9.
18. Lo CW, Wu MY, et al. Comparing the Efficacy of OnabotulinumtoxinA, Sacral Neuromodulation, and Peripheral Tibial Nerve Stimulation as Third Line Treatment for the Management of Overactive Bladder Symptoms in Adults: Systematic Review and Network Meta-Analysis. *Toxins (Basel)*. 2020 Feb 18;12(2):128. Doi: 10.3390/toxins12020128. PMID: 32085542.
19. Moore CK, Rueb JJ, et al. What Is New in Neuromodulation? *Curr Urol Rep*. 2019 Aug 7;20(9):55. Doi: 10.1007/s11934-019-0920-6. PMID: 31388779.
20. Mowatt G, Glazener C, Jarrett M. Sacral nerve stimulation for faecal incontinence and constipation in adults. *Cochrane Database of Systematic Reviews* 2007, Issue 3. Art. No.: CD004464. DOI: 10.1002/14651858.CD004464.pub2.
21. National Collaborating Centre for Women's and Children's Health, Urinary Incontinence: The Management of Urinary Incontinence in Women, London (UK), Royal College of Obstetricians and Gynaecologists (RCOG), October 2006, accessed at guideline.gov. 04/16/10.
22. National Institute for Clinical Excellence (NICE), Faecal Incontinence (QS54) Quality Standards, Issued February 2014. Accessed at nice.org.uk.
23. National Institute for Clinical Excellence (NICE). Faecal Incontinence: The Management of Faecal Incontinence in Adults; NICE Clinical Guideline 49. 2007; accessed at nice.org.uk.
24. Oerlemans, D, Van Kerrebroeck P, Sacral Nerve Stimulation for Neuromodulation of the Lower Urinary Tract, *Neurourology and Urodynamics*, Vol 27 Issue 1, pages 28-33, 2008.
25. Paquette IM, Varma M, et al. The American Society of Colon and Rectal Surgeons' Clinical Practice Guideline for the Evaluation and Management of Constipation. *Dis Colon Rectum*. Jun 2016; 59(6): 479-92.
26. Paquette IM, Varma MG, et al. The American Society of Colon and Rectal Surgeons' Clinical Practice Guideline for the Treatment of Fecal Incontinence. *Dis Colon Rectum*. Jul 2015; 58(7): 623-36.
27. Pilkington SA, Emmett C, Knowles CH, et al. Surgery for constipation: systematic review and practice recommendations: Results V: Sacral Nerve Stimulation. *Colorectal Dis*. Sep 2017;19(Suppl 3):92-100.
28. Rao SS, American College of Gastroenterology Practice Guidelines-Diagnosis and Management of Fecal Incontinence, 2004; accessed at s3.gi.org/physicians/guidelines/FecalIncontinence.pdf 08/26/15.

29. Roth TJ, Vandersteen DR, Hollatz P, Sacral Neuromodulation for the Dysfunctional Elimination Syndrome: A Single Center Experience with 20 Children, The Journal of Urology, Vol 180, Issue 1, pages 306-311, July 2008.
30. Satish SC, Practice Guidelines- Diagnosis and Management of Fecal Incontinence, American Journal of Gastroenterology, 2004, 1585-1604.
31. Southwell BR. Electro-Neuromodulation for Colonic Disorders-Review of Meta-Analyses, Systematic Reviews, and RCTs. Neuromodulation. 2020 Feb 3. Doi: 10.1111/ner.13099. Online ahead of print. PMID: 32017319.
32. Tjandra JJ, et al, Practice Parameters for the Treatment of Fecal Incontinence, Dis colon Rectum 2007; 50: 1497-1507, accessed at fascrs.org 04/23/14.
33. White, WM, Dobmeyer-Dittrich C, Klein FA, Wallace LS, Sacral Nerve Stimulation for Treatment of Refractory Urinary Retention: Long-Term Efficacy and Durability, Urology, Vol 71, Issue 1, Pages 71-74, January 2008.

COMMITTEE APPROVAL:

This Medical Coverage Guideline (MCG) was approved by the Florida Blue Medical Policy and Coverage Committee on 12/08/23.

GUIDELINE UPDATE INFORMATION:

| | |
|----------|---------------------------------------------------------------------------------------------------------------------------------------------------|
| 01/01/01 | New Medical Coverage Guideline. |
| 01/01/02 | Annual HCPCS coding update. |
| 07/25/02 | Reviewed. |
| 08/15/03 | Reviewed; no changes in coverage statement MCG changed to Active but no longer scheduled for routine review. |
| 01/01/05 | Annual HCPCS coding update: consisting of the revision of 64590, 95970, 95971, 95972 and 95973. |
| 01/01/06 | Annual HCPCS coding update: consisting of the deletion of E0752, E0754, E0756 and E0759 and the addition of L8680, L8681, L8682, L8683 and L8684. |
| 01/01/07 | Annual HCPCS coding update: consisting of the revision of 64590 and 64595. |
| 09/15/07 | Review and revision of guideline consisting of updated references and reformatted guideline. |
| 09/15/08 | Review and revision of guideline consisting of updated references. |
| 01/01/09 | Annual HCPCS coding update: revised descriptor for code L8681. |
| 07/15/09 | Annual review: position statements maintained, coding and references updated. |
| 06/15/10 | Annual review: position statements maintained and references updated. |
| 10/15/10 | Revision: formatting changes and related ICD-10 codes added. |
| 08/15/11 | Revision; formatting changes. |
| 10/01/11 | Revision; formatting changes. |
| 01/01/12 | Annual HCPCS update. Revised descriptor for codes 64561, 64581, & 95970-95973. |
| 05/15/12 | Annual review; title, position statements, coding/billing section, and references updated; formatting changes. |
| 10/15/12 | Permanent implantation criteria updated; formatting changes. |
| 01/01/13 | Annual HCPCS update. Revised descriptor for code 64561. |

| | |
|----------|--------------------------------------------------------------------------------------------------------------------------|
| 06/15/13 | Annual review; position statement section and references updated; formatting changes. |
| 01/01/14 | Annual HCPCS update. Added code L8679. |
| 06/15/14 | Annual review; Update position statements, coding, and references; formatting changes. |
| 01/01/15 | Annual HCPCS/CPT update. Revised code 95972. |
| 10/15/15 | Annual review; position statements, coding, & references updated; formatting changes. |
| 01/01/16 | Annual HCPCS/CPT update; code 95972 revised, code 95973 deleted. |
| 01/01/17 | Annual CPT/HCPCS update. Revised 95972; formatting changes. |
| 04/15/17 | Revision; position statements maintained, description section and references updated. |
| 07/15/18 | Review; description, position statements, coding, and references updated. |
| 01/01/19 | Annual CPT/HCPCS coding update. Revised codes 95970-95972. |
| 07/15/20 | Review; position statements maintained and references updated. |
| 01/01/22 | Annual CPT/HCPCS coding update. Code 64581 revised. |
| 06/15/22 | Review: Position statements maintained; coding and references updated. |
| 05/25/23 | Update to Program Exceptions section. |
| 01/01/24 | Position statements maintained. Annual CPT/HCPCS coding update. Codes 0786T- 0789T added; codes 64590, 64595 revised. |