02-61000-34

Original Effective Date: 11/15/07

Reviewed: 03/27/25

Revised: 04/15/25

Subject: Neurolysis/Ablation

THIS MEDICAL COVERAGE GUIDELINE IS NOT AN AUTHORIZATION, CERTIFICATION, EXPLANATION OF BENEFITS, OR A GUARANTEE OF PAYMENT, NOR DOES IT SUBSTITUTE FOR OR CONSTITUTE MEDICAL ADVICE. ALL MEDICAL DECISIONS ARE SOLELY THE RESPONSIBILITY OF THE PATIENT AND PHYSICIAN. BENEFITS ARE DETERMINED BY THE GROUP CONTRACT, MEMBER BENEFIT BOOKLET, AND/OR INDIVIDUAL SUBSCRIBER CERTIFICATE IN EFFECT AT THE TIME SERVICES WERE RENDERED. THIS MEDICAL COVERAGE GUIDELINE APPLIES TO ALL LINES OF BUSINESS UNLESS OTHERWISE NOTED IN THE PROGRAM EXCEPTIONS SECTION.

Position Statement	Billing/Coding	Reimbursement	Program Exceptions	<u>Definitions</u>	Related Guidelines
<u>Other</u>	References	<u>Update</u>			

DESCRIPTION:

Percutaneous radiofrequency (RF) facet neurolysis/ablation/denervation is used to treat neck and back pain originating in facet joints with degenerative changes. Diagnosis of facet joint pain is confirmed by response to nerve blocks. The goal of facet denervation is long-term pain relief. However, the nerves regenerate and, therefore, repeat procedures may be required.

Trigeminal neuralgia is one of the most common forms of severe facial pain. It is generally caused by neurovascular compression. Well established TN treatment protocols for trigeminal neuralgia include pharmacotherapy, neurotoxin denervation, nerve ablation and microvascular decompression, with high rates of relief.

Summary and Analysis of Evidence: UpToDate review "Interventional therapies for chronic pain" (Copenhaver et al, 2025) states, "the most common indications for radiofrequency neurotomy/radiofrequency ablation (RFN) are axial neck or back pain when the facet joints have been identified as the etiology of pain (eg, non-radicular axial spine pain with possible paraspinal tenderness in the location of the facet joints) ... efficacy of neurolytic procedures and the potential for adverse effects vary based on the speed with which the individual's nerves regenerate and whether regeneration results in normal neural function, acute neuritis, or chronic aberrant sensory changes. Various studies, including randomized controlled trials, have found that RFA procedures may be highly effective for pain relief. As an example, reported rates of positive outcome (ie, >50 percent pain relief at more than three months) are seen in more than 65 percent in patients who had effective diagnostic blocks prior to RFA. If pain recurs, repeat RFA can be performed. Studies have found that repeat medial branch RFA was successful in up to 65 to 90 percent of patients after an initially successful block."

The American Society of Interventional Pain Physicians (ASIPP) 2020 consensus guideline for facet joint interventions (Manchikanti et al) states that for the cervical spine, "the level of evidence is II with moderate strength of recommendation for cervical radiofrequency ablation with the inclusion of one randomized controlled trial with positive results and two observational studies with long-term improvement". For the thoracic spine, the guideline states "the level of evidence is III with weak to moderate strength of recommendation with emerging evidence for thoracic radiofrequency ablation with the inclusion of one relevant randomized controlled trial and three observational studies." For the lumbar

spine, the guideline states "The level of evidence is II with moderate strength of recommendation for lumbar radiofrequency ablation with inclusion of 11 relevant randomized controlled trials (RCTs) with 2 negative studies and 4 studies with long-term improvement."

UpToDate review "Trigeminal neuralgia" (Ho et al, 2025) states, "percutaneous ablation procedures may be preferred when imaging shows no neurovascular contact ... other ablative strategies for refractory idiopathic TN include radiosurgery with gamma knife or rhizotomy with radiofrequency, balloon, or glycerol ... patients with medically refractory TN symptoms who are felt to be at a high surgical risk due to concomitant medical comorbidities may be candidates for percutaneous ablation procedures like gamma knife stereotactic radiosurgery or radiofrequency thermocoagulation rhizotomy."

Texakalidis et al (2019) conducted a meta-analysis of 14 studies that covered more than 2,500 participants, comparing clinical efficacy of three treatments for trigeminal neuralgia: radiofrequency denervation, glycerol rhizotomy, and balloon compression. Radiofrequency denervation was associated with higher likelihood of distribution of trigeminal anesthesia and immediate pain relief. The authors concluded that radiofrequency denervation is as good as, or better than, other widely used treatments for trigeminal neuralgia.

There is insufficient published clinical evidence to support the safety and effectiveness of the following denervation interventions:

- Destruction of peripheral nerves of the extremities (Arnold et al, 2024; Finneran et al, 2020;
 McLean et al, 2020)
- Destruction of a nerve plexus or other truncal nerve (eg, brachial plexus, pudendal nerve) (Hampton et al, 2023; Shah et al, 2021)
- Destruction of the intercostal nerve (Eldredge et al, 2023; Gabriel et al, 2020; Humble 2015)
- Destruction of the genicular nerve branches (UpToDate, "Investigational approaches to the management of osteoarthritis", Yu, 2025); Liu et al, 2022; American Academy of Orthopedic Surgeons, 2021; Chen, Mullen et al, 2021)
- Destruction of the nerves innervating the sacroiliac joint (UpToDate "Subacute and chronic low back pain: Nonsurgical interventional treatment", Chou, 2025); Chou et al, 2021; Chappell et al, 2020; Chen et al, 2019; Juch et al, 2017; van Tilburg et al, 2016)
- Destruction of the pudendal nerve (Conic et al, 2025; Collard et al, 2021; Petrov-Kondratov et al, 2017)
- Destruction of the celiac plexus (Kong et al, 2022; Galafassi et al, 2020)
- Destruction of the superior hypogastric plexus (Bagchi et al, 2022; Fisher et al, 2021'; Rocha et al, 2020)

POSITION STATEMENT:

Paravertebral facet joint neurolysis/ablation

Facet joint pain

Non-pulsed radiofrequency neurolysis (ablation) of the paravertebral facet joints for pain suggestive of facet joint origin **meets the definition of medical necessity** when all of the following are met:

No prior spinal fusion surgery in the vertebral level being treated, AND

- Disabling back pain suggestive of facet joint origin, AND
- No nerve root compression documented in the medical record or on radiographic evaluations,
 AND
- The pain is not radicular, AND
- Pain has failed to respond to 3 months of conservative management, which may consist of therapies such as nonsteroidal anti-inflammatory medications, acetaminophen, manipulation, physical therapy, and a home exercise program, AND
- There has been a successful trial of controlled medial branch blocks, consisting of:
 - o 2 separate positive blocks on 2 different days, **OR**
 - A placebo-controlled series of blocks, under fluoroscopic guidance, that has resulted in at least a 50% reduction in pain for the duration of the local anesthetic used, AND
- If there has been a prior successful radiofrequency denervation, a minimum time of 6 months has elapsed since prior radiofrequency treatment*

***NOTE:** If there has been a prior successful radiofrequency denervation, additional diagnostic medial branch blocks for the same level of the spine are not required.

Trigeminal nerve neurolysis/ablation

Non-pulsed radiofrequency neurolysis (ablation) of the trigeminal nerve **meets the definition of medical necessity** when all of the following are met:

- Member is diagnosed with trigeminal neuralgia, AND
- Failure of at least 3 months of conservative treatment with medications, **OR** inability to tolerate side effects of the medications

All other methods of facet neurolysis and trigeminal nerve neurolysis are considered **experimental or investigational**, including but not limited to pulsed radiofrequency neurolysis, cooled radiofrequency neurolysis, laser neurolysis, chemical neurolysis and cryoneurolysis.

Radiofrequency neurolysis (all types), laser neurolysis, and cryoneurolysis are considered **experimental or investigational** for all other nerves, joints, and conditions, including but not limited to pain associated with the sacroiliac joint (SI) joints and osteoarthritis. The available scientific evidence does not support conclusions regarding safety, effectiveness, and net outcomes.

BILLING/CODING INFORMATION:

The following codes may be used to describe neurolysis:

NOTE: Per CPT coding guidelines, codes 64633, 64634, 64635, and 64636 exclude pulsed radiofrequency methods.

CPT Coding

0440T	Ablation, percutaneous, cryoablation, includes imaging guidance; upper extremity
	distal/peripheral nerve (investigational)
0441T	Ablation, percutaneous, cryoablation, includes imaging guidance; lower extremity
	distal/peripheral nerve (investigational)
0442T	Ablation, percutaneous, cryoablation, includes imaging guidance; nerve plexus or other
	truncal nerve (eg, brachial plexus, pudendal nerve) (investigational)
64600	Destruction by neurolytic agent, trigeminal nerve; supraorbital, infraorbital, mental, or
	inferior alveolar branch
64605	Destruction by neurolytic agent, trigeminal nerve; second and third division branches at
	foramen ovale
64610	Destruction by neurolytic agent, trigeminal nerve; second and third division branches at
	foramen ovale under radiologic monitoring
64620	Destruction by neurolytic agent, intercostal nerve (investigational)
64624	Destruction by neurolytic agent, genicular nerve branches including imaging guidance,
	when performed (investigational)
64625	Radiofrequency ablation, nerves innervating the sacroiliac joint, with image guidance (ie,
	fluoroscopy or computed tomography) (investigational)
64630	Destruction by neurolytic agent; pudendal nerve (investigational)
64633	Destruction by neurolytic agent, paravertebral facet joint nerve(s) with imaging guidance
	(fluoroscopy or CT); cervical or thoracic, single facet joint
64634	Destruction by neurolytic agent, paravertebral facet joint nerve(s) with imaging guidance
	(fluoroscopy or CT); cervical or thoracic, each additional facet joint (list separately in
	addition to code for primary procedure)
64635	Destruction by neurolytic agent, paravertebral facet joint nerve(s) with imaging guidance
	(fluoroscopy or CT); lumbar or sacral, single facet joint
64636	Destruction by neurolytic agent, paravertebral facet joint nerve(s) with imaging guidance
	(fluoroscopy or CT); lumbar or sacral, each additional facet joint (list separately in addition
	to code for primary procedure)
64640	Destruction by neurolytic agent, other peripheral nerve or branch
64680	Destruction by neurolytic agent, with or without radiologic monitoring; celiac plexus
	(investigational)
64681	Destruction by neurolytic agent, with or without radiologic monitoring; superior
	hypogastric plexus (investigational)

HCPCS Coding

C9809	Cryoablation needle (e.g., iovera system), including needle/tip and all disposable system
	components, non-opioid medical device (must be a qualifying medicare non-opioid
	medical device for post-surgical pain relief in accordance with section 4135 of the caa,
	2023) (Investigational)

ICD-10 Diagnosis Codes That Support Medical Necessity (64633, 64634, 64635, 64636)

M47.011 – M47.014	Anterior spinal artery compression syndromes, occipital, cervical,
	cervicothoracic, thoracic regions
M47.015 M47.016	Anterior spinal artery compression syndromes, thoracolumbar, lumbar
	regions
M47.021 – M47.022	Vertebral artery compression syndromes, occipital and cervical regions
M47.11 – M47.14	Other spondylosis with myelopathy, occipital, cervical, cervicothoracic,
	thoracic regions
M47.15 – M47.16	Other spondylosis with myelopathy, thoracolumbar, lumbar regions
M47.21 – M47.24	Other spondylosis with radiculopathy, occipital, cervical, cervicothoracic,
	thoracic regions
M47.25 – M47.28	Other spondylosis with radiculopathy, thoracolumbar, lumbar,
	lumbosacral, sacral and sacrococcygeal regions
M47.811 – M47.814	Spondylosis without myelopathy or radiculopathy, occipital, cervical,
	cevicothoracic, thoracic regions
M47.815 M47.818	Spondylosis without myelopathy or radiculopathy, thoracolumbar,
	lumbar, lumbosacral, sacral and sacrococcygeal regions
M47.891 – M47.894	Other spondylosis, occipital, cervical, cervicothoracic, thoracic regions
M47.895 M47.898	Other spondylosis, thoracolumbar, lumbar, lumbosacral, sacral and
	sacrococcygeal regions
M54.2	Cervicalgia
M54.31 – M54.32	Sciatica
M54.40 – M54.42	Lumbago with sciatica
M54.50, M54.51,	Low back pain, including vertebrogenic low back pain
M54.59	
M54.6	Pain in thoracic spine
M96.1	Post-laminectomy syndrome, not elsewhere classified

ICD-10 Diagnosis Codes That Support Medical Necessity for trigeminal neuralgia (64400, 64405, 64410)

G50.0, G50.8, G50.9	Disorders of trigeminal nerve
---------------------	-------------------------------

REIMBURSEMENT INFORMATION:

Percutaneous non-pulsed radiofrequency neurolysis for facet joint pain is limited to two (2) facet neurolysis procedures per region, every 12 months, with code limitations as follows:

64633 and **64634** in any combination, up to 12 in 12 months.

64635 and 64636 in any combination, up to 12 in 12 months.

Percutaneous non-pulsed radiofrequency neurolysis for trigeminal neuralgia is limited to one (1) procedure per branch, per 12 months, with code limitations as follows:

Code 64600 is limited to 1 per 12 months.

Code 64605 is limited to 1 per 12 months.

Code 64610 is limited to 1 per 12 months.

NOTE: Services in excess of the limitations shown above are subject to medical review of documentation. The following information is required documentation to support medical necessity: physician history and physical, radiology study reports, physician progress notes with documentation of conservative treatment, treatment plan including narrative, physician operative report. Documentation must support "Position Statement" criteria and provide rationale for additional procedures.

LOINC Codes:

Documentation Table	LOINC Codes	LOINC Time Frame Modifier Code	LOINC Time Frame Modifier Codes Narrative
Physician history and physical	28626-0	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.
Attending physician progress note	18741-9	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.
Radiology	18726-0	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.
Treatment plan, plan of treatment	18776-5	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.
Physical therapy initial assessment	18735-1	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.
Physical therapy progress note	11508-9	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.
Current, discharge, or administered medications	34483-8	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.
Physician operative report	28573-4	18805-2	Include all data of the selected type that represents observations made six months or

	fewer before starting date of service for the	
	claim.	

PROGRAM EXCEPTIONS:

Federal Employee Program (FEP): Follow FEP guidelines.

State Account Organization (SAO): Follow SAO guidelines.

Medicare Advantage Products: The following National Coverage Determination (NCD) was reviewed on the last guideline review date: Induced Lesions of Nerve Tracts (160.1), located at cms.gov.

The following Local Coverage Determinations (LCD) was reviewed on the last guideline reviewed date: Facet Joint Interventions for Pain Management (L33930), located at cms.gov.

If this Medical Coverage Guideline contains a step therapy requirement, in compliance with Florida law 627.42393, members or providers may request a step therapy protocol exemption to this requirement if based on medical necessity. The process for requesting a protocol exemption can be found at Coverage Protocol Exemption Request.

DEFINITIONS:

Cryoneurolysis: destruction of a nerve with the use of extreme cold; also called cryosurgery, cryoablation, cryodenervation (e.g., iovera treatment)

Facet joint: each of four joints formed above and below and on either side of a vertebra by bony projections (articular processes). The smooth surface at the end of the bony projections is called a facet. Each vertebra has a bony projection on either side which angles downward on its lower side and a bony projection that angles upward on either side. The lower projections of one vertebra meet the upper projections of the vertebra below it, forming facet joints.

Genicular nerve: a sensory nerve that surrounds the knee and provides innervation for the joint.

Laser ablation: destruction of a nerve with a powerful beam of light that produces intense heat when focused at close range.

Medial branch block: injection of local anesthetic near the very small nerve branches that control sensation to the facet joints.

Neurolysis, ablation, denervation, rhizotomy, neurotomy, or neuroablation: destruction of a nerve.

Radiofrequency neurolysis (radiofrequency lesioning): destruction of a nerve with the use of heat.

RELATED GUIDELINES:

Nerve Block Injections, 02-61000-29

Facet Joint Injections, 02-61000-30

OTHER:

None applicable.

REFERENCES:

- AHRQ National Guideline Clearinghouse. Guideline Summary NGC-7269. Low back pain. Early management of persistent non-specific low back pain. National Institute for Health and Clinical Excellence (NICE); 2009 May.
- 2. AHRQ National Guideline Clearinghouse. Guideline Summary NGC-7479. Diagnosis and treatment of forefoot disorders. Section 3. Morton's intermetatarsal neuroma. Clinical Practice Guideline Forefoot Disorders Panel. March 2009.
- AHRQ National Guideline Clearinghouse. Guideline Summary NGC-7951. Practice guidelines for chronic pain management. An updated report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. Anesthesiology 2010 Apr;112(4):810-33.
- 4. AHRQ National Guideline Clearinghouse. Guideline Summary NGC-8744. Management of acute low back pain. Michigan Quality Improvement Consortium. September 2011.
- AHRQ National Guideline Clearinghouse. Guideline Summary NGC-9259. Guideline for the evidence-informed primary care management of low back pain. Toward Optimized Practice; 2011.
- American Academy of Family Physicians. Plantar Fasciitis: A Common Cause of Heel Pain. ©2000
 2010.
- American Academy of Orthopaedic Surgeons Management of Osteoarthritis of the Knee (NonArthroplasty) Evidence-Based Clinical Practice Guideline. https://www.aaos.org/oak3cpg Published 08/31/2021.
- 8. American Chronic Pain Association. ACPA Consumer Guide to Pain Medication & Treatment 2011. Epidural, Nerve & Facet Blocks & Radiofreguency Ablation (Rhizotomy). (Accessed on 02/21/11).
- 9. American Podiatric Medical Association. Treatment by Your Podiatric Physician: Neuromas. ©2010 American Podiatric Medical Association, Inc.
- 10. American Podiatric Medical Association. Treatment by Your Podiatric Physician: Heel Pain. ©2010 American Podiatric Medical Association, Inc.
- 11. American Podiatric Medical Association. What is a neuroma? Accessed at http://www.apma.org/ on 02/05/13.
- 12. American Society of Anesthesiologists Task Force on Chronic Pain Management, American Society of Regional Anesthesia and Pain Medicine. Practice guidelines for chronic pain management: an updated report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. Anesthesiology 2010 Apr;112(4):810-33.
- Arnold DMJ, Ho ES, Switzer-McIntyre S, Amara C, Agur AMR. Anatomical study of the innervation of the posterior elbow joint capsule: implications for ultrasound-guided peripheral nerve block and radiofrequency ablation procedures. Reg Anesth Pain Med. 2024 Dec 3:rapm-2024-106098. doi: 10.1136/rapm-2024-106098. Epub ahead of print. PMID: 39627023.
- 14. Avanos Medical, Inc. Clinical dossier: COOLIEF* Cooled Thermal Radiofrequency Ablation for the Genicular Nerves of the Knee.
- 15. Bagchi S, Kemp R, Green CJ, Stephenson J. Superior Hypogastric Plexus Neurolysis to Manage Metastatic Pelvic Pain in a Forty-Five-Year-Old Spina Bifida Patient: A Case Report. J Med Cases. 2022 Feb;13(2):51-55. doi: 10.14740/jmc3800. Epub 2022 Feb 16.

- 16. Ball RD. The science of conventional and water-cooled monopolar lumbar radiofrequency rhizotomy: an electrical engineering point of view. Pain Physician. 2014: 17 (2): E175-211.
- 17. Becker S, Hadjipavlou A, Heggeness MH. Ablation of the basivertebral nerve for treatment of back pain: a clinical study. Spine J. 2017;17(2):218–223. doi:10.1016/j.spinee.2016.08.032.
- 18. Blue Cross Blue Shield Association Evidence Positioning System®. 6.01.23 Diagnosis and Treatment of Sacroiliac Joint Pain, 12/24.
- 19. Blue Cross Blue Shield Association Evidence Positioning System®. 7.01.116 -- Facet Joint Denervation, 12/24.
- 20. Blue Cross Blue Shield Association Evidence Positioning System®. 7.01.147 Minimally Invasive Ablation Procedures for Morton and Other Peripheral Neuromas, 08/24.
- 21. Blue Cross Blue Shield Association Evidence Positioning System®. 7.01.154 Ablation of Peripheral Nerves to Treat Pain, 10/24.
- 22. Blue Cross Blue Shield Association Evidence Positioning System®. 2.01.97-- Alcohol Injections for Treatment of Peripheral Neuromas (Sunset Status July 2022; refer to policy 7.01.147).
- 23. Boswell MV, Colson JD, Sehgal N, Dunbar EE, Epter R. A systematic review of therapeutic facet joint interventions in chronic spinal pain. Pain Physician. 2007 Jan; 10(1): 229-53.
- 24. Boswell MV, Trescot AM, Datta S, Schultz DM, Hansen HC, Abdi S, Sehgal N, Shah RV, Singh V, Benyamin RM, Patel VB, Buenaventura RM, Colson JD, Cordner HJ, Epter RS, Jasper JF, Dunbar EE, Atluri SL, Bowman RC, Deer TR, Swicegood JR, Staats PS, Smith HS, Burton AW, Kloth DS, Giordano J, Manchikanti L; American Society of Interventional Pain Physicians. Interventional techniques: evidence-based practice guidelines in the management of chronic spinal pain. Pain Physician. 2007 Jan; 10 (1): 7-111.
- 25. Burnham RS, Holitski S, Dinu I. A Prospective Outcome Study on the Effects of Facet Joint Radiofrequency Denervation on Pain, Analgesic Intake, Disability, Satisfaction, Cost, and Employment. Arch Phys Med Rehabil Vol 90, February 2009.
- 26. Bykowski JL, Wong WHW. Role of Facet Joints in Spine Pain and Image- Guided Treatment: A Review. Am J Neuroradiology 33:1419 –26 Sep 2012.
- 27. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determination (LCD): Destruction of Paravertebral Facet Joint Nerve(s) (L33814) (10/01/15) (Retired 04/25/21).
- 28. Centers for Medicare and Medicaid Services (CMS). Local Coverage Determination (LCD) Facet Joint Interventions for Pain Management (L33930) (10/01/15) (Revised 08/11/24).
- 29. Centers for Medicare and Medicaid Services (CMS). National Coverage Determination (NCD) Induced Lesions of Nerve Tracts (160.1) (Longstanding NCD).
- 30. Chappell ME, Lakshman R, Trotter P, Abrahams M, Lee M. Radiofrequency denervation for chronic back pain: a systematic review and meta-analysis. BMJ Open. 2020 Jul 21;10(7):e035540. doi: 10.1136/bmjopen-2019-035540.
- 31. Chen AF, Mullen K, Casambre F, Visvabharathy V, Brown GA. Thermal Nerve Radiofrequency Ablation for the Nonsurgical Treatment of Knee Osteoarthritis: A Systematic Literature Review. J Am Acad Orthop Surg. 2021 May 1;29(9):387-396. doi: 10.5435/JAAOS-D-20-00522.
- 32. Chen AF, Khalouf F, et al. Cooled Radiofrequency Ablation Compared with a Single Injection of Hyaluronic Acid for Chronic Knee Pain. A Multicenter, Randomized Clinical Trial Demonstrating Greater Efficacy and Equivalent Safety for Cooled Radiofrequency Ablation. J Bone Joint Surg Am. 2020;102:1501-10.
- 33. Chen A, Khalouf F, Zora K et al. Cooled radiofrequency ablation provides extended clinical utility in the management of knee osteoarthritis: 12-month results from a prospective, multi-center, randomized, cross-over trial comparing cooled radiofrequency ablation to a single hyaluronic acid injection. BMC Musculoskeletal Disorders (2020) 21:363.

- 34. Chen AF, Mullen K, Casambre F, Visvabharathy V, Brown GA. Thermal Nerve Radiofrequency Ablation for the Nonsurgical Treatment of Knee Osteoarthritis: A Systematic Literature Review [published online ahead of print, 2020 Jul 22]. J Am Acad Orthop Surg. 2020;10.5435/JAAOS-D-20-00522. doi:10.5435/JAAOS-D-20-00522. PMID: 32701684.
- 35. Chen CH, Weng PW, Wu LC, Chiang YF, Chiang CJ. Radiofrequency neurotomy in chronic lumbar and sacroiliac joint pain: A meta-analysis. Medicine (Baltimore). 2019 Jun;98(26):e16230. doi: 10.1097/MD.0000000000016230.
- 36. Chou R, et al. Diagnosis and Treatment of Low Back Pain: A Joint Clinical Practice Guideline from the American College of Physicians and the American Pain Society. Ann Intern Med. 2007;147:478-491.
- 37. Chou R, Fu R, Dana T, Pappas M, Hart E, Mauer KM. Interventional Treatments for Acute and Chronic Pain: Systematic Review. Comparative Effectiveness Review No. 247. (Prepared by the Pacific Northwest Evidence-based Practice Center under Contract No. 75Q80120D00006.) AHRQ Publication No. 21-EHC030. Rockville, MD: Agency for Healthcare Research and Quality; September 2021. DOI: https://doi.org/10.23970/AHRQEPCCER247.
- 38. Chou R, Huffman LH. American Pain Society Clinical Guideline for the Evaluation and Management of Low Back Pain: Evidence Review. 2009.
- 39. Civelek E, et al. Comparison of Effectiveness of Facet Joint Injection and Radiofrequency Denervation in Chronic Low Back Pain. Turkish Neurosurgery 2012, Vol. 22, No. 2, 200-206.
- 40. ClinicalTrials.gov NCT01743326: RFD Versus Cervical Medial Branch Blocks in Chronic Degenerative Neck Pain. Maastricht University Medical Center. Last updated: December 13, 2012.
- 41. Clinical trials.gov. NCT03630133: Intraosseous Basivertebral Nerve Ablation for the Treatment of Chronic Low Back Pain (CLBP). Relievant Medsystems, Inc. (October 2019).
- 42. Chua NHL, Vissers KC, Sluijter ME. Pulsed radiofrequency treatment in interventional pain management: mechanisms and potential indications—a review. Acta Neurochir (2011) 153:763–771.
- 43. Cohen SP, Bhaskar A, Bhatia A, et al. Consensus practice guidelines on interventions for lumbar facet joint pain from a multispecialty, international working group. Reg Anesth Pain Med. 2020 Jun;45(6):424-467. doi: 10.1136/rapm-2019-101243. Epub 2020 Apr 3.
- 44. Collard MD, Xi Y, Patel AA, Scott KM, Jones S, Chhabra A. Initial experience of CT-guided pulsed radiofrequency ablation of the pudendal nerve for chronic recalcitrant pelvic pain. Clin Radiol. 2019 Nov;74(11):897.e17-897.e23. doi: 10.1016/j.crad.2019.06.028. Epub 2019 Aug 22.
- 45. Conic RRZ, Kaur P, Kohan LR. Pudendal Neuralgia: A Review of the Current Literature. Curr Pain Headache Rep. 2025 Jan 28;29(1):38. doi: 10.1007/s11916-024-01354-z. PMID: 39873912.
- 46. Davis T, Loudermilk E, et al. Prospective, Multicenter, Randomized, Crossover Clinical Trial Comparing the Safety and Effectiveness of Cooled Radiofrequency Ablation with Corticosteroid Injection in the Management of Knee Pain from Osteoarthritis. Reg Anesth Pain Med. 2018; 43(1):84-91.
- 47. Dobrogowski J, Wrzosek A, Wordliczek J. Radiofrequency denervation with or without addition of pentoxifylline or methylprednisolone for chronic lumbar zygapophysial joint pain. Pharmacol Rep. 2005 Jul-Aug; 57(4):475-80.
- 48. ECRI Health Technology Assessment Information Services. Custom Hotline Response. Radiofrequency Neuroablation for Low-Back Pain. Updated 10/20/06.
- 49. ECRI Health Technology Assessment Information Services. Evidence Report. Radiofrequency Ablation for Chronic Spinal Pain. 04/07/10.
- 50. Eldredge RS, McMahon L. Intercostal nerve cryoablation therapy for the repair of pectus excavatum: a systematic review. Front Surg. 2023 Aug 24;10:1235120. doi: 10.3389/fsurg.2023.1235120.

- 51. Erdem Y, Sir E. The Efficacy of Ultrasound-Guided Pulsed Radiofrequency of Genicular Nerves in the Treatment of Chronic Knee Pain Due to Severe Degenerative Disease or Previous Total Knee Arthroplasty. Med Sci Monit. 2019 Mar 12;25:1857-1863. doi: 10.12659/MSM.915359.
- 52. Finneran JJ, Swisher MW, Gabriel RA, Fiallo A, Hamilton JR, Lee JG, Ilfeld BM. Ultrasound-Guided Lateral Femoral Cutaneous Nerve Cryoneurolysis for Analgesia in Patients With Burns. J Burn Care Res. 2020 Jan 30;41(1):224-227. doi: 10.1093/jbcr/irz192.
- 53. Fischgrund JS, Rhyne A, et al. Intraosseous Basivertebral Nerve Ablation for the Treatment of Chronic Low Back Pain: 2-Year Results From a Prospective Randomized Double-Blind Sham-Controlled Multicenter Study. International Journal of Spine Surgery, Vol. 13, No. 2, 2019, pp. 110–119.
- 54. Fischgrund JS, Rhyne A, et al. Intraosseous basivertebral nerve ablation for the treatment of chronic low back pain: a prospective randomized double-blind sham-controlled multi-center study. European Spine Journal. 2018 May 1;27(5):1146-56.
- 55. Fisher K, Daoud J, Gonzalez C, Reyes Md J, Lopez D, Desyatnikov O. Fluoroscopic-Guided Bilateral Superior Hypogastric Plexus Neurolysis in the Treatment of Intractable Neoplasm-Related Penile Pain. Cureus. 2021 Nov 29;13(11):e19991. doi: 10.7759/cureus.19991.
- 56. Florida Medicare Part B Local Coverage Determination. L29132 Destruction of Paravertebral Facet Joint Nerve(s) (Retired 09/30/15).
- 57. Florida Medicare Part B Local Coverage Determination. L29130 Destruction by neurolytic agent; interdigital nerve of the foot Morton's Neuroma. 02/02/09. (Retired 09/30/15).
- 58. Gabriel RA, Finneran JJ, Swisher MW, Said ET, Sztain JF, Khatibi B, Wallace AM, Hosseini A, Trescot AM, Ilfeld BM. Ultrasound-guided percutaneous intercostal cryoanalgesia for multiple weeks of analgesia following mastectomy: a case series. Korean J Anesthesiol. 2020 Apr;73(2):163-168. doi: 10.4097/kja.19332. Epub 2019 Sep 2.
- 59. Galafassi GZ, de Aguiar PH, de Oliveira NM, Rossi TG, de Aguiar PH. Celiac plexus neurolysis: systematical review. Rev. Chil. Neurocirugía. 2020;46:37-44.
- 60. Gazioğlu Türkyılmaz G, Rumeli Ş. The effect of consecutive facet medial branch radiofrequency denervation and dorsal root ganglion pulse radiofrequency therapy on lumbar facet joint pain. Agri. 2023 Oct;35(4):220-227. English. doi: 10.14744/agri.2022.42713. PMID: 37886865.
- 61. Gerwin R. Chronic Facial Pain: Trigeminal Neuralgia, Persistent Idiopathic Facial Pain, and Myofascial Pain Syndrome-An Evidence-Based Narrative Review and Etiological Hypothesis. Int J Environ Res Public Health. 2020 Sep 25;17(19):7012. doi: 10.3390/ijerph17197012.
- 62. Gofeld M, Jitendra J, Faclier G. Radiofrequency denervation of the lumbar zygapophysial joints: 10-year prospective clinical audit. Pain Physician. 2007 Mar; 10(2): 291-300.
- 63. Gupta A, Huettner DP, Dukewich M. Comparative Effectiveness Review of Cooled Versus Pulsed Radiofrequency Ablation for the Treatment of Knee Osteoarthritis: A Systematic Review. Pain Physician. 2017 Mar;20(3):155-171.
- 64. Gupta S, Gupta M, Nath S, Hess GM. Survey of European Pain Medicine Practice. Pain Physician 2012; 15:E983-E994.
- 65. Hampton H, Kalava A. Ischiorectal Approach to Cryoablation of the Pudendal Nerve Using a Handheld Device: A Report of Two Cases. Cureus. 2023 Aug 30;15(8):e44377. doi: 10.7759/cureus.44377.
- 66. Hayes, Inc. Hayes Medical Technology Directory. Radiofrequency Ablation for Chronic Low Back Pain. Lansdale, PA: Hayes, Inc.; March 2007.
- 67. Hayes, Inc. Hayes Medical Technology Directory. Radiofrequency Ablation for Cervical and Thoracic Back Pain. Lansdale, PA: Hayes, Inc.; March 2007.

- 68. Hong T, Wang H, Li G, Yao P, Ding Y. Systematic Review and Meta-Analysis of 12 Randomized Controlled Trials Evaluating the Efficacy of Invasive Radiofrequency Treatment for Knee Pain and Function. Biomed Res Int. 2019 Jun 26;2019:9037510.
- 69. Humble SR, Dalton AJ, Li L. A systematic review of therapeutic interventions to reduce acute and chronic post-surgical pain after amputation, thoracotomy or mastectomy. Eur J Pain. 2015 Apr;19(4):451-65. doi: 10.1002/ejp.567. Epub 2014 Aug 4.
- 70. Hurley RW, Adams MCB, Barad M, et al. Consensus practice guidelines on interventions for cervical spine (facet) joint pain from a multispecialty international working group. Pain Med. 2021 Nov 26;22(11):2443-2524. doi: 10.1093/pm/pnab281.
- 71. Institute for Clinical Systems Improvement (ICSI). Percutaneous Radiofrequency Ablation for Facet-Mediated Neck and Back Pain. ICSI Technology Assessment Report # 88. Bloomington, MN: ICSI; January 2005.
- 72. International Society for Advancement of Spine Surgery. ISASS Guideline Intraosseous Ablation of the Basivertebral Nerve for the Relief of Chronic Low Back Pain (December 2019). Accessed at https://www.isass.org/isass-guideline-intraosseous-ablation-of-the-basivertebral-nerve-for-the-relief-of-chronic-low-back-pain/.
- 73. Juch JNS, Maas ET, Ostelo RWJG, Groeneweg JG, Kallewaard JW, Koes BW, Verhagen AP, van Dongen JM, Huygen FJPM, van Tulder MW. Effect of Radiofrequency Denervation on Pain Intensity Among Patients With Chronic Low Back Pain: The Mint Randomized Clinical Trials. JAMA. 2017 Jul 4;318(1):68-81. doi: 10.1001/jama.2017.7918. Erratum in: JAMA. 2017 Sep 26;318(12):1188.
- 74. Kapural L, et al. Long-term retrospective assessment of clinical efficacy of radiofrequency ablation of the knee using a cooled RF system. Pain Physician. 2019; 22(5):489-494.
- 75. Kasparek M, Schneider W. Surgical treatment of Morton's neuroma: clinical results after open excision. Int Orthop. 2013 Sep; 37(9): 1857–1861.
- 76. Khalil JG, Smuck M, Koreckij T, Keel J, Beall D, Goodman B, Kalapos P, Nguyen D, Garfin S, INTRACEPT Trial Investigators. A prospective, randomized, multicenter study of intraosseous basivertebral nerve ablation for the treatment of chronic low back pain. The Spine Journal. 2019 Oct 1:19(10):1620-32.
- 77. Kim DH, Choi SS, Yoon SH, Lee SH, Seo DK, Lee IG, Choi WJ, Shin JW. Ultrasound-Guided Genicular Nerve Block for Knee Osteoarthritis: A Double-Blind, Randomized Controlled Trial of Local Anesthetic Alone or in Combination with Corticosteroid. Pain Physician. 2018 Jan;21(1):41-52.
- 78. Klessinger S, Casser HR, Gillner S, Koepp H, Kopf A, Legat M, Meiler K, Norda H, Schneider M, Scholz M, Slotty PJ, Tronnier V, Vazan M, Wiechert K. Radiofrequency Denervation of the Spine and the Sacroiliac Joint: A Systematic Review based on the Grades of Recommendations, Assesment, Development, and Evaluation Approach Resulting in a German National Guideline. Global Spine J. 2024 Sep;14(7):2124-2154. doi: 10.1177/21925682241230922. Epub 2024 Feb 6.
- 79. Kolasinski SL, Neogi T, Hochberg MC, et al. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Management of Osteoarthritis of the Hand, Hip, and Knee. Arthritis Care Res (Hoboken). 2020 Feb;72(2):149-162. doi: 10.1002/acr.24131. Epub 2020 Jan 6.
- 80. Kong FL, Bie ZX, Li B, Guo RQ, Xu S, Li XG. Comparison of analgesic effect in celiac plexus neurolysis: Ethanol injection with or without iodine-125 radioactive seeds implantation. Journal of Cancer Research and Therapeutics. 2022 Sep 1;18(5):1306-11.
- 81. Leggett LE, Soril LJ, Lorenzetti DL, Noseworthy T, Steadman R, Tiwana S, Clement F. Radiofrequency ablation for chronic low back pain: a systematic review of randomized controlled trials. Pain Res Manag. 2014 Sep-Oct;19(5):e146-53.
- 82. Li SJ, Zhang SL, Feng D. A comparison of pulsed radiofrequency and radiofrequency denervation for lumbar facet joint pain. J Orthop Surg Res. 2023 May 5;18(1):331. doi: 10.1186/s13018-023-03814-5.

- 83. Liu J, Wang T, Zhu ZH. Efficacy and safety of radiofrequency treatment for improving knee pain and function in knee osteoarthritis: a meta-analysis of randomized controlled trials. J Orthop Surg Res. 2022 Jan 15;17(1):21. doi: 10.1186/s13018-021-02906-4.
- 84. Lord SM, Barnsley L, Wallis BJ, McDonald GJ, Bogduk N. Percutaneous radio-frequency neurotomy for chronic cervical zygapophyseal-joint pain. N Engl J Med. 1996 Dec 5; 335(23): 1721-6.
- 85. Lorio M, Clerk-Lamalice O, Beall DP, Julien T. International Society for the Advancement of Spine Surgery Guideline-Intraosseous Ablation of the Basivertebral Nerve for the Relief of Chronic Low Back Pain. Int J Spine Surg. 2020;14(1):18–25. Published 2020 Feb 29. doi:10.14444/7002.
- 86. Maletis GB. Is Cooled Radiofrequency Ablation the Hot New Treatment for Knee Osteoarthritis? J Bone Joint Surg Am. 2020;102:e103(1-2).
- 87. Manchikanti KN, Atluri S, Singh V, Geffert S, Sehgal N, Falco FJE. Systematic Review: An Update of Evaluation of Therapeutic Thoracic Facet Joint Interventions. Pain Physician 2012; 15:E463-E481.
- 88. Manchikanti L MD, Boswell M MD PhD, et al. American Society of Interventional Pain Physicians. IPM Guidelines; Comprehensive Evidence-Based Guidelines for Interventional Techniques in the Management of Chronic Spinal Pain. Pain Physician 2009; 12:699-802. (Accessed 02/05/13).
- 89. Manchikanti L, Damron K, Cash K, Manchukonda R, Pampati V. Therapeutic cervical medial branch blocks in managing chronic neck pain: a preliminary report of a randomized, double-blind, controlled trial: clinical trial NCT0033272. Pain Physician. 2006 Oct; 9(4): 333-46.
- 90. Manchikanti L, Kaye AD, Soin A, Albers SL, et al. Comprehensive Evidence-Based Guidelines for Facet Joint Interventions in the Management of Chronic Spinal Pain: American Society of Interventional Pain Physicians (ASIPP) Guidelines Facet Joint Interventions 2020 Guidelines. Pain Physician. 2020 May;23(3S):S1-S127.
- 91. Manchikanti L, Manchikanti KN, Manchukonda R, Cash KA, Damron KS, Pampati V, McManus CD. Evaluation of lumbar facet joint nerve blocks in the management of chronic low back pain: preliminary report of a randomized, double-blind controlled trial: clinical trial NCT00355914. Pain Physician. 2007 May; 10(3): 425-40.
- 92. Manchikanti L, Singh V, Vilims BD, Hansen HC, Schultz DM, Kloth DS. Medial branch neurotomy in management of chronic spinal pain: systematic review of the evidence. Pain Physician. 2002 Oct; 5(4): 405-18.
- 93. Manchikanti, L., Singh, V., Kloth, D. Interventional Pain Management Practice Policies; Facet Joint Neurolytic Blocks. American Society of Interventional Pain Physicians. Accessed 02/21/11.
- 94. Manchikanti L, Datta S, Gupta S, Munglani R, Bryce DA, Ward SP, Benyamin RM, Sharma ML, Helm II S, Fellows B, Hirsch JA. Health Policy Review. A Critical Review of the American Pain Society Clinical Practice Guidelines For Interventional Techniques: Part 2. Therapeutic Interventions. Pain Physician 2010; 13:E215-E264.
- 95. Manchikanti L, Singh V, Falco FJE, Cash KA, Fellows B. Randomized trial. Comparative Outcomes of a 2-Year Follow-Up of Cervical Medial Branch Blocks in Management of Chronic Neck Pain: A Randomized, Double-Blind Controlled Trial. Pain Physician 2010; 13:437-450.
- 96. Manchikanti L, Singh V, Falco FJE, Cash KA, Pampati V, Fellows B. Randomized trial. Comparative Effectiveness of a One-Year Follow-Up of Thoracic Medial Branch Blocks in Management of Chronic Thoracic Pain: A Randomized, Double-Blind Active Controlled Trial. Pain Physician 2010; 13:535-548.
- 97. Manjunath PS, Jayalakshmi TS, Dureja GP, Prevost AT. Management of lower limb complex regional pain syndrome type 1: an evaluation of percutaneous radiofrequency thermal lumbar sympathectomy versus phenol lumbar sympathetic neurolysis a pilot study. Anesth Analg. 2008 Feb;106(2):647-9.

- 98. Mansano AM, Frederico TN, Valentin REB, Carmona MJC, Ashmawi HA. Percutaneous Radiofrequency Ablation for Trigeminal Neuralgia Management: A Randomized, Double-blind, Sham-Controlled Clinical Trial. Pain Med. 2022 Aug 27:pnac132. doi: 10.1093/pm/pnac132. Epub ahead of print. PMID: 36029256.
- 99. Mazoch MJ, et al. Effects of alcohol injection in rat sciatic nerve as a model for Morton's neuroma treatment. Foot Ankle Int. 2014 Nov;35(11):1187-91.
- 100. McLean BC, Nguyen CD, Newman DP. Cryoablation of the Infrapatellar Branch of the Saphenous Nerve Identified by Non-Invasive Peripheral Nerve Stimulator for the Treatment of Non-Surgical Anterior Knee Pain: A Case Series and Review of the Literature. Cureus. 2020 Jun 21;12(6):e8747. doi: 10.7759/cureus.8747.
- 101. Murtagh J, Foerster V. Radiofrequency neurotomy for lumber pain. Ottawa: Canadian Agency for Drugs and Technologies in Health (CADTH). 2006:4. Canadian Agency for Drugs and Technologies in Health (CADTH).
- 102. National Podiatric Medical Association. Foot Health Topics. Neuromas. ©2008 National Podiatric Medical Association.
- 103. Niemisto L, Kalso E, Malmivaara A, Seitsalo S, Hurri H. Radiofrequency denervation for neck and back pain. Cochrane Database of Systematic Reviews 2003, Issue 1. Art. No.: CD004058. DOI: 10.1002/14651858.CD004058.
- 104. Oladeji LO, Cook JL. Cooled Radio Frequency Ablation for the Treatment of Osteoarthritis-Related Knee Pain: Evidence, Indications, and Outcomes. J Knee Surg. 2019 Jan;32(1):65-71. doi: 10.1055/s-0038-1675418. Epub 2018 Nov 5. PMID: 30396206.
- 105. Orhurhu V, Urits I, Grandhi R, Abd-Elsayed A. Systematic Review of Radiofrequency Ablation for Management of Knee Pain. Curr Pain Headache Rep. 2019 Jul 8;23(8):55. doi: 10.1007/s11916-019-0792-y.
- 106. Pace A, et al. The outcome of Morton's neurectomy in the treatment of metatarsalgia. International Orthopaedics (SICOT) (2010) 34:511–515.
- 107. Pasquali C, et al. Ultrasound-Guided Alcohol Injection for Morton's Neuroma. Foot & Ankle International® 2015, Vol. 36(1) 55–59.
- 108. Petrov-Kondratov V, Chhabra A, Jones S. Pulsed Radiofrequency Ablation of Pudendal Nerve for Treatment of a Case of Refractory Pelvic Pain. Pain Physician. 2017 Mar;20(3):E451-E454.
- 109. Radoš I. TREATMENT OPTIONS FOR TRIGEMINAL NEURALGIA. Acta Clin Croat. 2022 Sep;61(Suppl 2):96-102. doi: 10.20471/acc.2022.61.s2.12. PMID: 36824641; PMCID: PMC9942467.
- 110. Ramasubba C, Cohen SP. Case Report: Cooled Sacroiliac Radiofrequency Denervation for the Treatment of Pain Secondary to Tumor Infiltration: A Case-Based Focused Literature Review. Pain Physician 2013; 16:1-8.
- 111. Rocha A, Plancarte R, Nataren RGR, Carrera IHS, Pacheco VALR, Hernandez-Porras BC. Effectiveness of Superior Hypogastric Plexus Neurolysis for Pelvic Cancer Pain. Pain Physician. 2020 Mar;23(2):203-208.
- 112. Rupert MP, Lee M, Manchikanti L, Datta S, Cohen SP. Evaluation of Sacroiliac Joint Interventions: A Systematic Appraisal of the Literature. Pain Physician 2009; 12:399-418.
- 113. Sehgal N, Dunbar EE, Shah RV, Colson J. Systematic review of diagnostic utility of facet (zygapophysial) joint injections in chronic spinal pain: an update. Pain Physician. 2007 Jan; 10(1): 213-28.
- 114. Shabat S, Leitner Y, Bartal G, Folman Y. Radiofrequency treatment has a beneficial role in reducing low back pain due to facet syndrome in octogenarians or older. Clin Interv Aging. 2013;8:737-40.

- 115. Shah N, Engle AM, Raggi E, Alter B, Emerick T. Pulsed Radiofrequency Ablation: An Alternative Treatment Modality for Radiation-Induced Brachial Plexopathy. Pain Med. 2021 Mar 18;22(3):749-753. doi: 10.1093/pm/pnaa382.
- 116. Texakalidis P, Xenos D, Tora MS, et al. Comparative safety and efficacy of percutaneous approaches for the treatment of trigeminal neuralgia: A systematic review and meta-analysis. Clin Neurol Neurosurg. 2019; 182:112-122.
- 117. Truumees E, Macadaeg K, Pena E, et al. A prospective, open-label, single-arm, multi-center study of intraosseous basivertebral nerve ablation for the treatment of chronic low back pain. Eur Spine J. 2019;28(7):1594–1602. doi:10.1007/s00586-019-05995-2.
- 118. UpToDate. Interventional therapies for chronic pain. 2025. Accessed at uptodate.com.
- 119. UpToDate. Investigational approaches to the management of osteoarthritis. 2025. Accessed at uptodate.com.
- 120. UpToDate. Management of knee osteoarthritis. 2025. Accessed at uptodate.com.
- 121. UpToDate. Management of nonradicular neck pain in adults. 2025. Accessed at uptodate.com
- 122. UpToDate. Subacute and chronic low back pain: Nonsurgical interventional treatment. 2025. Accessed at uptodate.com.
- 123. UpToDate. Trigeminal neuralgia. 2025. Accessed at uptodate.com.
- 124. U.S. Food & Drug Administration (FDA). 510(k) Premarket Notification K153272: Intracept Intraosseous Nerve Ablation System (July 9, 2016).
- 125. Van Oosterwyck W, Vander Cruyssen P, Castille F, Van de Kelft E, Decaigny V. Lumbar Facet Joint Disease: What, Why, and When? Life (Basel). 2024 Nov 14;14(11):1480. doi: 10.3390/life14111480.
- 126. van Tilburg CW, Schuurmans FA, Stronks DL, Groeneweg JG, Huygen FJ. Randomized Sham-controlled Double-Blind Multicenter Clinical Trial to Ascertain the Effect of Percutaneous Radiofrequency Treatment for Sacroiliac Joint Pain: Three-month Results. Clin J Pain. 2016 Nov;32(11):921-926. doi: 10.1097/AJP.000000000000351.
- 127. Walega D, McCormick Z, Manning D, Avram M. Radiofrequency ablation of genicular nerves prior to total knee replacement has no effect on postoperative pain outcomes: a prospective randomized sham-controlled trial with 6-month follow-up. Reg Anesth Pain Med. 2019 Apr 25:rapm-2018-100094. doi: 10.1136/rapm-2018-100094. Epub ahead of print. PMID: 31023931.
- 128. Xiao L, Shu F, Xu C, Zhang Z, Huang L, Wang X, Shang Y. Highly selective peripheral nerve radio frequency ablation for the treatment of severe knee osteoarthritis. 2018. Experimental and Therapeutic Medicine 16: 3973-3977.
- 129. Xu R, Xie ME, Jackson CM. Trigeminal Neuralgia: Current Approaches and Emerging Interventions. J Pain Res. 2021 Nov 3;14:3437-3463. doi: 10.2147/JPR.S331036.
- 130. Zeitlinger L, Kopinski J, DiPasquale T. Genicular nerve ablation: a systematic review of procedure outcomes for chronic knee pain. Current Orthopaedic Practice. Volume 30, Number 5, September/October 2019.
- 131. Zhang X, Peng L, Liu D. Radiofrequency Therapies for Trigeminal Neuralgia: A Systematic Review and Updated Meta-analysis. Pain Physician. 2022 Dec;25(9):E1327-E1337.
- 132. Zundert JV, Vanelderen P, Kessels A, van Kleef M. Radiofrequency Treatment of Facet-related Pain: Evidence and Controversies. Curr Pain Headache Rep (2012) 16:19–25.

COMMITTEE APPROVAL:

This Medical Coverage Guideline (MCG) was approved by the Florida Blue Medical Policy and Coverage Committee on 03/27/25.

GUIDELINE UPDATE INFORMATION:

11/15/07	New Medical Coverage Guideline.
05/15/09	Scheduled review; update to description section to include medical necessity
03/13/03	management statement, update position statement to include coverage criteria, and
	update to reimbursement statement limitations.
09/15/09	Update position statement.
07/15/10	Revisions consisting of Billing and Coding section changes to include coding guidelines.
11/15/10	Revision; MCG title changed to "Neurolysis"; updated description section to include
11/13/10	chemical neurolysis for foot pain; updated position statement to include coverage
	criteria for neurolysis for foot pain; revised CPT coding to include 64632; revised coding
	notes; updated ICD-9 coding to include 355.5, 355.6, 355.79, 355.8 and 728.71; added
	related ICD-10 codes; revised reimbursement section; added Medicare exception; added
	MCG 02-61000-29 as a related guideline; updated references; reformatted guideline.
05/15/11	Scheduled review; position statement unchanged; references updated.
07/15/11	Revision; formatting changes.
10/15/11	Revision; added experimental/investigational coverage statement for neurolysis of
10/15/11	sacroiliac (SI) joints; formatting changes.
01/01/12	Annual HCPCS coding update. Added 64633, 64634, 64635 and 64636. Deleted 64622,
01/01/12	64623, 64626 and 64627. Updated Coding Notes and Reimbursement Information
	sections.
03/15/12	Scheduled review. Revised description section, position statement and ICD9/ICD10
00, 10, 12	coding sections; deleted Medicare Advantage Program Exception; updated references
	and reformatted guideline.
07/15/12	Revision; added program exception for Medicare Advantage products.
03/15/13	Scheduled review. Revised position statement (chemical neurolysis is E/I for facet
, , ,	neurolysis). Revised description, ICD10 coding and Medicare Advantage program
	exception (added ICD9 and HCPCS codes). Updated references and reformatted
	guideline.
05/11/14	Revision: Program Exceptions section updated.
07/01/15	Scheduled review. Revised description and position statement. Updated references.
10/01/15	Revision; updated ICD10 coding section.
11/01/15	Revision: ICD-9 Codes deleted.
07/01/16	Quarterly CPT/HCPCS update. Added codes 0440T, 0441T, AND 0442T.
08/15/16	Revision; updated ICD10 coding section.
10/01/16	ICD-10 coding update: added code G57.63.
02/15/17	Revision; updated Reimbursement Information section.
04/15/17	Revision: updated pain relief criteria and frequency of treatment criteria for facet joint
	neurolysis. Updated references.
06/15/17	Revision: added codes 64620 and 64630.
10/15/17	Revision: Revised MCG title and description section. Added clarifying language to
·	Position Statement regarding when neurolysis is considered E/I. Revised ICD10 coding
ļ	section, Reimbursement Information section, and definitions. Updated references.

08/15/18	Revision: added coverage statement (E/I) for intralesional alcohol injections for
	treatment of Morton's neuroma. Updated program exceptions section and references.
03/15/19	Revision: updated frequency of treatment section and references.
07/15/19	Scheduled review. Revised home exercise program requirements. Updated references.
09/01/19	Revision: clarified what constitutes an "active" modality.
01/01/20	Annual CPT/HCPCS coding update. Added 64624, 64625.
04/15/20	Revision; updated reimbursement information section.
05/15/20	Revision; updated ICD10 coding section.
06/15/20	Revision: added codes C9752, C9753 (investigational). Updated references.
07/15/20	Scheduled review. Maintained position statement and updated references.
12/15/20	Unscheduled review. Maintained position statement, revised definitions and updated
	references.
05/15/21	Revision. Added Intracept procedure for clarity; revised ICD10 coding.
06/15/21	Scheduled review. Maintained position statement and updated references.
10/01/21	ICD10 coding update: added codes M54.50, M54.51, M54.59; deleted code M54.5.
01/01/22	Annual CPT/HCPCS coding update. Added 64628, 64629. Deleted C9752, C9753.
02/15/22	Deleted codes 64628, 64629 (refer to MCG 02-61000-20, Percutaneous Intradiscal
	Electrothermal Annuloplasty, Radiofrequency Annuloplasty, and Biacuplasty).
08/15/22	Revision. Updated CPT coding (added codes 64600, 64605, 64610, 64680, 64681).
06/10/23	Scheduled review. Revised description, position statement, ICD10 coding, CPT coding,
	Reimbursement Information section, Billing and Coding Information section, and
	Medicare Advantage program exception. Updated references.
08/21/23	Update to Program Exceptions section.
04/15/24	Scheduled review. Revised description, maintained position statement, and updated
	references.
06/15/24	Revision. Updated criteria for controlled medial branch blocks.
01/01/25	Annual CPT/HCPCS coding update. Added C9809.
04/15/25	Scheduled review. Revised description, maintained position statement and updated references.