

02-61000-36

Original Effective Date: 03/15/10

Reviewed: 09/27/18

Revised: 01/01/20

## Subject: Minimally Invasive Fusion Techniques

THIS MEDICAL COVERAGE GUIDELINE IS NOT AN AUTHORIZATION, CERTIFICATION, EXPLANATION OF BENEFITS, OR A GUARANTEE OF PAYMENT, NOR DOES IT SUBSTITUTE FOR OR CONSTITUTE MEDICAL ADVICE. ALL MEDICAL DECISIONS ARE SOLELY THE RESPONSIBILITY OF THE PATIENT AND PHYSICIAN. BENEFITS ARE DETERMINED BY THE GROUP CONTRACT, MEMBER BENEFIT BOOKLET, AND/OR INDIVIDUAL SUBSCRIBER CERTIFICATE IN EFFECT AT THE TIME SERVICES WERE RENDERED. THIS MEDICAL COVERAGE GUIDELINE APPLIES TO ALL LINES OF BUSINESS UNLESS OTHERWISE NOTED IN THE PROGRAM EXCEPTIONS SECTION.

[Position Statement](#)

[Billing/Coding](#)

[Reimbursement](#)

[Program Exceptions](#)

[Definitions](#)

[Related Guidelines](#)

[Other](#)

[References](#)

[Updates](#)

### **DESCRIPTION:**

#### **Axial Lumbar Interbody Fusion (AxiaLIF)**

The procedure for 1-level axial lumbosacral interbody fusion (axial LIF) is as follows (1): Under fluoroscopic monitoring, a blunt guide pin introducer is passed through a 15- to 20-mm incision lateral to the coccyx and advanced along the midline of the anterior surface of the sacrum. A guide pin is introduced and tapped into the sacrum. A series of graduated dilators are advanced over the guide pin, and a dilator sheath attached to the last dilator is left in place to serve as a working channel for the passage of instruments. A cannulated drill is passed over the guide pin into the L5-S1 disc space to rest on the inferior endplate of L5. It is followed by cutters alternating with tissue extractors, and the nucleus pulposus is debulked under fluoroscopic guidance. Next, bone graft material is injected to fill the disc space. The threaded rod is placed over the guide pin and advanced through the sacrum into L5. The implant is designed to distract the vertebral bodies and restore disc and neural foramen height. Additional graft material is injected into the rod, where it enters into the disc space through holes in the axial rod. A rod plug is then inserted to fill the cannulation of the axial rod. Percutaneous placement of pedicle or facet screws may be used to provide supplemental fixation. An advantage of axial LIF is that it allows preservation of the annulus and all paraspinal soft tissue structures. However, there is an increased need for fluoroscopy and an inability to address intracanal pathology or visualize the discectomy procedure directly. Complications of the axial approach may include perforation of the bowel and injury to blood vessels and/or nerves.

#### **Sacroiliac joint fusion/stabilization**

**Sacroiliac fusion** involves bony fusion of the sacroiliac joint for stabilization. This has been proposed for treatment of chronic sacroiliac pain. Several percutaneous or minimally invasive fixation/fusion devices have received marketing clearance by the FDA. These include the SI-FIX Sacroiliac Joint Fusion System (cortical implant with cancellous autograft), the IFUSE® Implant System (titanium triangular implant), the

Slmmetry® Sacroiliac Joint Fusion System (system of threaded screws), the Silex™ (cylindrical implant) and the SI-LOK® (hydroxyapatite coated screw system).

Surgical devices for annular repair or modulation after spinal surgery, according to FDA documentation, are intended for use in soft tissue approximation, or repair of annular defect, for procedures such as general and orthopedic surgery. There are several FDA-approved, commercially marketed devices.

## **POSITION STATEMENT:**

**Axial lumbosacral interbody fusion (Axial LIF)** is considered **experimental or investigational** for **all** indications. There is insufficient scientific evidence to permit conclusions concerning the effect of this technology on net health outcomes.

**Open sacroiliac joint fusion/stabilization (27280)** meets the definition of **medical necessity** for any of the following indications:

- A tumor involving the sacrum and/or sacroiliac joint
- As adjunctive treatment of sacroiliac joint infection, following successful treatment of the infection
- Following traumatic injury of the sacroiliac joint (e.g., following pelvic ring fracture)
- When performed as a part of multi-segment long fusion to correct spinal deformity associated with scoliosis or kyphosis

**Open sacroiliac joint fusion/stabilization** for the treatment of all other indications, including chronic back pain, is considered **experimental or investigational**. The available scientific evidence remains insufficient to permit conclusions concerning the effect of this technology on net health outcomes.

**Minimally invasive sacroiliac joint fusion/stabilization** using a titanium triangular implant **meets the definition of medical necessity** when **ALL** of the following criteria have been met:

- Pain is at least 5 on a 0 to 10 rating scale; and pain impacts quality of life or limits activities of daily living, **AND**
- There is an absence of generalized pain behavior (eg, somatoform disorder) or generalized pain disorders (eg, fibromyalgia), **AND**
- Patients have undergone and failed a minimum 6 months of intensive nonoperative treatment that must include medication optimization, activity modification, and 6 weeks of active physical therapy targeted at the lumbar spine, pelvis, sacroiliac joint, and hip, including a home exercise program, **AND**
- Pain is caudal to the lumbar spine (L5 vertebra), localized over the posterior sacroiliac joint, and consistent with sacroiliac joint pain, **AND**
- A thorough physical examination demonstrates localized tenderness with palpation over the sacral sulcus (Fortin's point) in the absence of tenderness of similar severity elsewhere, **AND**
- There is a positive response to a cluster of 3 provocative tests (eg, thigh thrust test, compression test, Gaenslen sign, distraction test, Patrick test, or posterior provocation test), **AND**
- Diagnostic imaging studies include **ALL** of the following:
  - Imaging (plain radiographs and computed tomography or magnetic resonance imaging) of the sacroiliac joint excludes the presence of destructive lesions (eg, tumor, infection) or inflammatory arthropathy of the sacroiliac joint, **AND**
  - Imaging of the pelvis (anteroposterior plain radiograph) rules out concomitant hip pathology, **AND**

- Imaging of the lumbar spine (computed tomography or magnetic resonance imaging) is performed to rule out neural compression or other degenerative condition that can be causing low back or buttock pain, **AND**
- Imaging of the sacroiliac joint indicates evidence of injury and/or degeneration, **AND**
- There is at least a 75% reduction in pain for the expected duration of the anesthetic used following an image-guided, contrast-enhanced intra-articular sacroiliac joint injection on 2 separate occasions, **AND**
- A trial of a therapeutic sacroiliac joint injection (ie, corticosteroid injection) has been performed at least once.

Minimally invasive fusion/stabilization of the sacroiliac joint for the treatment of back pain presumed to originate from the sacroiliac joint is considered **experimental or investigational** under all other conditions, and with any devices other than a titanium triangular implant.

The use of surgical devices for annular repair/modulation following spinal surgery (e.g. Inclose™ Surgical Mesh System, Xclose™ Tissue Repair System) is considered **experimental or investigational** due to the lack of scientific peer-reviewed literature demonstrating improvement in health outcomes.

### **BILLING/CODING INFORMATION:**

#### **CPT Coding:**

22586	Arthrodesis, pre-sacral interbody technique, including disc space preparation, discectomy, with posterior instrumentation, with image guidance, includes bone graft when performed, L5-S1 interspace ( <b>investigational</b> )
27279	Arthrodesis, sacroiliac joint, percutaneous or minimally invasive (indirect visualization), with image guidance, includes obtaining bone graft when performed, and placement of transfixing device
27280	Arthrodesis, open, sacroiliac joint, including obtaining graft, including instrumentation, when performed

#### **HCPCS Coding:**

C9757	Laminotomy (hemilaminectomy), with decompression of nerve root(s), including partial facetectomy, foraminotomy and excision of herniated intervertebral disc, and repair of annular defect with implantation of bone anchored annular closure device, including annular defect measurement, alignment and sizing assessment, and image guidance; 1 interspace, lumbar ( <b>investigational</b> )
-------	--

### **REIMBURSEMENT INFORMATION:**

Refer to section entitled **POSITION STATEMENT**.

### **PROGRAM EXCEPTIONS:**

**Federal Employee Program (FEP):** Follow FEP guidelines.

**State Account Organization (SAO):** Follow SAO guidelines.

**Medicare Advantage:** The following coverage article was reviewed on the last guideline reviewed date: Local Coverage Article: Medical review article for percutaneous minimally invasive fusion/stabilization of the SACROILIAC JOINT (A55120) located at cms.gov.

The following Local Coverage Determination (LCD) was reviewed on the last guideline review date: Local Coverage Determination (LCD) L33777: Noncovered Services. located at fcso.com.

## **DEFINITIONS:**

**Anulus fibrosus:** The ring of fibrocartilage and fibrous tissue forming the circumference of the intervertebral disc; surrounds the nucleus pulpous, which can herniate when the anulus is diseased or injured.

**Fibromyalgia:** A disorder that causes muscle pain and fatigue in the form of "tender points" on the body in response to applied pressure; generally on the neck, shoulders, back, hips, arms, and legs.

**Interbody:** Between the bodies of two adjacent vertebra.

**Interspace:** Any space between two similar objects.

**Somatoform disorder:** A form of mental illness that causes one or more bodily symptoms, including pain; symptoms may or may not be traceable to a physical cause including general medical conditions, other mental illnesses, or substance abuse; also known as somatic symptom disorder or somatization disorder.

## **RELATED GUIDELINES:**

[Bone Morphogenetic Protein \(BMP\), 02-20000-32](#)  
[Interspinous and Interlaminar Stabilization/Distracton \(Spacers\) and Fixation \(Fusion\) Devices, 02-20000-36](#)  
[Automated Percutaneous Discectomy, Laser Discectomy, Endoscopic Discectomy, or DISC Nucleoplasty™ as Techniques of Intervertebral Disc Decompression, 02-61000-32](#)

## **OTHER:**

None applicable

## **REFERENCES:**

1. Ackerman SJ, Polly Jr DW, Knight T, Schneider K, Holt T, Cummings J. Comparison of the costs of nonoperative care to minimally invasive surgery for sacroiliac joint disruption and degenerative sacroiliitis in a United States Medicare population: potential economic implications of a new minimally-invasive technology. ClinicoEconomics and outcomes research: CEOR (2013), 5, 575.
2. AHRQ national Guideline Clearinghouse. Guideline Summary NGC-8134. Laminectomy and fusion for the treatment of cervical degenerative myelopathy. American Association of Neurological Surgeons; Congress of Neurological Surgeons. 2009 Aug.
3. AHRQ national Guideline Clearinghouse. Guideline Summary NGC-8766. Diagnosis and treatment of degenerative lumbar spinal stenosis. North American Spine Society (NASS); 2011.
4. AHRQ national Guideline Clearinghouse. Guideline Summary NGC-9903. Clinical guidelines for diagnosis and treatment of lumbar disc herniation with radiculopathy. North American Spine Society; 2012.

5. AHRQ national Guideline Clearinghouse. Guideline Summary NGC-9327. Low back disorders. American College of Occupational and Environmental Medicine (ACOEM); 2011.
6. AHRQ national Guideline Clearinghouse. Guideline Summary NGC-10118: Hip & pelvis (acute & chronic). Work Loss Data Institute; 2013.
7. AHRQ national Guideline Clearinghouse. Guideline Summary NGC-10121. Low back - lumbar & thoracic (acute & chronic). Work Loss Data Institute; 2013.
8. American Academy of Orthopaedic Surgeons. Anterior Lumbar Interbody Fusion. Accessed at <http://orthoinfo.aaos.org/topic.cfm?topic=A00595>. Copyright 2010 American Academy of Orthopaedic Surgeons.
9. American Academy of Orthopaedic Surgeons. Posterior Lumbar Interbody Fusion and Transforaminal Lumbar Interbody Fusion. Accessed at <http://orthoinfo.aaos.org/topic.cfm?topic=A00596>. Copyright 2010 American Academy of Orthopaedic Surgeons.
10. Anand N, Baron EM. Minimally invasive approaches for the correction of adult spinal deformity. *Eur Spine J*. 2013 Mar;22 Suppl 2:S232-41. doi: 10.1007/s00586-012-2344-6. Epub 2012 May 10.
11. Aryan HE, Newman CB, Gold JJ, Acosta Jr. FL, Coover C, Ames CP. Percutaneous Axial Lumbar Interbody Fusion (AxialLIF) of the L5-S1 Segment: Initial Clinical and Radiographic Experience. *Minim Invas Neurosurg* 2008; 51: 225 – 230.
12. Ashman B, Norvell DC, Hermsmeyer JT. Chronic sacroiliac joint pain: fusion versus denervation as treatment options. *Evidence-Based Spine-Care Journal* Volume 1/Issue 3 — 2010.
13. Bahtia NN, Timon S, Wieser ES, Wang JC. TLIF, XLIF or ALIF for adjacent segment degenerative disc disease. *NASS SpineLine*, May/June 2008.
14. Bailey A, Messer J, Griffith SL. (2010). Prospective, Randomized Controlled Study of Repairing the Anulus Fibrosus after Lumbar Discectomy: A Single Surgeon's Experience. *The Spine Journal*, 10(9), S127.
15. Blue Cross Blue Shield Association. Medical Policy Reference Manual. 6.01.23 Diagnosis and Treatment of Sacroiliac Joint Pain. December 2017.
16. Blue Cross Blue Shield Association. Medical Policy Reference Manual. 7.01.115 Percutaneous Axial Anterior Lumbar Fusion. (Archived 11/10/11).
17. Blue Cross Blue Shield Association. Medical Policy Reference Manual. 7.01.130 Axial Lumbosacral Interbody Fusion. April 2018.
18. Blumenthal, SL, Guyer RD. (2010, October). Anular Repair After Lumbar Discectomy: Preliminary Observations From An Ongoing, Prospective, Randomized, Controlled Clinical Trial: 41. In *Spine Journal Meeting Abstracts* (p. 41).
19. Boachie-Adjei O, Cho W, King AB. Axial lumbar interbody fusion (AxialLIF) approach for adult scoliosis. *Eur Spine J*. 2013 Mar;22 Suppl 2:S225-31.
20. Bohinski RJ, Jain VV, Tobler WD. Presacral retroperitoneal approach to axial lumbar interbody fusion: a new, minimally invasive technique at L5-S1: Clinical outcomes, complications, and fusion rates in 50 patients at 1-year follow-up. *SAS J*. 2010 Jun 1;4(2):54-62. doi: 10.1016/j.esas.2010.03.003. eCollection 2010.
21. Bradley WD, et al. Minimally invasive trans-sacral approach to L5-S1 interbody fusion: Preliminary results from 1 center and review of the literature. *Int J Spine Surg*. 2012 Dec 1;6:110-4. doi: 10.1016/j.ijssp.2011.12.005. eCollection 2012.
22. Capobianco R, et al. Safety and effectiveness of minimally invasive sacroiliac joint fusion in women with persistent post-partum posterior pelvic girdle pain: 12-month outcomes from a prospective, multi-center trial. *Springerplus*. 2015 Oct 5;4:570.

23. Caputo AM, Michael KW, Chapman Jr TM, Massey GM, Howes CR, Isaacs RE, Brown CR. Clinical outcomes of extreme lateral interbody fusion in the treatment of adult degenerative scoliosis. *ScientificWorldJournal*. 2012;2012:680643.
24. Cher DJ, Polly DW. Improvement in Health State Utility after Sacroiliac Joint Fusion: Comparison to Normal Populations. *Global Spine Journal*. 2016 Mar;6(2):100.
25. Cher DJ, Reckling WC, Capobianco RA. Implant survivorship analysis after minimally invasive sacroiliac joint fusion using the iFuse Implant System (®). *Medical devices (Auckland, NZ)*. 2015 Nov 23;8:485.
26. Cho CB, et al, Anterior Lumbar Interbody Fusion with Stand-Alone Interbody Cage in Treatment of Lumbar Intervertebral Foraminal Stenosis: Comparative Study of Two Different Types of Cages, *J Korean Neurosurg Soc*, 47: 352-357, 2010.
27. Choy W-S, Kim KJ, Lee SK, Park HJ. Anterior Pelvic Plating and Sacroiliac Joint Fixation in Unstable Pelvic Ring Injuries. *Yonsei Med J* 53(2):422-426, 2012.
28. ClinicalTrials.gov, A Clinical Study of the GO-LIF® Approach for Lumbar Spinal Fixation. NCT00810433. Sponsored by Mazor Surgical Technologies, Ltd, last updated: February 6, 2011. Accessed 12/28/10.
29. ClinicalTrials.gov. Osteocel® Plus in eXtreme Lateral Interbody Fusion (XLIF®), NCT00948532. Sponsored by NuVasive, last updated June 10, 2011. Accessed 01/02/13.
30. ClinicalTrials.gov. XLIF® vs. MAS®TLIF for the Treatment of Symptomatic Lumbar Degenerative Spondylolisthesis With or Without Central Stenosis. NCT01024699. Sponsored by NuVasive, accessed 01/06/12. last updated on June 10, 2011. Accessed 01/06/12.
31. ClinicalTrials.gov. Randomized Study of Anular Repair with the Xclose Tissue Repair System. NCT00760799. Sponsored by Anulex Technologies, Inc, last updated: May 1, 2012. Accessed 12/28/10.
32. ClinicalTrials.gov. NCT00965380. Trinity Evolution in Posterior or Transforaminal Lumbar Interbody Fusion (PLIF/TLIF) (TLF), Sponsored by Orthofix Inc., last updated on May 22, 2012. Accessed 01/16/12.
33. ClinicalTrials.gov. Osteocel® Plus in Anterior Lumbar Interbody Fusion (ALIF). NCT00948831. NuVasive, last updated June 10, 2011. Accessed 01/02/13.
34. ClinicalTrials.gov. NCT01640353: Sacroiliac Joint Fusion With iFuse Implant System (SIFI). August 2014.
35. ClinicalTrials.gov. NCT01681004: Investigation of Sacroiliac Fusion Treatment (INSITE). September 2014.
36. ClinicalTrials.gov. NCT01741025: iFuse Implant System® Minimally Invasive Arthrodesis (iMIA). January 2014.
37. ClinicalTrials.gov. NCT01861899: Treatment of Sacroiliac Dysfunction With SI-LOK® Sacroiliac Joint Fixation SI-SI-LOK. May 2013.
38. ClinicalTrials.gov. NCT02074761: Evolusion Study Using the Zyga Slimmetry Sacroiliac Joint Fusion System. July 2015.
39. ClinicalTrials.gov. NCT02064322: SAIF: Sacroiliac Fusion Study. April 2015.
40. ClinicalTrials.gov. NCT02270203: LOIS: Long-Term Follow-Up in INSITE/SIFI (LOIS) (SI-Bone, Inc.) (August 2016).
41. Cohen A, Miller LE, Block JE. Minimally invasive presacral approach for revision of an Axial Lumbar Interbody Fusion rod due to fall-related lumbosacral instability: a case report. *J Med Case Rep*. 2011 Sep 29;5:488. doi: 10.1186/1752-1947-5-488,

42. Cummings Jr J, Capobianco RA. Minimally invasive sacroiliac joint fusion: one-year outcomes in 18 patients. *Ann Surg Innov Res*. 2013 Sep 16;7(1):12.
43. Dengler J, Duhon B, et al. Predictors of Outcome in Conservative and Minimally Invasive Surgical Management of Pain Originating From the Sacroiliac Joint: A Pooled Analysis. *Spine (Phila Pa 1976)*. 2017 Nov 1;42(21):1664-1673.
44. Dengler JD, Kools D, et al. 1-Year Results of a Randomized Controlled Trial of Conservative Management vs. Minimally Invasive Surgical Treatment for Sacroiliac Joint Pain. *Pain Physician*. 2017 Sep;20(6):537-550.
45. Derman PB, Albert TJ. Interbody Fusion Techniques in the Surgical Management of Degenerative Lumbar Spondylolisthesis. *Curr Rev Musculoskelet Med*. 2017 Dec;10(4):530-538. doi: 10.1007/s12178-017-9443-2.
46. DeVine JG, Gloystein D, Singh N. A novel alternative for removal of the AxiaLif (TranS1) in the setting of pseudarthrosis of L5–S1. *The Spine Journal* 9 (2009) 910–915.
47. Duhon BS, Cher DJ, Wine KD, Lockstadt H, Kovalsky D, Soo CL. Safety and 6-month effectiveness of minimally invasive sacroiliac joint fusion: a prospective study. *Med Devices (Auckl)*. 2013 Dec 13;6:219-29.
48. Duhon BS, et al. Triangular Titanium Implants for Minimally Invasive Sacroiliac Joint Fusion: A Prospective Study. *Global Spine Journal*. June 30, 2015.
49. Duhon BS, et al. Triangular Titanium Implants for Minimally Invasive Sacroiliac Joint Fusion: 2-Year Follow-Up from a Prospective Multicenter Trial. *Int J Spine Surg*. 2016 Apr 20;10:13.
50. Duhon BS, et al. Triangular Titanium Implants for Minimally Invasive Sacroiliac Joint Fusion: A Prospective Study. *Global spine journal*. 2016 May;6(3):257-69.
51. Eck JC, Hodges S, Humphreys SC. Minimally invasive lumbar spinal fusion. *J Am Acad Orthop Surg*. 2007; 15(6):321-329.
52. ECRI Institute Health Technology Information Service: Product Brief. iFuse Implant System (SI-Bone, Inc.) for Minimally Invasive Sacroiliac Joint Fusion (May 2016).
53. Endres S, Ludwig E. Outcome of distraction interference arthrodesis of the sacroiliac joint for sacroiliac arthritis. *Indian journal of orthopaedics* (2013), 47(5), 437.
54. Erkan S, et al, Biomechanical Evaluation of a New AxiaLIF Technique for Two-Level Lumbar Fusion, *Eur Spine J* (2009), 18:807-814.
55. Florida Medicare Part B Local Coverage Determination. Non-covered Services (L29288). 02/02/09. Retired 09/30/15.
56. First Coast Service Options, Inc. (FCSO). Local Coverage Article: Medical review article for percutaneous minimally invasive fusion/stabilization of the SACROILIAC JOINT (A55120). Accessed at cms.gov.
57. First Coast Service Options, Inc. (FCSO). Local Coverage Determination (LCD) L33777: Noncovered Services. 07/03/18.
58. Fokter SK. Update review and clinical presentation on chronic low back pain treated by AxiaLIF. *Eur J Orthop Surg Traumatol* (2011) 21:39–42.
59. Garber T, Ledonio CG, Polly Jr DW. How Much Work Effort is Involved in Minimally Invasive Sacroiliac Joint Fusion?. *International journal of spine surgery*. 2015 Nov 6;9:58.
60. Griffith SL, Davis RJ, Hutton WC. Repair of the annulus fibrosus of the lumbar disc. In: Davis RJ, Girardi FP, Cammisa FP editor. *Nucleus arthroplasty technology in spinal care, volume II: biomechanics and development*. Bloomington, MN: Raymedica Inc; 2007;p. 41–48.
61. Gundanna MI, Miller LE, Block JE. Complications with axial presacral lumbar interbody fusion: A 5-year postmarketing surveillance experience. *SAS Journal* 5 (2011) 90–94.

62. Ha KY, Lee JS, Kim KW. Degeneration of sacroiliac joint after instrumented lumbar or lumbosacral fusion: a prospective cohort study over five-year follow-up. *Spine (Phila Pa 1976)*. 2008 May 15;33(11):1192-8.
63. Habib A, Smith ZA, Lawton CD, Fessler RG. Minimally Invasive Transforaminal Lumbar Interbody Fusion: A Perspective on Current Evidence and Clinical Knowledge. *Minimally Invasive Surgery Volume 2012*, Article ID 657342.
64. Hadjipavlou A, Alpantaki K, Katonis P, Vastardis G, Tzermiadianos M, Benardos N. Safety and effectiveness of retrorectal presacral approach for lumbosacral axial instrumentation. A clinical study. *Acta Orthop Belg*. 2013 Apr;79(2):222-9.
65. Heiney J, Capobianco R, Cher D. A systematic review of minimally invasive sacroiliac joint fusion utilizing a lateral transarticular technique. *Int J Spine Surg*. 2015 Jul 22;9:40.
66. Inamasu J, Guiot BH. Laparoscopic Anterior Lumbar Interbody Fusion: A Review of Outcome Studies. *Minim Invas Neurosurg* 2005; 48: 340±347.
67. International Society for the Advancement of Spine Surgery (ISASS). Proposed Recommendations for "Coverage Criteria": Minimally Invasive Sacroiliac Joint Fusion. March 2014. Accessed at <http://www.isass.org/>.
68. International Society for the Advancement of Spine Surgery. Statement on Coding Changes for Minimally Invasive SI Joint Fusion July 2013). Accessed at [www.isass.org](http://www.isass.org).
69. InterQual® 2013.2. CP Procedures: Fusion, Lumbar Spine.
70. Issack PS, et al. Axial lumbosacral interbody fusion appears safe as a method to obtain lumbosacral arthrodesis distal to long fusion constructs. *HSS J*. 2012 Jul; 8(2): 116–121.
71. Ledonio CGT, Polly DW, Swiontkowski MF, Cummings Jr JT. Comparative effectiveness of open versus minimally invasive sacroiliac joint fusion. *Medical Devices: Evidence and Research* 2014;7 187–193.
72. Ledonio CG, Polly DW Jr, Swiontkowski MF. Minimally invasive versus open sacroiliac joint fusion: are they similarly safe and effective? *Clin Orthop Relat Res*. 2014 Jun;472(6):1831-8.
73. Lee KH, Yue WM, Yeo W, Soeharno H, Tan SB. Clinical and radiological outcomes of open versus minimally invasive transforaminal lumbar interbody fusion. *Eur Spine J*. 2012 Nov;21(11):2265-70.
74. Lindley EM, McCullough MA, Burger EL, Brown CW, Patel VV. Complications of axial lumbar interbody fusion. *J Neurosurg Spine* 15:273–279, 2011.
75. Lindsey DP, et al. Evaluation of a minimally invasive procedure for sacroiliac joint fusion – an in vitro biomechanical analysis of initial and cycled properties. *Medical Devices: Evidence and Research* 2014;7 131–137.
76. Lindsey DP, Kiapour A, Yerby SA, Goel VK. Sacroiliac Joint Fusion Minimally Affects Adjacent Lumbar Segment Motion: A Finite Element Study. *International journal of spine surgery*. 2015 Nov 13;9:64.
77. Lorio MP. ISASS Policy 2016 Update - Minimally Invasive Sacroiliac Joint Fusion. *Int J Spine Surg*. 2016 Jul 13;10:26.
78. Lorio MP, Polly Jr DW, Ninkovic I, Ledonio CGT, Hallas K, Andersson G. Utilization of Minimally Invasive Surgical Approach for Sacroiliac Joint Fusion in Surgeon Population of ISASS and SMISS Membership. *The Open Orthopaedics Journal*, 2014, 8, 1-6.
79. Lorio MP, Rashbaum R. ISASS Policy Statement – Minimally Invasive Sacroiliac Joint Fusion. *International Journal of Spine Surgery Volume 8 Article 25*.
80. MacMillan M, et al. Description of a transosseous approach to the L5-S1 disc and 2 clinical case reports. *Int J Spine Surg*. 2012 Dec 1;6:178-83. doi: 10.1016/j.ijssp.2012.06.001. eCollection 2012.



81. Malham GM, Ellis NJ, Parker RM, Seex KA. Clinical outcome and fusion rates after the first 30 extreme lateral interbody fusions. *ScientificWorldJournal*. 2012;2012:246989.
82. Manchikanti L, et al. An Update of Comprehensive Evidence-Based Guidelines for Interventional Techniques in Chronic Spinal Pain. Part II: Guidance and Recommendations. *Pain Physician* 2013; 16:S49-S283 • ISSN 1533-3159.
83. Marawar S, et al. Changes in neuroforaminal height with 2 level axial presacral lumbar interbody fusion at L4-S1. *Int J Spine Surg*. 2014 Dec 1;8. doi: 10.14444/1002. eCollection 2014.
84. Marchi L, Abdala N, Oliveira L, Amaral R, Coutinho E, Pimenta L. Stand-Alone Lateral Interbody Fusion for the Treatment of Low-Grade Degenerative Spondylolisthesis. *ScientificWorldJournal*. 2012; 2012:456346.
85. Marotta N, Cosar M, Pimenta L, et al. A novel minimally invasive presacral approach and instrumentation technique for anterior L5 – S1 intervertebral discectomy and fusion: technical description and case presentations. *Neurosurg Focus*. 2006 Jan 15; 20(1):E9.
86. Melgar MA, et al. Segmental and global lordosis changes with two-level axial lumbar interbody fusion and posterior instrumentation. *Int J Spine Surg*. 2014 Dec 1;8. doi: 10.14444/1010. eCollection 2014.
87. Miller LE, Block JE. Minimally invasive arthrodesis for chronic sacroiliac joint dysfunction using the SImmetry SI Joint Fusion system. *Medical Devices: Evidence and Research* 2014;7 125–130.
88. Miller LE, Reckling WC, Block JE. Analysis of postmarket complaints database for the iFuse SI Joint Fusion System®: a minimally invasive treatment for degenerative sacroiliitis and sacroiliac joint disruption. *Med Devices (Auckl)*. 2013 May 29;6:77-84.
89. Nandyala SV, Fineberg SJ, Pelton M, Singh K. Minimally invasive transforaminal lumbar interbody fusion: one surgeon's learning curve. *Spine J*. 2013 Oct 3. pii: S1529-9430(13)01493-9.
90. National Institute for Health and Clinical Excellence (NICE), Lateral (Including Extreme, Extra and Direct Lateral) Interbody Fusion in the Lumbar Spine- Interventional Procedure Guidance 321, Issue Date: November 2009, accessed at [guidance.nice.org.uk](http://guidance.nice.org.uk) 01/28/14.
91. National Institute for Health and Clinical Excellence (NICE). Interventional Procedures Guidance 387, Transaxial interbody lumbosacral fusion. Issue date: March 2011.
92. National Institute for Health and Clinical Excellence (NICE). Transaxial Interbody Lumbosacral Fusion- consultation Document, London UK: NICE; December 2010.
93. National Institute for Health and Care Excellence (NICE). Interventional Procedures Guidance 578: Minimally invasive sacroiliac joint fusion surgery for chronic sacroiliac pain (April 2017).
94. [NeurosurgeryToday.org](http://NeurosurgeryToday.org). Minimally Invasive Spine Surgery (MIS). January 2009.
95. North American Spine Society. Percutaneous Sacroiliac Joint Fusion: Defining Appropriate Coverage Positions. May 31, 2015.
96. North American Spine Society (NASS). Public Education Series. Spinal Fusion. 2006 – 2009 North American Spine Society. Accessed 01/09/12.
97. North American Spine Society Coverage Policy Recommendations: Percutaneous Sacroiliac Joint Fusion (June 2015). Accessed at <https://www.spine.org/>.
98. Official Disability Guidelines (ODG). Medical Treatment and Return to Work Guidelines. Evidence-Based Decision Support: Percutaneous Sacroiliac Joint Fusion.
99. Osman, SG. Endoscopic transforaminal decompression, interbody fusion, and percutaneous pedicle screw implantation of the lumbar spine: A case series report. *Int J Spine Surg*. 2012; 6: 157–166.
100. Ozgur BM, Aryan HE, Pimenta L, Taylor WR. Extreme Lateral Interbody Fusion (XLIF): a novel surgical technique for anterior lumbar Interbody fusion. *Spine J*. 2006 July – August; 6(4): 435-43.

101. Park P, Foley KT. Minimally invasive transforaminal lumbar interbody fusion with reduction of spondylolisthesis: technique and outcomes after a minimum of 2 years' follow-up. *Neurosurg Focus* 25 (2):E16, 2008.
102. Patil SS, Lindley EM, et al. Clinical and Radiological Outcomes of Axial Lumbar Interbody Fusion, Spine Center, Department of Orthopedics, University of Colorado Denver, Colorado, accessed at orthosupersite.com 01/05/11.
103. Petersen T, Laslett M, Juhl C. Clinical classification in low back pain: best-evidence diagnostic rules based on systematic reviews. *BMC Musculoskelet Disord*. 2017 May 12;18(1):188.
104. Polly DW, et al. Does Level of Response to SI Joint Block Predict Response to SI Joint Fusion? *Int J Spine Surg*. 2016 Jan 21;10:4.
105. Polly DW, et al. *Neurosurgery*. 2015 Nov;77(5):674-90; discussion 690-1. Randomized Controlled Trial of Minimally Invasive Sacroiliac Joint Fusion Using Triangular Titanium Implants vs Nonsurgical Management for Sacroiliac Joint Dysfunction: 12-Month Outcomes.
106. Polly DW, et al. Two-Year Outcomes from a Randomized Controlled Trial of Minimally Invasive Sacroiliac Joint Fusion vs. Non-Surgical Management for Sacroiliac Joint Dysfunction. *Int J Spine Surg*. 2016 Aug 23;10:28.
107. Rapp SM, Miller LE, Block JE. AxiaLIF system: minimally invasive device for presacral lumbar interbody spinal fusion. *Med Devices (Auckl)*. 2011;4:125-31.
108. Resnick DK, Choudhri TF, Dailey AT, Groff MW, Khoo L, Matz PG, et al. Guidelines for the performance of fusion procedures for degenerative disease of the lumbar spine. Part 11: interbody techniques for lumbar fusion. *J Neurosurg Spine*. 2005 June;2(6):692-9.
109. Rudolf L. MIS Fusion of the SI Joint: Does Prior Lumbar Spinal Fusion Affect Patient Outcomes? *The open orthopaedics journal* (2013), 7, 163.
110. Rudolf L. Sacroiliac joint arthrodesis-MIS technique with titanium implants: report of the first 50 patients and outcomes. *The open orthopaedics journal* (2012), 6, 495.
111. Rudolph L, Capobianco R. Five-Year Clinical and Radiographic Outcomes After Minimally Invasive Sacroiliac Joint Fusion Using Triangular Implants. *The Open Orthopaedics Journal*, 2014, 8, 375-383.
112. Saavoss JD, Koenig L, Cher DJ. Productivity benefits of minimally invasive surgery in patients with chronic sacroiliac joint dysfunction. *ClinicoEconomics and outcomes research: CEOR*. 2016 Apr 11;8:77.
113. Sachs D, Capobianco R. One year successful outcomes for novel sacroiliac joint arthrodesis system. *Ann Surg Innov Res*. 2012 Dec 27;6(1):13.
114. Sachs D, Capobianco R. Minimally invasive sacroiliac joint fusion: one-year outcomes in 40 patients. *Advances in orthopedics*, 2013.
115. Sachs D, et al. One-year outcomes after minimally invasive sacroiliac joint fusion with a series of triangular implants: a multicenter, patient-level analysis. *Medical Devices: Evidence and Research* 2014;7 299–304.
116. Sachs D, et al. Durable intermediate-to long-term outcomes after minimally invasive transiliac sacroiliac joint fusion using triangular titanium implants. *Med Devices (Auckl)*. 2016 Jul 13;9:213-22.
117. Schroeder JE, Cunningham ME, Ross T, Boachie-Adjei O. Early Results of Sacro-Iliac Joint Fixation Following Long Fusion to the Sacrum in Adult Spine Deformity. *HSS Journal®* (2014) 10(1), 30-35.
118. Shaffrey CI, Smith JS. Editorial: Stabilization of the sacroiliac joint. *Neurosurg Focus (Suppl)* 35:Editorial, 2013.

119. Sharma AK, Kepler CK, Girardi FP, Cammisa FP, Huang RC, Sama AA. Lateral lumbar interbody fusion: clinical and radiographic outcomes at 1 year: a preliminary report. *J Spinal Disord Tech*. 2011 Jun;24(4):242-
120. Shen FH, Samartzis D, Khanna AJ, et al. Minimally invasive techniques for lumbar interbody fusions. *Orthop Clin North Am*. 2007 Jul; 38(3):373-86.
121. Smith AG, Capobianco R, Cher D, Rudolf L, Sachs D, Gundanna M, Shamie AN. Open versus minimally invasive sacroiliac joint fusion: a multi-center comparison of perioperative measures and clinical outcomes. *Ann Surg Innov Res*. 2013 Oct 30;7(1):14. doi: 10.1186/1750-1164-7-14.
122. Society for Minimally Invasive Spine Surgery. Position Statement on Presacral Lumbar Interbody Fusion (February 29, 2012). Accessed at [www.smis.org](http://www.smis.org).
123. Spain K, Holt T. Surgical Revision after Sacroiliac Joint Fixation or Fusion. *Int J Spine Surg*. 2017 Jan 19;11:5.
124. Stark, John G., J. Abner Fuentes, Tania I. Fuentes, and Christopher Idemmili. The history of sacroiliac joint arthrodesis: a critical review and introduction of a new technique. *Current Orthopaedic Practice* 22, no. 6 (2011): 545-557.
125. Stippler M, Turka M, & Gerszten P. C. Outcomes after Percutaneous TranS1 AxiaLIF® L5 – S1 Interbody Fusion for Intractable Lower Back Pain. *The Internet Journal of Spine Surgery*. 2009 Volume 5, Number 1.
126. Stuber KJ. Specificity, sensitivity, and predictive values of clinical tests of the sacroiliac joint: a systematic review of the literature. *J Can Chiropr Assoc*. 2007 Mar;51(1):30-41.
127. Stuesson B, et al. Six-month outcomes from a randomized controlled trial of minimally invasive SI joint fusion with triangular titanium implants vs conservative management. *Eur Spine J*. 2016 May 14.
128. Szadek KM, van der Wurff P, van Tulder MW, Zuurmond WW, Perez RS. Diagnostic validity of criteria for sacroiliac joint pain: a systematic review. *J Pain*. 2009 Apr;10(4):354-68.
129. Tang S. Does TLIF Aggravate Adjacent Segmental Degeneration More Adversely than ALIF? A Finite Element Study. *Turkish Neurosurgery* 2012, Vol: 22, No: 3, 324-328.
130. Tender GC, et al. Percutaneous pedicle screw reduction and axial presacral lumbar interbody fusion for treatment of lumbosacral spondylolisthesis: A case series. *J Med Case Rep*. 2011 Sep 12;5:454. doi: 10.1186/1752-1947-5-454.
131. Tobler WD, Ferrara LA. The presacral retroperitoneal approach for axial lumbar interbody fusion: A prospective study of clinical outcomes, complications and fusion rates at a follow-up of two years in 26 patients. *J Bone Joint Surg [Br]* 2011; 93-B:955-60.
132. Tobler WD, Melgar MA, Raley TJ, Anand N, Miller LE, Nasca RJ. Clinical and radiographic outcomes with L4-S1 axial lumbar interbody fusion (AxiaLIF) and posterior instrumentation: a multicenter study. *Medical Devices (Auckland, NZ)* (2013), 6, 155.
133. Tobler WD, Gerszten PC, Bradley WD, Raley TJ, Nasca RJ, Block JE. Clinical Case Series: Minimally Invasive Axial Presacral L5–S1 Interbody Fusion. Two-Year Clinical and Radiographic Outcomes. *SPINE* Volume 36, Number 20, pp 1–6, September 2011.
134. Tohmeh AG, Watson B, Tohmeh M, Zielinski XJ. Allograft cellular bone matrix in extreme lateral interbody fusion: preliminary radiographic and clinical outcomes. *ScientificWorldJournal*. 2012;2012:263637.
135. Tsahtsarlis A, Wood M. Minimally invasive transforaminal lumbar interbody fusion and degenerative lumbar spine disease. *Eur Spine J*. 2012 Nov;21(11):2300-5.
136. U.S. National Library of Medicine MedlinePlus. Fibromyalgia. Accessed at <https://medlineplus.gov/fibromyalgia.html>.

137. Vanaclocha VV, Verdú-López F, Sánchez-Pardo M, Gozalbes-Esterelles L, Herrera JM. (2014). Minimally Invasive Sacroiliac Joint Arthrodesis: Experience in a Prospective Series with 24 Patients. *J Spine*, 3(185), 2.
138. Vleeming A, Albert HB, Ostgaard HC, Sturesson B, Stuge B. European guidelines for the diagnosis and treatment of pelvic girdle pain. *Eur Spine J*. 2008 Jun;17(6):794-819.
139. Wang X, Xu J, Zhu Y, Li J, Zhou S, Tian S, Xiang Y, Liu X, Zheng Y, Pan T. Biomechanical analysis of a newly developed shape memory alloy hook in a transforaminal lumbar interbody fusion (TLIF) in vitro model. *PLoS One*. 2014 Dec 4;9(12):e114326.
140. WebMD. Somatic Symptom and Related Disorders. Accessed at <https://www.webmd.com/mental-health/somatoform-disorders-symptoms-types-treatment#1>.
141. Whang P, et al. Sacroiliac Joint Fusion Using Triangular Titanium Implants vs. Non-Surgical Management: Six-Month Outcomes from a Prospective Randomized Controlled Trial. *Int J Spine Surg*. 2015 Mar 5;9:6.
142. Whang PG, Sasso RC, Patel VV, Ali RM, Fischgrund JS. Comparison of axial and anterior interbody fusions of the L5-S1 segment: a retrospective cohort analysis. *J Spinal Disord Tech*. 2013 Dec;26(8):437-43.
143. Wong CK, Johnson EK. A narrative review of evidence-based recommendations for the physical examination of the lumbar spine, sacroiliac and hip joint complex. *Musculoskelet. Care* 10 (2012) 149–161.
144. Wu WJ, Liang Y, Zhang XK, Cao P, Zheng T. Complications and clinical outcomes of minimally invasive transforaminal lumbar interbody fusion for the treatment of one- or two-level degenerative disc diseases of the lumbar spine in patients older than 65 years. *Chinese Medical Journal* 2012;125(14):2505-2510.
145. Zeilstra DJ, Miller LE, Block JE. Axial lumbar interbody fusion: a 6-year single-center experience. *Clinical interventions in aging* (2013), 8, 1063.

### **COMMITTEE APPROVAL:**

This Medical Coverage Guideline (MCG) was approved by the Florida Blue Medical Policy & Coverage Committee on 09/27/18.

### **GUIDELINE UPDATE INFORMATION:**

03/15/10	New Medical Coverage Guideline.
02/15/11	Annual review; position statement maintained and references updated.
01/26/12	Scheduled review. Position statement maintained; updated description section and references.
01/01/13	Annual CPT coding update. Added 22586 and 0309T. Revised code descriptors for 0195T and 0196T.
02/15/13	Scheduled review. Position statement maintained. Revised description and updated references.
03/15/14	Scheduled review. Revised description, position statement, program exceptions and CPT coding. Updated references.

12/15/14	Unscheduled review. Revised position statement (coverage for open SI joint fusion). Updated references.
01/01/15	Annual CPT/HCPCS update. Added 27279. Revised 27280 descriptor. Deleted 0334T.
03/15/15	Scheduled review. Position statement maintained. Updated references. Reformatted guideline.
03/15/16	Scheduled review. Position statement maintained. Updated references.
02/15/17	Scheduled review. Position statement maintained. Updated references.
01/01/18	Annual CPT/HCPCS coding update: deleted code 0309T. Reformatted guideline.
02/15/18	Scheduled review. Revised description section. Added coverage criteria for minimally invasive sacroiliac joint fusion/stabilization. Revised definitions and related guidelines sections. Updated references.
06/15/18	Revision: deleted bracing as a requirement for minimally invasive sacroiliac joint fusion. Updated Medicare Advantage program exception and references.
10/15/18	Unscheduled review. Position statement maintained. Revised program exceptions. Updated references.
01/01/19	Annual CPT/HCPCS coding update. Deleted 0195T, 0196T.
01/01/20	Annual CPT/HCPCS coding update. Added C9757.