04-77260-22

Original Effective Date: 06/15/11

Reviewed: 04/24/25

Revised: 05/15/25

Subject: Intensity-Modulated Radiation Therapy (IMRT)

THIS MEDICAL COVERAGE GUIDELINE IS NOT AN AUTHORIZATION, CERTIFICATION, EXPLANATION OF BENEFITS, OR A GUARANTEE OF PAYMENT, NOR DOES IT SUBSTITUTE FOR OR CONSTITUTE MEDICAL ADVICE. ALL MEDICAL DECISIONS ARE SOLELY THE RESPONSIBILITY OF THE PATIENT AND PHYSICIAN. BENEFITS ARE DETERMINED BY THE GROUP CONTRACT, MEMBER BENEFIT BOOKLET, AND/OR INDIVIDUAL SUBSCRIBER CERTIFICATE IN EFFECT AT THE TIME SERVICES WERE RENDERED. THIS MEDICAL COVERAGE GUIDELINE APPLIES TO ALL LINES OF BUSINESS UNLESS OTHERWISE NOTED IN THE PROGRAM EXCEPTIONS SECTION.

Position Statement	Billing/Coding	Reimbursement	Program Exceptions	Definitions	Related Guidelines
<u>Other</u>	References	<u>Updates</u>			

DESCRIPTION:

Intensity-modulated radiation therapy (IMRT) is an advanced mode of high-precision radiotherapy that utilizes computer-controlled x-ray accelerators to deliver precise radiation doses to a malignant tumor or specific areas within the tumor. IMRT allows for the radiation dose to conform more precisely to the three-dimensional (3D) shape of the tumor by modulating or controlling the intensity of the radiation beam. IMRT also allows higher radiation doses to be focused to regions within the tumor while minimizing the dose to surrounding normal critical structures. Treatment is planned by using 3D computed tomography (CT) images of the patient in conjunction with computerized dose calculations to determine the dose intensity pattern that will best conform to the tumor shape. IMRT treatment may be delivered using several delivery methods, including, for example multiple static segment treatment (Step-and-shoot), dynamic segment treatment (sliding window), binary-collimator tomotherapy and intensity-modulated arc techniques.

IMRT is complex and requires precision and accuracy, and involves multiple medical specialists (e.g., radiation oncologist, medical physicist, radiation therapists, dosimetrist). IMRT requires multiple or fractionated treatment sessions and different radiation doses. Several factors determine the number of treatment sessions and radiation dose; the type, location and size of the tumor, doses to critical normal structures and the individual's health. An IMRT schedule may consist of five days a week for five to ten weeks. At the beginning of each treatment, the individual is positioned on the treatment table guided by the marks on the skin defining the treatment area; the individual may be repositioned during the IMRT treatment. Imaging systems on the IMRT treatment delivery systems may be used to check positioning and marker location, molded devices may be used to help the individual maintain proper position. IMRT treatment may take between 10 and 30 minutes.

There are several radiation therapy devices in use for IMRT that have received U.S. Food and Drug Administration (FDA) 510(k) clearance to market, including the TiGRT TPS (Radiation Treatment Planning System, Innocure Intensity Modulating Radiation Therapy Compensators, micro-Multi-Leaf Collimator and the Genesis[™] IMRT Medical Linear Accelerator.

Summary of Evidence: Intensity-modulated radiation therapy (IMRT) for abdomen and pelvic conditions, breast cancer, tumors of the central nervous system, head and neck cancer, lung cancer and prostate cancer, the evidence includes retrospective studies, comparative studies, evaluation studies, multi-center studies and practice guidelines (Chuong 2013; Fuller 2007, Caglar 2007; Graff 2007; Appel 2019; Choi 2005; Murshed 2004; Cahlon 2008; Zaorsky 2017; Bai 2003; Beriwal 2007; Bhatnaga 2004; 2006; Freedman 2009; Czito 2009;Harkenrider 2023; Shaitelman 2024).

POSITION STATEMENT:

Intensity modulated radiation therapy (IMRT) **meets the definition of medical necessity** for the following indications.

Abdomen and Pelvis

Intensity-modulated radiotherapy **meets the definition of medical necessity** for members with cancer of the anus and anal canal.

Intensity-modulated radiotherapy **meets the definition of medical necessity** for the treatment of cancer of the abdomen and pelvis, including but not limited to the following when dosimetric planning with standard 3-dimensional conformal radiotherapy predicts that the radiation dose to an adjacent organ would result in unacceptable normal tissue toxicity:

- Esophageal cancer
- Gastrointestinal cancer (e.g., cholangiocarcinoma, gastric, liver, pancreas)
- Gynecologic tumors (e.g., cervical, endometrial, uterine, vaginal, vulvar)
- Hepatobiliary tract
- Rectal locations.

Intensity-modulated radiotherapy is considered **experimental or investigational** for all other indications for abdomen and pelvis. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

Breast Cancer

Intensity-modulated radiotherapy (IMRT) **meets the definition of medical necessity** as a technique to deliver whole-breast irradiation in members receiving treatment for left-sided breast cancer after breast-conserving surgery when **ALL** the following conditions have been met:

- Cardiac radiation exposure cannot be avoided using alternative radiotherapy; AND
- IMRT dosimetry demonstrates significantly reduced cardiac target volume radiation exposure.

IMRT **meets the definition of medical necessity** in members with large breasts when treatment planning with 3-dimensional conformal radiotherapy results in hot spots (focal regions with dose variation > 10% of target) and the hot spots can be avoided with IMRT.

Central Nervous System

Intensity-modulated radiotherapy **meets the definition of medical necessity** for members with malignant or benign brain tumors when the tumor is proximate to organs at risk (e.g., brain stem, spinal cord, cochlea and eye structures including optic nerve and chiasm, lens and retina) and 3-dimensional conformal radiotherapy planning is not able to meet dose-volume constraints for normal tissue tolerance.

Hippocampal-avoiding intensity-modulated radiotherapy **meets the definition of medical necessity** for members with brain tumor metastases outside a 5-mm margin around either hippocampus and expected survival is \geq 4 months.

Intensity-modulated radiotherapy is considered **experimental or investigational** for the treatment of tumors of the central nervous system not meeting the above criteria. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Head and Neck Cancers

Intensity-modulated radiotherapy **meets the definition of medical necessity** for the treatment of head and neck cancers.

Intensity-modulated radiotherapy **meets the definition of medical necessity** for the treatment of thyroid cancers in close proximity to organs at risk (e.g., esophagus, salivary glands, spinal cord) and 3-dimensional conformal radiotherapy planning is not able to meet dose volume constraints for normal tissue tolerance.

Lung Cancer

Intensity-modulated radiotherapy (IMRT) **meets the definition of medical necessity** as a technique to deliver radiotherapy in members with lung cancer when **ALL** of the following conditions are met:

- IMRT dosimetry demonstrates a reduction in the V20 to at least 10% below the V20 that is achieved with the 3-dimensional plan (e.g., from 40% down to 30% or lower);
- Radiotherapy is being given with curative intent; AND
- Three-dimensional conformal radiotherapy will expose > 35% of normal lung tissue to more than a 20-gray (Gy) dose-volume (V20).

Prostate Cancer

Intensity-modulated radiotherapy **meets the definition of medical necessity** in the treatment of localized prostate cancer.

Intensity-modulated radiotherapy **meets the definition of medical necessity** after radical prostatectomy for the following indications:

- Adjuvant therapy when there are adverse pathologic findings at prostatectomy or with a persistently detectable prostate-specific antigen level after prostatectomy; **OR**
- Salvage therapy when there is evidence of biochemical or local recurrence when there is no evidence of distant metastatic disease.

Intensity-modulated radiotherapy **meets the definition of medical necessity** for the following indications:

- Bladder cancer (primary non-metastatic bladder carcinoma and treatment intent is curative)
- Head and neck cancers (e.g., lymphomas, glottic cancer (stage III and IV))
- Mediastinal tumors (e.g., lymphomas, thymoma, thymic carcinoma)
- Pancreatic cancer
- Pediatric members (below age 18) (radiosensitive tumors, retinoblastoma)
- Sarcoma (pelvic sarcoma, retroperitoneal sarcoma, extremity sarcoma)
- Skin cancer (head and neck region when regional lymphatics will be treated, cutaneous melanoma)
- Thoracic malignancies
- Thyroid cancer (anaplastic, node-positive, node-recurrent).

Intensity modulated radiotherapy to treat a previously irradiated field **meets the definition of medical necessity** for the following indications:

- Benign brain lesion
- Benign brain tumors (acoustic neuromas, craniopharyngiomas, pineal gland tumors, schwannomas)
- Bone metastases
- Breast cancer
- Central nervous system tumor (medulloblastoma, supratentorial primitive neuroectodermal tumors (PNET), ependymoma)
- CNS lymphoma
- Colorectal cancer and anal cancer
- Gastrointestinal cancers (cholangiocarcioma, esophageal, gastric)
- Genitourinary cancers (bladder, penile, testicular)
- Glioma (high grade. low grade)
- Gynecologic cancers (cervical, fallopian tube, ovarian, uterine, vulvar, vaginal)
- Head and neck cancers (including thyroid)
- Hepatocellular carcinoma

- Intracranial arteriovenous malformations
- Liver metastases
- Lymphoma (Hodgkin and Non-Hodgkin)
- Meningioma
- Metastatic brain lesion
- Metastatic lesions in the lung
- Pancreatic cancer
- Pituitary adenomas
- Primary lung cancers (non-small cell, small cell, mestatic lung lesions)
- Prostate cancer (low, intermediate and high risk of recurrence, post-prostatectomy)
- Retinoblastoma
- Sarcoma
- Skin cancer
- Spine lesions (primary or metastatic)
- Thymoma and thymic carcinoma
- Trigeminal neuralgia
- Uveal melanoma.

BILLING/CODING INFORMATION:

Note: Procedure and diagnosis codes may not be all inclusive.

CP	Т	Co	di	ng	-
----	---	----	----	----	---

77301	Intensity modulated radiotherapy plan, including dose-volume histograms for
	target and critical structure partial tolerance specifications
77338	Multi-leaf collimator (MLC) device(s) for intensity modulated radiation
	therapy (IMRT), design and construction per IMRT plan
77385	Intensity modulated radiation treatment delivery (IMRT), includes guidance
	and tracking, when performed; simple
77386	Intensity modulated radiation treatment delivery (IMRT), includes guidance
	and tracking, when performed; complex

HCPCS Coding:

G6015	Intensity modulated treatment delivery, single or multiple fields/arcs,via
	narrow spatially and temporally modulated beams, binary, dynamic mlc, per
	treatment session

G6016	Compensator-based beam modulation treatment delivery of inverse planned
	treatment using 3 or more high resolution (milled or cast) compensator,
	convergent beam modulated fields, per treatment session

ICD-10 Diagnosis Codes That Support Medical Necessity:

C00.0 – C00.9	Malignant neoplasm of lip
C01	Malignant neoplasm of base of tongue
C02.0 – C02.9	Malignant neoplasm of other and unspecified parts of tongue
C03.0 – C03.9	Malignant neoplasm of gum
C04.0 - C04.9	Malignant neoplasm of floor of mouth
C06.0 – C06.2	Malignant neoplasm of other and unspecified parts of mouth
C06.80 – C06.89	Malignant neoplasm of overlapping sites of other and unspecified parts of
	mouth
C06.9	Malignant neoplasm of mouth , unspecified
C07	Malignant neoplasm of parotid gland
C08.0 – C08.9	Malignant neoplasm of other and unspecified major salivary glands
C09.0 – C09.9	Malignant neoplasm of tonsil
C10.0 – C10.9	Malignant neoplasm of oropharynx
C11.0 - C11.9	Malignant neoplasm of nasopharynx
C12	Malignant neoplasm of pyriform sinus
C13.0 – C13.9	Malignant neoplasm of hypopharynx
C14.0 – C14.8	Malignant neoplasm of other and ill-defined sites in the lip, oral cavity and
	pharynx
C15.3 – C15.9	Malignant neoplasm of esophagus
C16.0 – C16.9	Malignant neoplasm of stomach
C18.0 – C18.9	Malignant neoplasm of colon
C19	Malignant neoplasm of rectosigmoid junction
C20	Malignant neoplasm of rectum
C21.0 – C21.8	Malignant neoplasm of anus and anal canal
C22.0	Liver cell carcinoma
C22.1	Intrahepatic bile duct carcinoma
C24.0	Malignant neoplasm of extrahepatic bile duct
C25.0 – C25.9	Malignant neoplasm of pancreas
C30.0, C30.1	Malignant neoplasm of nasal cavity and middle ear
C31.0 – C31.9	Malignant neoplasm of accessory sinuses
C32.0 – C32.9	Malignant neoplasm of larynx
C34.00 – C34.02	Malignant neoplasm of main bronchus
C34.10 – C34.12	Malignant neoplasm of upper lobe, bronchus or lung
C34.2	Malignant neoplasm of middle lob, right bronchus or lung
C34.30 – C34.32	Malignant neoplasm of lower lobe, bronchus or lung
C34.80 – C34.82	Malignant neoplasm of overlapping sites of bronchus and lung
C34.90 – C34.92	Malignant neoplasm of bronchus or lung, unspecified
C37	Malignant neoplasm of thymus

C41.2	Malignant neoplasm of vertebral column
C43.0-C43.9	Malignant melanoma
C44.0-C44.99	Other and unspecified malignant neoplasm of skin
C48.0-C48.8	Malignant neoplasm of retroperitoneum and peritoneum
C49.10-C49.12	Malignant neoplasm of connective and soft tissue of upper limb, including
	shoulder
C49.20-C49.22	Malignant neoplasm of connective and soft tissue of lower limb, including hip
C49.4	Malignant neoplasm of connective and soft tissue of abdomen
C49.5	Malignant neoplasm of connective and soft tissue of pelvis
C50.011 – C50.019	Malignant neoplasm of nipple and areola, female
C50.021 – C50.029	Malignant neoplasm of nipple and areola, male
C50.111 – C50.119	Malignant neoplasm of central portion of breast, female
C50.121 – C50.129	Malignant neoplasm of central portion of breast, male
C50.211 – C50.219	Malignant neoplasm of upper-inner quadrant of breast, female
C50.221 – C50.229	Malignant neoplasm of upper-inner quadrant of breast, male
C50.311 – C50.319	Malignant neoplasm of lower-inner quadrant of breast, female
C50.321 – C50.329	Malignant neoplasm of lower-inner quadrant of breast, male
C50.411 – C50.419	Malignant neoplasm of upper-outer quadrant of breast, female
C50.421 – C50.429	Malignant neoplasm of upper-outer quadrant of breast, male
C50.511 – C50.519	Malignant neoplasm of lower-outer quadrant of breast, female
C50.521 – C50.529	Malignant neoplasm of lower-outer quadrant of breast, male
C50.611 – C50.619	Malignant neoplasm of axillary tail of breast, female
C50.621 – C50.629	Malignant neoplasm of axillary tail of breast, male
C50.811 – C50.819	Malignant neoplasm of overlapping sites of breast, female
C50.821 – C50.829	Malignant neoplasm of overlapping sites of breast, male
C50.911 – C50.919	Malignant neoplasm of breast of unspecified site, female
C50.921 – C50.929	Malignant neoplasm of breast of unspecified site, male
C51.0 – C51.9	Malignant neoplasm of vulva
C52	Malignant neoplasm of vagina
C53.0 – C53.9	Malignant neoplasm of cervix uteri
C54.0 – C54.9	Malignant neoplasm of corpus uteri
C55	Malignant neoplasm of uterus, part unspecified
C56.1 – C56.9	Malignant neoplasm of ovary
C57.00 – C57.02	Malignant neoplasm of fallopian tube
C57.7	Malignant neoplasm of other specified female genital organs
C57.8	Malignant neoplasm of overlapping sites of female genital organs
C57.9	Malignant neoplasm of female genital organ, unspecified
C60.0 – C60.9	Malignant neoplasm of penis
C62.00 - C62.02	Malignant neoplasm of undescended testis
C62.10 - C62.12	Malignant neoplasm of descended testis
C62.90 – C62.92	Malignant neoplasm of testis, unspecified whether descended or
	undescended
C61	Malignant neoplasm of prostate

C67.0 – C67.9	Malignant neoplasm of bladder
C69.00 - C69.02	Malignant neoplasm of conjunctiva
C69.10 - C69.12	Malignant neoplasm of cornea
C69.20 – C69.22	Malignant neoplasm of retina
C69.30 – C69.32	Malignant neoplasm of choroid
C69.40 – C69.42	Malignant neoplasm of ciliary body
C69.50 – C69.52	Malignant neoplasm of lacrimal gland and duct
C69.60 – C69.62	Malignant neoplasm of orbit
C69.80 - C69.82	Malignant neoplasm of overlapping sites of eye and adnexa
C69.90 – C69.92	Malignant neoplasm of eye, unspecified
C71.0 – C71.9	Malignant neoplasm of brain
C73	Malignant neoplasm of thyroid gland
C76.0	Malignant neoplasm of head, face and neck
C78.00 – C78.02	Secondary malignant neoplasm of lung
C78.7	Secondary malignant neoplasm of liver and intrahepatic bile duct
C79.31 – C79.32	Secondary malignant neoplasm of brain and cerebral meninges
C79.40 – C79.49	Secondary malignant neoplasm of other and unspecified parts of nervous
	system
C79.51 – C79.52	Secondary malignant neoplasm of bone and bone marrow
C79.81	Secondary malignant neoplasm of breast
C79.82	Secondary malignant neoplasm of genital organs
C80.1	Malignant (primary) neoplasm, unspecified
C81.02	Nodular lymphocyte predominant Hodgkin lymphoma, intrathoracic lymph
	nodes
C81.12	Nodular sclerosis Hodgkin lymphoma, intrathoracic lymph nodes
C81.22	Mixed cellularity Hodgkin lymphoma, intrathoracic lymph nodes
C81.32	Lymphocyte depleted Hodgkin lymphoma, intrathoracic lymph nodes
C81.42	Lymphocyte-rich Hodgkin lymphoma, intrathoracic lymph nodes
C81.72	Other Hodgkin lymphoma, intrathoracic lymph nodes
C81.92	Hodgkin lymphoma, unspecified, intrathoracic lymph nodes
C82.02	Follicular lymphoma grade I, intrathoracic lymph nodes
C82.12	Follicular lymphoma grade II, intrathoracic lymph nodes
C82.22	Follicular lymphoma grade III, unspecified, intrathoracic lymph nodes
C82.32	Follicular lymphoma grade IIIa, intrathoracic lymph nodes
C82.42	Follicular lymphoma grade IIIb, intrathoracic lymph nodes
C82.52	Diffuse follicle center lymphoma, intrathoracic lymph nodes
C82.62	Cutaneous follicle center lymphoma, intrathoracic lymph nodes
C82.82	Other types of follicular lymphoma, intrathoracic lymph nodes
C82.92	Follicular lymphoma, unspecified, intrathoracic lymph nodes
C83.02	Small cell B-cell lymphoma, intrathoracic lymph nodes
C83.12	Mantle cell lymphoma, intrathoracic lymph nodes
C83.32	Diffuse large B-cell lymphoma, intrathoracic lymph nodes
C83.52	Lymphoblastic (diffuse) lymphoma, intrathoracic lymph nodes

C83.72	Burkitt lymphoma, intrathoracic lymph nodes
C83.82	Other non-follicular lymphoma, intrathoracic lymph nodes
C83.92	Non-follicular (diffuse) lymphoma, unspecified, intrathoracic lymph nodes
C84.02	Mycosis fungoides, intrathoracic lymph nodes
C84.12	Sézary disease, intrathoracic lymph nodes
C84.42	Peripheral T-cell lymphoma, not classified, intrathoracic lymph nodes
C84.62	Anaplastic large cell lymphoma, ALK-positive, intrathoracic lymph nodes
C84.72	Anaplastic large cell lymphoma, ALK-negative, intrathoracic lymph nodes
C84.92	Mature T/NK-cell lymphomas, unspecified, intrathoracic lymph nodes
C84.A2	Cutaneous T-cell lymphoma, unspecified, intrathoracic lymph nodes
C84.Z2	Other mature T/NK-cell lymphomas, intrathoracic lymph nodes
C85.12	Unspecified B-cell lymphoma, intrathoracic lymph nodes
C85.22	Mediastinal (thymic) large B-cell lymphoma, intrathoracic lymph nodes
C85.82	Other specified types of non-Hodgkin lymphoma, intrathoracic lymph nodes
C85.92	Non-Hodgkin lymphoma, unspecified, intrathoracic lymph nodes
C85.81	Other specified types of non-Hodgkin lymphoma, lymph nodes of head, face,
	and neck
C85.81	Other specified types of non-Hodgkin lymphoma, lymph nodes of head, face,
	and neck
D05.00 - D05.02	Lobular carcinoma in situ of breast
D05.10 - D05.12	Intraductal carcinoma in situ of breast
D05.80 - D05.82	Other specified type of carcinoma in situ of breast
D05.90 - D05.92	Unspecified type of carcinoma in situ of breast
D15.0	Benign neoplasm of thymus
D33.0 – D33.2	Benign neoplasm of brain and other parts of central nervous system
D35.2	Benign neoplasm of pituitary gland
D35.4	Benign neoplasm of pineal gland
G50.0	Trigeminal neuralgia
Q28.2	Arteriovenous malformation of cerebral vessels
R97.2	Elevated prostate specific antigen [PSA]
Z08	Encounter for follow-up examination after completed treatment for
	malignant neoplasm
Z51.0	Malignant neoplasm of vertebral column
Z51.5	Encounter for palliative care
Z92.3	Personal history of irradiation
Z85.820	Personal history of malignant melanoma of skin
Z98.89	Other specified postprocedural states

LOINC Codes:

The following information may be required documentation to support medical necessity: physician history and physical, physician progress notes, plan of treatment and reason for intensity-modulated radiation therapy (IMRT).

DOCUMENTATION	LOINC	LOINC TIME	LOINC TIME FRAME MODIFIER CODES
TABLE	CODES	FRAME MODIFIER	NARRATIVE
		CODE	
Physician history and	28626-0	18805-2	Include all data of the selected type that
physical			represents observations made six months
			or fewer before starting date of service
			for the claim
Attending physician	18741-9	18805-2	Include all data of the selected type that
progress note			represents observations made six months
			or fewer before starting date of service
			for the claim
Plan of treatment	18776-5	18805-2	Include all data of the selected type that
			represents observations made six months
			or fewer before starting date of service
			for the claim
Physician history and	28626-0	18805-2	Include all data of the selected type that
physical			represents observations made six months
			or fewer before starting date of service
			for the claim

REIMBURSEMENT INFORMATION:

Refer to section entitled **POSITION STATEMENT**.

PROGRAM EXCEPTIONS:

Federal Employee Program (FEP): Follow FEP guidelines.

State Account Organization (SAO): Follow SAO guidelines.

Medicare Advantage products: No National Coverage Determination (NCD) and/or Local Coverage Determination (LCD) were found at the time of the last guideline reviewed date.

If this Medical Coverage Guideline contains a step therapy requirement, in compliance with Florida law 627.42393, members or providers may request a step therapy protocol exemption to this requirement if based on medical necessity. The process for requesting a protocol exemption can be found at <u>Coverage</u> <u>Protocol Exemption Request</u>.

DEFINITIONS:

Active surveillance: ongoing monitoring, exams and tests may be done to monitor.

Tomotherapy IMRT: a type of imaged-guided IMRT.

3D conformal radiation therapy (radiotherapy) (**3D-CRT**): a form of <u>external beam therapy</u> that more precisely conforms the radiation to the tumor, allowing a higher radiation dose to be safely delivered.

3D planning (3D plan): treatment plan.

RELATED GUIDELINES:

Image-Guided Radiation Therapy for Treatment Planning and Delivery, 04–77260–19

OTHER:

Other names used to describe intensity modulated radiation therapy (IMRT):

Compensatory-based IMRT Helical Tomotherapy IMRT Tomotherapy Inverse Treatment Planning Segment Radiation Therapy (RT) Sliding Window Technique Step-and-Shoot Inverse IMRT

REFERENCES:

- 1. Abrams RA, winter KA, Regine WF et al. Failure to adhere to protocol specified radiation therapy guidelines was associated with decreased survival in RTOG 9704--a phase III trial of adjuvant chemotherapy and chemoradiotherapy for patients with resected adenocarcinoma of the pancreas. International Journal of Oncology, Biology, Physics 2012; 82(2): 809-16.
- 2. Anal Carcinoma National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology. Version 2.2011, 02/02/11.
- 3. American Society for Radiation Oncology (ASTRO) Quantitative Analysis of Normal Tissue Effects in the Clinic (QUANTEC), accessed 2016.
- 4. Apisarnthanarax S, Barry A, Cao M, et al. External Beam Radiation Therapy for Primary Liver Cancers: An ASTRO Clinical Practice Guideline. Pract Radiat Oncol. 2022 Jan-Feb;12(1):28-51.
- Appel S, Bar J, Ben-Nun A, Perelman M, et al. Comparative effectiveness of intensity modulated radiation therapy to 3-dimensional conformal radiation in locally advanced lung cancer: pathological and clinical outcomes. Br J Radiol. 2019 May;92(1097):20180.
- Bai YR, Wu GH, Guo WJ et al. Intensity modulated radiation therapy and chemotherapy for locally advanced pancreatic cancer: results of feasibility study. World Journal of Gastroenterology 2003; 9(11): 2561-2564.
- 7. Beriwal S, Gan GN, Heron DE et al. Early clinical outcome with concurrent chemotherapy and extend-field, intensity-modulated radiotherapy for cervical cancer. International Journal Radiation Oncology Biology Physics 2007; 68(1): 166-171.
- 8. Bhatia A, Rao A, Ang KK et al. Anaplastic thyroid cancer: Clinical outcomes with conformal radiotherapy. Head Neck 2010; 32(7): 829-836.
- 9. Bhatnagar AK, Brandner E, Sonnik D, et al. Intensity-modulated radiation therapy (IMRT) reduces the dose to the contralateral breast when compared to conventional tangential fields for primary breast irradiation: initial report. The Cancer Journal 2004; 10(6): 381-385.
- 10. Bhatnagar AK, Brandner E, Sonnik D, et al. Intensity modulated radiation therapy (IMRT) reduces the dose to the contralateral breast when compared to conventional tangential fields for primary breast irradiation. Breast Cancer Research and Treatment 2006; 96(1): 41-46.
- 11. Blue Cross Blue Shield Association Evidence Positioning System®. 8.01.49 Intensity-Modulated Radiotherapy: Abdomen, Pelvis and Chest, 08/24.

- 12. Blue Cross Blue Shield Association Evidence Positioning System®. 8.01.46 Intensity-Modulated Radiotherapy of the Breast and Lung, 08/24
- 13. Blue Cross Blue Shield Association Evidence Positioning System®. 8.01.59 Intensity-Modulated Radiotherapy: Central Nervous System Tumors, 08/24 (Arcchived).
- 14. Blue Cross Blue Shield Association Evidence Positioning System®. 8.01.48 Intensity-Modulated Radiotherapy of the Head and Neck or Thyroid,
- 15. Blue Cross Blue Shield Association Evidence Positioning System®. 8.01.47 Intensity-Modulated Radiotherapy of the Prostate, 08/24 (Archived).
- Cabrera AR, Kirkpatrick JP, Fiveash JB et al. Radiation therapy for glioblastoma: Executive summary of an American Society for Radiation Oncology Evidence-Based Clinical Practice Guideline. Practical Radiation Oncology 2016; 6(4): 217-225.
- 17. Caglar HB, Allen AM. Intensity-modulated radiotherapy for head and neck cancer. Clinical Advances in Hematology & Oncology 2007; 5(6): 425-431.
- 18. Cahlon O, Hunt M, Zelefsky MJ. Intensity-modulated radiation therapy: supportive data for prostate cancer. Seminars in Radiation Oncology 2008; 18(1): 48-57.
- Choi Y, Kim JK, Lee HS et al. Impact of intensity-modulated radiation therapy as a boost treatment on the lung-dose distributions for non-small-cell lung cancer. International Journal Radiation Oncology Biology Physics 2005; 63(3): 683-689.
- Christian JA, Bedford JL, Webb S, Brada M. Comparison of inverse-planned three-dimensional conformal radiotherapy and intensity-modulated radiotherapy for non-small-cell lung cancer. International Journal Radiation Oncology Biology Physics 2007; 67(3): 735-741.
- Chuong MD, Freilich JM, Hoffe SE et al. Intensity-Modulated Radiation Therapy vs. 3D Conformal Radiation Therapy for Squamous Cell Carcinoma of the Anal Canal. Gastrointestinal Cancer Research: GCR 2013; 6(2): 39-45.
- 22. Czito BG, Pepek JM, Meyer JJ et al. Intensity-modulated radiation therapy for anal cancer. Oncology 2009; 23(12):1082-1089.
- 23. Das IJ, Cheng CW, Chopra KL et al. Intensity-modulated radiation therapy dose prescription, recording, and delivery: patterns of variability among institutions and treatment planning systems. Journal of the National Cancer Institute 2008; 100(5): 300-307.
- 24. Dawson LA, Jaffray DA. Advances in image-guided radiation therapy. Journal of Clinical Oncology 2007; 25(8): 938-946.
- 25. De Meerleer G O, Vakaet L A, De Gersem WR, et al. Radiotherapy of prostate cancer with or without intensity modulated beams: a planning comparison. International Journal Radiation Oncology Biology Physiology 2000; 47(3): 639-648.
- 26. Ding M, Newman F, Raben D. New radiation therapy techniques for the treatment of head and neck cancer. Otolaryngologic Clinics of North America 2005; 38:371-395.
- Donovan E, Bleakly N, Denholm E et al. Randomised trial of standard 3D radiotherapy (RT) versus intensity modulated radiotherapy (IMRT) in patients prescribed breast radiotherapy. Radiotherapy and Oncology 2007; 82(3): 254-264.
- Eccles CL, Bissonnette JP, Craig T et al. Treatment planning study to determine potential benefit of intensity-modulated radiotherapy versus conformal radiotherapy for unresectable hepatic malignancies. International Journal Radiation Oncology Biology Physics 2008; 72(2): 582-588.
- 29. Fang FM, Tsai WL, Chen HC et al. Intensity-modulated or conformal radiotherapy improves the quality of life of patients with nasopharyngeal carcinoma: comparisons of four radiotherapy techniques. Cancer 2007; 109(2): 313-321.

- 30. Fenwick JD, Riley SW, Scott AJ. Advances in intensity-modulated radiotherapy delivery. Cancer Treatment Research 2008; 139: 193-214.
- Freedman GM, Li T, Nicolaou N et al. Breast intensity-modulated radiation therapy reduces time spent with acute dermatitis for women of all breast sizes during radiation. International Journal Radiation Oncology Biology Physics 2009; 74(3): 689-694.
- Freedman GM, Anderson PR, Li J et al. Intensity modulated therapy (IMRT) decreases acute skin toxicity for women receiving radiation for breast cancer. American Journal Clinical Oncology 2006; 29(1): 66-70.
- 33. Fuller CD, Choi M, Forthuber B et al. Standard fractionation intensity modulated radiation therapy (IMRT) of primary and recurrent glioblastoma multiforme. Radiation Oncology 2007; 2(26).
- 34. Girinsky T, Pichenot C, Beaudre A, et al. Is intensity-modulated radiotherapy better than conventional radiation treatment and three-dimensional conformal radiotherapy for mediastinal masses in patients with Hodgkin's disease, and is there a role for beam orientation optimization and dose constraints assigned to virtual volumes? International Journal Radiation Oncology Biology Physics 2006; 64(1): 218-226.
- 35. Gondi V, Pugh SL, Tome WA et al. Preservation of memory with conformal avoidance of the hippocampal neural stem-cell compartment during whole-brain radiotherapy for brain metastases (RTOG 0933): a phase II multi-institutional trial. Journal of Clinical Oncology 2014; 32(34):3810-3816.
- 36. Gong Y, Wang J, Bai S et al. Conventionally-fractionated image-guided intensity modulated radiotherapy (IG-IMRT): a safe and effective treatment for cancer spinal metastasis. Radiation Oncology 2008; 3:11.
- 37. Graff P, Lapeyre M, Desandes E et al. Impact of intensity-modulated radioatherapy on health-related quality of life for head and neck cancer patients: matched-pair comparison with conventional radiotherapy. International Journal Radiation Oncology Biology Physics 2007; 67(5): 1309-1317.
- Guerrero Urbano MT, Henrys AJ, Adams EJ et al. Intensity-modulated radiotherapy in patients with locally advanced rectal cancer reduces volume of bowel treated to high dose levels. International Journal Radiation Oncology Biology Physics 2006; 65(3): 907-916.
- 39. Gupta V, McGunigal M, Prasad-Hayes M et al. Adjuvant radiation therapy is associated with improved overall survival in high-intermediate risk stage I endometrial cancer: A national cancer data base analysis. Gynecologic Oncology 2017; 144 (1):119–124.
- 40. Haffty BG. Should intensity-modulated radiation therapy be the standard of care in the conservatively managed breast cancer patient? Journal of clinical Oncology 2008; 26(13): 2072-2074.
- Harkenrider MM, Abu-Rustum N, Albuquerque K, et al. Radiation Therapy for Endometrial Cancer: An American Society for Radiation Oncology Clinical Practice Guideline. Pract Radiat Oncol. 2023 Jan-Feb;13(1):41-65.
- 42. Hatano K, Araki H, Sakai M, et al. Current status of intensity-modulated radiation therapy (IMRT). International Journal of Clinical Oncology 2007; 12(6): 408-415.
- 43. luchi T, Hatano K, Narita Y et al. Hypofractionated high-dose irradiation for the treatment of malignant astrocytomas using simultaneous integrated boost technique by IMRT. International Journal Radiation Oncology Biology Physics 2006; 64(5): 1317-1324.
- 44. James ND, Jenkins P, Tremlett J et al. Radiotherapy with or without Chemotherapy in Muscle-Invasive Bladder Cancer. New England Journal of Medicine 2012; 366(16): 1477-1488.
- 45. Kam MK, Leung SF, Zee B et al. Prospective randomized study of intensity-modulated radiotherapy on salivary gland function in early-stage nasopharyngeal carcinoma patients. Journal of Clinical Oncology 2007; 25(31): 4873-4879.
- 46. Kaufman DS, Winter KA, Shipley WU et al. The initial results in muscle-invading bladder cancer of RTOG 95-06: phase I/II trial of transurethral surgery plus radiation therapy with concurrent cisplatin

and 5-fluorouracil followed by selective bladder preservation or cystectomy depending on the initial response. Oncologist 2000; 5(6): 471-476.

- 47. Kavanagh BD, Schefter TE, Wu, O, et al. Clinical application of intensity-modulated radiotherapy for locally advanced cervical cancer. Seminars in Radiation Oncology 2002; 12(3): 260-271.
- Kestin LL, Sharpe MB, Frazier RC et al. Intensity modulation to improve dose uniformity with tangential breast radiotherapy: initial clinical experience. International Journal Radiation Oncology Biology Physics 200; 48(5): 1559-1568.
- 49. Kruger, EA, Fraas BA, Pierce LJ. Clinical aspects of intensity-modulated radiotherapy in the treatment of breast cancer. Seminars Radiation Oncology 2002; 12(3): 250-259.
- 50. Liao ZX, Komaki RR, Thames HD et al. Influence of technologic advances on outcomes in patients with unresectable, locally advanced non-small-cell lung cancer receiving concomitant chemoradiotherapy. International Journal Radiation Oncology Biology Physics 2010; 76(3): 775-781.
- Laskar S, Bahl G, Muckaden M et al. Nasopharyngeal carcinoma in children: comparison of conventional and intensity-modulated radiotherapy. International Journal Radiation Oncology Biology Physics 2008; 72(3): 728-736
- 52. Lee NY, Terezakis SA. Intensity-modulated radiation therapy. Journal of Surgical Oncology 2008; 97(8): 691-696.
- Leonard C, Carter D, Kercher J et al. Prospective trial of accelerated partial breast intensitymodulated radiotherapy. International Journal Radiation Oncology Biology Physics 2007; 67(5): 1291-1298.
- 54. Lips I, Dehnad H, Kruger AB et al. Health-related quality of life in patients with locally advanced prostate cancer after 76 Gy intensity-modulated radiotherapy vs. 70 Gy conformal radiotherapy in a prospective and longitudinal study. International Journal Radiation Oncology Biology Physiology 2007; 69(3): 656-661.
- 55. MacDonald SM, Ahmad S, Kachris S et al. Intensity modulated radiation therapy versus threedimensional conformal radiation therapy for the treatment of high grade glioma: a dosimetric comparison. Journal of Applied Clinical Medical Physics 2007; 8(2): 47-60.
- 56. Macdonald JS, Smalley SR, Benedetti J et al. Chemoradiotherapy after surgery compared with surgery alone for adenocarcinoma of the stomach or gastroesophageal junction. New England Journal of Medicine 2001; 345(10): 725-730.
- 57. Mak RH, Hunt D, Shipley WU et al. Long-term outcomes in patients with muscle-invasive bladder cancer after selective bladder-preserving combined-modality therapy: a pooled analysis of Radiation Therapy Oncology Group protocols 8802, 8903, 9506, 9706, 9906, and 0233. Journal of Clinical Oncology 2014; 32(34):3801-3809.
- Meattini I, Marrazzo L, Saieva C, et al.Accelerated Partial-Breast Irradiation Compared With Whole-Breast Irradiation for Early Breast Cancer: Long-Term Results of the Randomized Phase III APBI-IMRT-Florence Trial. J Clin Oncol. 2020 Dec 10;38(35):4175-4183.
- 59. Milano MT, Chmura SJ, Garofalo MC, et al. Intensity-modulated radiotherapy in treatment of pancreatic and bile duct malignancies: toxicity and clinical outcome. International Journal Radiation Oncology Biology Physics 2004; 2004; 59(2): 445-453.
- 60. Miller TP, Dahlberg S, Cassady JR et al. Chemotherapy alone compared with chemotherapy plus radiotherapy for localized intermediate- and high-grade non-Hodgkin's lymphoma. New England Journal of Medicine 1998; 339(1): 21-26.
- Murshed H, Liu HH, Liao Z et al. Dose and volume reduction for normal lung using intensitymodulated radiotherapy for advanced-stage non-small-cell lung cancer. International Journal Radiation Oncology Biology Physics 2004; 58(4): 1258-1267.

- 62. National Cancer Institute Retinoblastoma Treatment (PDQ[®])–Health Professional Version, Updated August 9, 2022.
- 63. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Breast Cancer. Version 4.2022.
- 64. National Comprehensive Cancer Network. Clinical Practice Guidelines in Oncology (NCCN). NCCN Clinical Practice Guidelines in Oncology. Colon Cancer, Version 2.2017.
- 65. National Comprehensive Cancer Network. Clinical Practice Guidelines in Oncology (NCCN). NCCN Clinical Practice Guidelines in Oncology. Esophageal and Esophagogastric Junction Cancers, Version 2.2017.
- 66. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Melanoma: Cutaneous, Version 1.2023.
- 67. National Comprehensive Cancer Network. Clinical Practice Guidelines in Oncology (NCCN). NCCN Clinical Practice Guidelines in Oncology. Ovarian Cancer Including Fallopian Tube Cancer and Primary Peritoneal Cancer, Version 2.2017.
- 68. Pawlik TM, Ahuja N, Herman JM. The role of radiation in retroperitoneal sarcomas: a surgical perspective. Current Opinion in Oncology 2007; 17(4): 359-366.
- 69. Penagaricano JA, Papanikolaou N, Yan Y, et al. Application of intensity-modulated radiation therapy for pediatric malignancies. Medical Dosimetry 2004; 29(4): 247-253.
- 70. Pepek JM, Willett CG, Czito BG. Radiation therapy advances for treatment of anal cancer. Journal of the National Comprehensive Cancer Network 2010 Jan; 8(1): 123-129.
- 71. Pignol JP, Olivotto I, Rakovitch E, et al. A multicenter randomized trial of breast intensity-modulated radiation therapy to reduce acute radiation dermatitis. Journal of Clinical Oncology 2008; 26(13): 2085-2092.
- 72. Ren F, Li S, Zhang Y, et al. Efficacy and safety of intensity-modulated radiation therapy versus threedimensional conformal radiation treatment for patients with gastric cancer: a systematic review and meta-analysis. Radiat Oncol. 2019 May 22;14(1):84.
- Salma JK, Mell LK, Schomas DA et al. Concurrent chemotherapy and intensity-modulated radiation therapy for anal canal cancer patients: a multicenter experience. Journal of Clinical Oncology 2007; 25(29): 4581-4586.
- Santanam L, Malinowski K, Hubenshmidt J et al. Fiducial-based translational localization accuracy of electromagnetic tracking system and on-board kilovoltage imaging system. International Journal Radiation Oncology Biology Physics 2008; 70(3): 892-799.
- 75. Shaitelman SF, Anderson BM, Arthur DW, et al. Partial Breast Irradiation for Patients with Early-Stage Invasive Breast Cancer or Ductal Carcinoma In Situ: An ASTRO Clinical Practice Guideline. Pract Radiat Oncol. 2024 Mar-Apr;14(2):112-132.
- Smalley SR, Benedetti JK, Haller DG et al. Updated analysis of SWOG-directed intergroup study 0116: a phase III trial of adjuvant radiochemotherapy versus observation after curative gastric cancer resection. Journal of Clinical Oncology 30(19): 2327-2333.
- 77. Stehman FB, Ali S, Keys HM et al. Radiation therapy with or without weekly cisplatin for bulky stage 1B cervical carcinoma: follow-up of a Gynecologic Oncology Group trial. American Journal of Obstetrics and Gynecology 2007; 197(5): 503.e1–503.e6.
- Straus DJ, Portlock CS, Qin J et al. Results of a prospective randomized clinical trial of doxorubicin, bleomycin, vinblastine, and dacarbazine (ABVD) followed by radiation therapy (RT) versus ABVD alone for stages I, II, and IIIA nonbulky Hodgkin disease. Blood 2004; 104 (12): 3483-3489.

- Sura S, Gupta V, Yorke E et al. Intensity-modulated radiation therapy (IMRT) for inoperable non-small cell lung cancer: the Memorial Sloan-Kettering Cancer Center (MSKCC) experience. Radiotherapy Oncology 2008; 87(1): 17-23.
- 80. Teh BS, Mai WY, Uhl BM et al. Intensity-modulated radiation therapy (IMRT) for prostate cancer with the use of a rectal balloon for prostate immobilization: acute toxicity and dose-volume analysis. International Journal Radiation Oncology Biology Physics 2001; 49(3): 705-712.
- 81. Tiong SS, Dickie C, Haas RL et al. The role of radiotherapy in the management of localized soft tissue sarcomas. Cancer Biology & Medicine 2016; 13(3): 373-383.
- Trofimov A, Nguyen PL, Coen JJ et al. Radiotherapy treatment of early-stage prostate cancer with IMRT and protons: a treatment planning comparison. International Journal Radiation Oncology Biology Physiology 2007; 69(4): 444-453.
- 83. U.S. Food and Drug Administration (FDA) 510(k) Summary-TiGRT TPS Radiation Treatment Planning System K090893, 07/15/09.
- 84. U.S. Food and Drug Administration (FDA) 510(k) Summary-Innocure Intensity Modulating Radiation Therapy Compensators K052383, 03/29/06.
- 85. U.S. Food and Drug Administration (FDA) 510(k) Summary-micro-Multi-Leaf Collimator K020860, 06/06/02.
- 86. U.S. Food and Drug Administration (FDA) 510(k) Summary-Genesis™ IMRT Medical Linear Accelerator K982502, 07/31/98.
- Vaidya JS, Wenz F, Bulsara et al. Risk-adapted targeted intraoperative radiotherapy versus wholebreast radiotherapy for breast cancer: 5-year results for local control and overall survival from the TARGIT-A randomised trial. Lancet 2014; 383 (9917): 603-613.
- 88. Veldeman L, Madani I, Hulstaert F et al. Evidence behind use of intensity-modulated radiotherapy: a systematic review of comparative clinical studies. Lancet Oncology 2008; 9(4): 367-375.
- Wang SJ, Choi M, Fuller CD et al. Intensity-modulated radiotherapy for patients with brain metastases: a mature outcomes analysis. Technology in Cancer Research and Treatment 2007; 6(3): 161-166.
- Wang D, Zhang Q, Eisenberg BL et al. Significant Reduction of Late Toxicities in Patients With Extremity Sarcoma Treated With Image-Guided Radiation Therapy to a Reduced Target Volume: Results of Radiation Therapy Oncology Group RTOG-0630 Trial. Journal of Clinical Oncology 2015; 33(20): 2231-2238.
- 91. Williams PC. IMRT: delivery techniques and quality assurance. The British Journal of Radiology 2003; 76:766-776.
- 92. Zagar TM, Willett CG, Czito BG. Intensity-Modulated Radiation Therapy for Anal Cancer: Toxicity versus Outcomes. Oncology (Williston Park) 2010; 24(9): 815-823.
- Zaorsky NG, Showalter TN, Ezzell GA, et al. ACR Appropriateness Criteria for external beam radiation therapy treatment planning for clinically localized prostate cancer, part II of II. Adv Radiat Oncol. 2017 Mar 20;2(3):437-454.
- 94. Zelefsky MJ, Fuks Z, Happensett L, et al. Clinical experience with intensity-modulated radiation therapy (IMRT) in prostate cancer. Radiotherapy and Oncology 2000; 55: 241-249.
- Zelefsky MJ, Fuks Z, Hunt M, et al. High dose radiation delivered by intensity modulated conformal radiotherapy improves the outcomes of localized prostate cancer. The Journal of Urology 2001; 166(3): 876-881.
- Zelefsky MJ, Fuks Z, Hunt M, et al. High-dose intensity modulated radiation therapy for prostate cancer: early toxicity and biochemical outcome in 772 patients. International Journal Radiation Oncology Biology Physics 2002b; 53(5): 1111-1116.

97. Zelefsky MJ, Fuks Z, Leibel SA. Intensity-modulated radiation therapy for prostate cancer. Seminars Radiation Oncology 2002a; 12:229-237.

COMMITTEE APPROVAL:

This Medical Coverage Guideline (MCG) was approved by the Florida Blue Medical Policy & Coverage Committee on 04/24/25.

GUIDELINE UPDATE INFORMATION:

06/15/11	New Medical Coverage Guideline.
08/15/11	Revision; formatting changes.
05/11/14	Revision: Program Exceptions section updated.
01/01/15	Annual HCPCS code update. Deleted 77418, 0073T. Added 77385, 77386, G6015 and
	G6016.
10/01/15	Revision; updated ICD9 and ICD10 coding section.
11/01/15	Revision: ICD-9 Codes deleted.
05/01/16	Added/revised indications: bone metastases, breast cancer, CNS cancers, intracranial
	lesions, medulloblastoma, supratentorial, PNET, ependymoma, CNS lymphoma,
	metastatic brain lesions, benign brain lesions, pituitary adenomas, meningioma, other
	benign brain tumors (acoustic neuromas, carniopharyngiomas, pineal gland tumors,
	schwannomas), ocular lesions, retinoblastoma, spine lesions (primary or metastatic),
	other neurologic indication (trigeminal neuralgia), colorectal and anal cancers,
	gastrointestinal cancers, non-colorectal (cholangiocarcinoma, esophageal, gastric, liver
	and pancreatic), genitourinary cancers (bladder, penile and testicular), gynecologic
	cancers (cervical, fallopian tube, ovarian, uterine, and vulvar/vaginal), head and neck
	cancers (including thyroid cancer), lung cancer (small cell and non-small cell), other
	tumor types (including sarcoma, pediatrics, and other malignancies), and prostate
	cancer; added 77280, 72285, 77290, 77295, 77338, 77402, 77407 and 77412; updated
	ICD-10 codes; updated program exception; updated references.
06/02/16	Code update; change code C50.519 in code range to C50.811-C50.819.
08/15/16	Updated program exceptions.
11/15/16	Revision; revised position statement. Added 77424, 77425 and 77469. Updated
	references.
01/01/17	Annual HCPCS code update. Revised 77402, 77407 and 77412 code descriptor.
10/15/17	Revision; revised position statement. Added Hodgkin and Non-Hodgkin lymphoma,
	thyoma and thymic carcinoma. Revised other malignancies. Updated references.
02/15/18	Revision; updated position statement, ICD-10 diagnoses codes, program exceptions and
	definitions.
03/15/21	Review/revision. Bone metastases: Revised retreatment with EBRT, deleted spinal cord
	and added adjacent organ. Breast cancer: Revised dose-volume constraints criteria.
	Central nervous system: Revised high grade and low grade gliomas criteria. Revised
	metastatic brain lesions criteria. Rectal cancer: Deleted criteria. Lymphoma: Hodgkin and
	Non-Hodgkin lymphoma: Added lymphomas of the head and neck region. Head and neck
	cancers: Revised criteria, deleted laryngeal and added glottic. Added "advanced" to head

	and neck cancers. Added lymphomas of the head and neck region. Primary lung cancers:
	non-small cell lung cancer: Added stage I and II disease to adjuvant or definitive
	treatment. Revised dose-volume constraints. Added adjuvant or definitive treatment of
	stage III disease. Added fractionation and criteria. Primary lung cancers: small cell lung
	cancer: Revised dose-volume constraints. Sarcoma: Revised extremity sarcoma, deleted
	prior to surgery, to spare a joint. Added radiosensitive tumor in children. Prostate
	cancer: risk of recurrence (low, intermediate, high): Revised statement for clarity.
	Prostate cancer: Revised local recurrence criteria. Added fractionation and criteria.
	Added skin cancer and criteria. Updated ICD-10 codes and references.
05/15/23	Review: revised position statement. Updated coding and references.
12/15/23	Updated related guidelines.
05/15/25	Review; no change in position statement. Updated references.