

05-86000-24

Original Effective Date: 03/15/05

Reviewed: 12/05/19

Revised: 01/15/20

Subject: Laboratory Tests for Heart and Kidney Transplant Rejection

THIS MEDICAL COVERAGE GUIDELINE IS NOT AN AUTHORIZATION, CERTIFICATION, EXPLANATION OF BENEFITS, OR A GUARANTEE OF PAYMENT, NOR DOES IT SUBSTITUTE FOR OR CONSTITUTE MEDICAL ADVICE. ALL MEDICAL DECISIONS ARE SOLELY THE RESPONSIBILITY OF THE PATIENT AND PHYSICIAN. BENEFITS ARE DETERMINED BY THE GROUP CONTRACT, MEMBER BENEFIT BOOKLET, AND/OR INDIVIDUAL SUBSCRIBER CERTIFICATE IN EFFECT AT THE TIME SERVICES WERE RENDERED. THIS MEDICAL COVERAGE GUIDELINE APPLIES TO ALL LINES OF BUSINESS UNLESS OTHERWISE NOTED IN THE PROGRAM EXCEPTIONS SECTION.

[Position Statement](#)

[Billing/Coding](#)

[Reimbursement](#)

[Program Exceptions](#)

[Definitions](#)

[Related Guidelines](#)

[Other](#)

[References](#)

[Updates](#)

DESCRIPTION:

Acute cellular rejection is most likely to occur in the first six months after transplantation, with a significant decline in the incidence of rejection after this time. Although immunosuppressants are required on a life-long basis, dosing is adjusted based on graft function and the grade of acute cellular rejection determined by histopathology. Endomyocardial biopsies are typically taken periodically during the first 6 to 12 months post-transplant. The interval between biopsies varies among clinical centers. A typical schedule is weekly for the first month, once or twice monthly for the following six months, and several times (monthly to quarterly) between six months and one year post-transplant. Surveillance biopsies may also be performed after the first postoperative year (eg, on a quarterly or semiannual basis). This practice, although common, has not been demonstrated to improve transplant outcomes. Some centers no longer routinely perform endomyocardial biopsies after one year in patients who are clinically stable.

While the endomyocardial biopsy is the criterion standard for assessing heart transplant rejection, it is limited by a high degree of interobserver variability in the grading of results and potential morbidity that can occur with the biopsy procedure. Also, the severity of rejection may not always coincide with the grading of the rejection by biopsy. Finally, a biopsy cannot be used to identify patients at risk of rejection, limiting the ability to initiate therapy to interrupt the development of rejection. For these reasons, an endomyocardial biopsy is considered a flawed criterion standard by many.

Noninvasive methods of detecting cellular rejection have been explored with the hope that noninvasive tests will assist in determining appropriate patient management and avoid overuse or underuse of treatment with steroids and other immunosuppressants that can occur with false-negative and false-positive biopsy reports.

Several commercially available laboratory tests assess heart transplant rejection, including the Heartsbreath test, which measures breath markers of oxidative stress, and the AlloMap test, which uses gene expression profiling. These tests create a score based on the expression of a variety of immunomodulatory genes and are proposed as an alternative or as an adjunct to invasive endomyocardial biopsy. Renal transplant rejection may be assessed by the AlloSure test, which measures the donor-derived cell-free DNA in peripheral blood and is proposed as an alternative or as an adjunct to invasive renal biopsy. A protein biomarker, soluble suppression of tumorigenicity-2 (sST2), has also elicited interest as a prognostic marker post heart transplantation and as a test to predict acute cellular rejection (graft-versus-host disease).

POSITION STATEMENT:

AlloMap® molecular expression testing **meets the definition of medical necessity** as a non-invasive method of determining the risk of rejection in heart transplant recipients (15 years or older) who are between 6 months and 5 years post-transplant.

AlloMap molecular expression testing is considered **experimental or investigational** for all other indications. The evidence is insufficient to determine the effects of the technology on health outcomes.

The use of all other molecular expression tests in the management of members after heart transplantation is considered **experimental or investigational**. The evidence is insufficient to determine the effects of the technology on health outcomes.

The measurement of volatile organic compounds (e.g. Heartsbreath® test) to assist in the detection of moderate grade 2R (formerly grade 3) heart transplant rejection is considered **experimental or investigational**. The evidence is insufficient to determine the effects of the technology on health outcomes.

The use of peripheral blood measurement of donor-derived cell-free DNA in the management of members after renal transplantation, including but not limited to the detection of acute renal transplant rejection or renal transplant graft dysfunction (e.g. AlloSure®) is considered **experimental or investigational**. The evidence is insufficient to determine the effects of the technology on health outcomes.

The use of the Presage® ST2 assay is considered **experimental or investigational** for all indications including, but not limited to, predicting prognosis and predicting acute cellular rejection in the post cardiac transplantation period. The evidence is insufficient to determine the effects of the technology on health outcomes.

BILLING/CODING INFORMATION:

CPT Coding:

81595	Cardiology (heart transplant), mRNA, gene expression profiling by real-time quantitative PCR of 20 genes (11 content and 9 housekeeping), utilizing subfraction of peripheral blood, algorithm reported as a rejection risk score
83006	Growth stimulation expressed gene 2 (ST2, Interleukin 1 receptor like-1) (Investigational)
0085T	Breath test for heart transplant rejection (Investigational)
0055U	Cardiology (heart transplant), cell-free DNA, PCR assay of 96 DNA target sequences (94 single nucleotide polymorphism targets and two control targets), plasma

	(Investigational)
0087U	Cardiology (heart transplant), mRNA gene expression profiling by microarray of 1283 genes, transplant biopsy tissue, allograft rejection and injury algorithm reported as a probability score (Investigational)
0088U	Transplantation medicine (kidney allograft rejection) microarray gene expression profiling of 1494 genes, utilizing transplant biopsy tissue, algorithm reported as a probability score for rejection (Investigational)
0118U	Transplantation medicine, quantification of donor-derived cell-free DNA using whole genome next-generation sequencing, plasma, reported as percentage of donor-derived cell-free DNA in the total cell-free DNA (Investigational)

ICD-10 Diagnosis Codes That Support Medical Necessity for 81595:

T86.20-T86.298	Complications of heart transplant
Z48.21	Encounter for aftercare following heart transplant
Z94.1	Heart transplant status

REIMBURSEMENT INFORMATION:

Refer to section entitled [POSITION STATEMENT](#).

PROGRAM EXCEPTIONS:

Federal Employee Program (FEP): Follow FEP guidelines.

State Account Organization (SAO): Follow SAO guidelines.

Medicare Advantage Products:

The following National Coverage Determination (NCD) was reviewed on the last guideline reviewed date: Heartsbreath Test for Heart Transplant Rejection (260.10) located at cms.gov.

The following was reviewed on the last guideline reviewed date and located at palmettogba.com:

- Local Coverage Article:MoIDX:AlloMap Billing and Coding Guidelines Update (A53099)
- Local Coverage Determination (LCD):MoIDX: AlloSure® Donor-Derived Cell-Free DNA Test (L37266).

DEFINITIONS:

No guideline specific definitions apply.

RELATED GUIDELINES:

None applicable.

OTHER:

None applicable.

REFERENCES:

1. Bloom RD, Bromberg JS, et al. Cell-Free DNA and Active Rejection in Kidney Allografts. *J Am Soc Nephrol.* 2017 Jul;28(7):2221-2232.
2. Blue Cross Blue Shield Association Medical Policy Reference Manual. 2.01.68 Laboratory Tests for Heart and Kidney Transplant Rejection, 11/19.
3. Blue Cross Blue Shield Association Medical Policy Reference Manual. 2.04.130 ST2 Assay for Chronic Heart Failure and Heart Transplant Rejection, 06/19.
4. Blue Cross Blue Shield Association Technology Evaluation Center (TEC). Gene expression profiling as a noninvasive method to monitor for cardiac allograft rejection. *TEC Assessments 2011*; volume 26, tab 8.
5. Centers for Medicare & Medicaid Services (CMS), National Coverage Determination (NCD) for Heartsbreath Test for Heart Transplant Rejection (260.10); accessed at cms.org.
6. ClinicalTrials.gov. Evaluation of Patient Outcomes From the Kidney Allograft Outcomes AlloSure Registry, sponsored by CareDx; accessed 10/14/19.
7. ClinicalTrials.gov. Outcomes AlloMap Registry Study: the Clinical Long-term Management and Outcomes of Heart Transplant Recipients With Regular Rejection Surveillance Including Use of AlloMap Gene-expression Profiling Testing, sponsored by CareDx; accessed 10/14/19.
8. ClinicalTrials.gov. Utility of Donor-Derived Cell-free DNA in Association With Gene-Expression Profiling (AlloMap®) in Heart Transplant Recipients (D-OAR), sponsored by CareDx; accessed 10/14/19.
9. Costanzo MR, Dipchand A, Starling R et al. The International Society of Heart and Lung Transplantation Guidelines for the care of heart transplant recipients. *J Heart Lung Transplant* 2010; 29(8):914-56 accessed at isHLT.org.
10. Crespo-Leiro MG, Stypmann J, Schulz U, et al. Clinical usefulness of gene-expression profile to rule out acute rejection after heart transplantation: CARGO II. *Eur Heart J.* Jan 7 2016.
11. Deng MC, Eisen HJ, Mehra MR, et al., Noninvasive Discrimination of Rejection in Cardiac Allograft Recipients Using Gene Expression Profiling, *American Journal of Transplantation*, Vol 6, Issue 1, page 150-160, January 2006.
12. Deng MC, Elashoff B, Pham MX et al. Utility of Gene Expression Profiling Score Variability to Predict Clinical Events in Heart Transplant Recipients. *Transplantation* 2014 Mar 27; 97(6): 708–714.
13. Fang, K, Clinical Utilities of Peripheral blood Gene Expression Profiling in the Management of Cardiac Transplant Patients, *Journal of Immunotoxicology*, Vol 4, July 2007, pages 209-217.
14. Fujita B, Proashovikj F, et al. Predictive value of gene expression profiling for long-term survival after heart transplantation. *Transpl Immunol.* 2017 Mar;41:27-31. doi: 10.1016/j.trim.2017.02.001. Epub 2017 Feb 4. PMID: 28167272.
15. JA T. Gene Expression Profiling for the Diagnosis of Heart Transplant Rejection. *California Technology Assessment Forum.* 2010, accessed at ctaf.org 07/18/12.
16. Jordan SC, Bunnapradist S, et al. Donor-derived Cell-free DNA Identifies Antibody-mediated Rejection in Donor Specific Antibody Positive Kidney Transplant Recipients. *Transplant Direct.* 2018 Aug 20;4(9):e379.
17. Loupy A, Woodward RN, et al. Distinct Patterns of Gene Expression Identified Longitudinally and within AlloMap Score Ranges are Associated with Clinical Outcomes. *Journal of Heart and Lung Transplantation*, Vol. 38, Issue 4, Supplement, April 2019 Page S51.
18. Mehra MR, Uber PA, Genomic Biomarkers and Heart Transplantation, *Heart Fail Clin.* 2007 Jan; 3(1): 83-6.

19. Moayed Y, Fouroutan F, et al. Risk evaluation using gene expression screening to monitor for acute cellular rejection in heart transplant recipients. J Heart Lung Transplant. 2019 Jan;38(1):51-58. doi: 10.1016/j.healun.2018.09.004.Epub 2018 Sep 12. PMID: 30352779.
20. Nikolaidis LA, Borzognia B, Bove AA, Limitations of Molecular Expression Profiling (Allomap Score) in Predicting Endomyocardial biopsy Findings in Cardiac Allograft Recipients: Two Year Follow-Up Surveillance Study, Circulation. October 2007; 116:II_505.
21. Palmetto GBA, Local Coverage Article:MoIDX: AlloMap Billing and Coding Guidelines Update (A53099);accessed at palmettogba.com.
22. Palmetto GBA, Local Coverage Determination (LCD):MoIDX: AlloSure® Donor-Derived Cell-Free DNA Test (L37266); accessed at palmettogba.com.
23. Pham MX, Deng MC, Kfoury AG, et al, Molecular Testing for Long-Term Rejection surveillance in Heart Transplant Recipients: Design of the Invasive Monitoring Attenuation Through Gene Expression (IMAGE) trial, J Heart Lung Transplant. 2007 Aug; 26(8): 808-14.
24. Stewart S, Winters GL, Fishbein MC, et al. Revision of the 1990 working formulation for the standardization of nomenclature in the diagnosis of heart rejection. J Heart Lung Transplant. Nov 2005;24(11):1710-1720; accessed at isHLT.org.
25. TAI Diagnostics - myTAI_{HEART} Test Dossier, October 2019.
26. U.S. Food and Drug Administration (FDA), accessed at fda.gov.
27. Yamani MH, Taylor DO, Haire C, Smedira N, Starling RC, Post-Transplant Ischemic Injury is Associated With Up-regulated AlloMap Gene Expression, Clin Transplant. 2007 Jul-Aug; 21(4): 523-5.
28. Yamani MH, Taylor DO, Rodriguez ER, et al, Transplant Vasculopathy is Associated with Increased AlloMap Gene Expression Score, J Heart Lung Transplant. 2007 Apr; 26(4): 403-6.
29. Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol. Oct 15 2013;62(16):e147-239.

COMMITTEE APPROVAL:

This Medical Coverage Guideline (MCG) was approved by the Florida Blue Medical Policy & Coverage Committee on 12/05/19.

GUIDELINE UPDATE INFORMATION:

03/15/04	New Medical Coverage Guideline.
03/15/06	Annual review; continue investigational status.
03/15/07	Scheduled review; no change in coverage statement; references.
06/15/07	Reformatted guideline.
11/15/07	Revision: title changed, description section updated, position statement updated, Medicare Advantage section updated, references updated.
02/15/08	Annual review: position statements maintained; description section updated; references updated.
02/15/09	Annual review: position statements maintained; description section and references updated.
12/15/09	Annual review: position statements maintained; description section and references updated.
09/15/12	Review; position statements maintained; program exceptions section and references

	updated.
06/15/13	Annual review; position statements maintained and references updated.
06/15/14	Annual review; position statements maintained, program exception and reference updated.
06/15/15	Annual review; position statements maintained and references updated.
01/01/16	Annual HCPCS/CPT update; code 81595 added.
07/15/16	Annual review; revise the position statement section, coding, program exception, and references.
12/15/17	Annual review; investigational position maintained; description, position statements, and references updated.
12/15/18	Annual review; Investigational position maintained; investigational statement for AlloSure test added; title, description, coding, and references updated.
07/01/19	Quarterly CPT/HCPCS update. Added codes 0087U & 0088U.
01/15/20	Review; position statements, coding, description, and references updated.