

05-86000-26

Original Effective Date: 10/15/05

Reviewed: 01/28/21

Revised: 01/01/22

Subject: Assays of Genetic Expression in Tumor Tissue as a Technique to Determine Prognosis in Patients with Breast Cancer

THIS MEDICAL COVERAGE GUIDELINE IS NOT AN AUTHORIZATION, CERTIFICATION, EXPLANATION OF BENEFITS, OR A GUARANTEE OF PAYMENT, NOR DOES IT SUBSTITUTE FOR OR CONSTITUTE MEDICAL ADVICE. ALL MEDICAL DECISIONS ARE SOLELY THE RESPONSIBILITY OF THE PATIENT AND PHYSICIAN. BENEFITS ARE DETERMINED BY THE GROUP CONTRACT, MEMBER BENEFIT BOOKLET, AND/OR INDIVIDUAL SUBSCRIBER CERTIFICATE IN EFFECT AT THE TIME SERVICES WERE RENDERED. THIS MEDICAL COVERAGE GUIDELINE APPLIES TO ALL LINES OF BUSINESS UNLESS OTHERWISE NOTED IN THE PROGRAM EXCEPTIONS SECTION.

[Position Statement](#)

[Billing/Coding](#)

[Reimbursement](#)

[Program
Exceptions](#)

[Definitions](#)

[Related Guidelines](#)

[Other](#)

[References](#)

[Updates](#)

DESCRIPTION:

An important part of treatment planning for patients with breast cancer involves determining which patients could benefit from adjuvant cytotoxic chemotherapy. For example, for patients with early-stage, invasive breast cancer (ie, cancer extending beyond the basement membrane of the mammary ducts into adjacent tissue), adjuvant cytotoxic chemotherapy consistently provides approximately a 30% relative risk reduction in 10-year breast cancer mortality regardless of patients' baseline prognosis. However, the absolute benefit of chemotherapy depends on the underlying or baseline risk of recurrence. Patients with the best prognosis have tumors that are small, early-stage, estrogen receptor–positive, and are lymph node–negative. Patients may have received no adjuvant treatment, or adjuvant tamoxifen and/or adjuvant chemotherapy. These patients have an approximately 15% 10-year risk of recurrence with tamoxifen alone, which means that approximately 85% of these patients could avoid the toxicity of adjuvant cytotoxic chemotherapy if they could be accurately identified. Conventional risk classifiers estimate recurrence risk by considering criteria such as tumor size, type, grade, and histologic characteristics; hormone receptor status; and number of affected lymph nodes. Consensus guidelines for defining receptor status exist, however, no single classifier is considered a criterion standard. As a result, more patients are treated with chemotherapy who fail to benefit. Better predictors of recurrence risk could help patient's decision making, some who may prefer to avoid chemotherapy if assured that their risk is low.

Laboratory tests have been developed that detect the expression, via messenger RNA, of different genes in breast tumor tissue and combine the results to determine prognosis in patients with breast cancer. Test

results may help providers and patients decide whether to include adjuvant chemotherapy in the postsurgical management of breast cancer, to alter treatment in patients with ductal carcinoma in situ (DCIS), or triple-negative (estrogen receptor, progesterone receptor, human epidermal growth factor receptor 2) breast cancer (TNBC), or to recommend extended endocrine therapy in patients who are recurrence-free at 5 years.

POSITION STATEMENT:

The use of the 21-gene reverse transcriptase-polymerase chain reaction (PT-PCR) assay (i.e., Oncotype DX[®]) to determine recurrence risk for deciding whether to undergo adjuvant chemotherapy **meets the definition of medical necessity** in members with newly diagnosed (6 months or less) primary, invasive breast cancer meeting **ALL** of the following criteria:

- Unilateral tumor;
- Node-negative (lymph nodes with [micrometastases](#) (≤ 2 mm in size) are considered node negative) **OR** with 1-3 involved ipsilateral axillary lymph nodes;
- Hormone receptor-positive ([ER-positive](#) or [PR-positive](#));
- [HER2-negative](#);
- Tumor size > 0.5 cm;
- Who have had surgery and full pathological examination of the tumor completed;
- The test result will determine if adjuvant chemotherapy or hormonal therapy will be used; **AND**
- The test is ordered by the treating physician.

Use of EndoPredict[™], the Breast Cancer Index[™], MammaPrint[®], and Prosigna[™] (also known as PAM50) to determine recurrence risk for deciding whether to undergo adjuvant chemotherapy **meets the definition of medical necessity** in members with primary, invasive breast cancer that meet the above criteria for for Oncotype DX.

The use of the MammaPrint assay to determine recurrence risk for deciding whether to undergo adjuvant chemotherapy **meets the definition of medical necessity** in members with primary, invasive breast cancer meeting **ALL** of the following characteristics:

- Unilateral tumor;
- 1-3 positive nodes;
- Hormone receptor-positive (ER-positive or PR-positive);
- HER2-negative;
- Stage T1, T2, or operable T3 at high clinical risk of recurrence*;
- The test result will determine if adjuvant chemotherapy or hormonal therapy will be used;
- Who have had surgery and full pathological examination of the tumor completed;
- When ordered within 6 months after diagnosis; **AND**
- The test is ordered by the treating physician.

*Tumor Grade	Nodes	Tumor Size
Well differentiated	1-3	≤2 cm or 2.1-5 cm
Moderately differentiated	1-3	Any size
Poorly differentiated or undifferentiated	1-3	Any size

The use of the Breast Cancer Index (BCI) assay to assist in decision of extending adjuvant hormonal therapy beyond 5 years of treatment **meets the definition of medical necessity** for members currently receiving adjuvant hormonal therapy and meeting **ALL** of the following criteria:

- Diagnosed with early-stage breast cancer
- Hormone receptor-positive (ER-positive or PR-positive);
- HER2-negative;
- Node-negative (lymph nodes with micrometastases (≤ 2 mm in size) are considered node negative) OR with 1-3 involved ipsilateral axillary lymph nodes;
- No evidence of distant breast cancer metastasis (i.e., non-relapsed);
- The test result will guide the decision regarding extended adjuvant hormonal therapy; AND
- The test is ordered by the treating physician.

All other indications for the 21-gene PT-PCR assay (i.e., Oncotype DX), EndoPredict, the Breast Cancer Index, MammaPrint, and Prosigna are considered **experimental or investigational**. There is a lack of clinical data in peer-reviewed literature to permit conclusions on safety, efficacy and net health outcomes.

Use of a subset of genes from the 21-gene RT-PCR assay for predicting recurrence risk in members with noninvasive ductal carcinoma in situ (i.e., Oncotype DX[®] DCIS Score) to inform treatment planning following excisional surgery is considered **experimental or investigational**. The evidence is insufficient to permit conclusions on efficacy and net health outcomes.

All other gene expression assays (e.g., Mammostrat[®], BreastOncPx[™], Insight[®] DX Breast Cancer Profile, NexCourse[®] Breast IHC4, BreastPRS[™], MapQuant Dx[™], and BreastOncPx[™]) are considered **experimental or investigational**. The evidence is insufficient to permit conclusions on efficacy and net health outcomes.

The use of Insight TNBCtype to aid in making decisions regarding chemotherapy in members with triple-negative breast cancer is considered **experimental or investigational**. The evidence is insufficient to permit conclusions on efficacy and net health outcomes.

The following are considered **experimental or investigational** as the evidence is insufficient to permit conclusions on efficacy and net health outcomes:

- The use of gene expression assays to molecularly subclassify breast cancer (eg, Blueprint[®])
- The use of gene expression assays for quantitative assessment of ER, PR and HER2 overexpression (eg, TargetPrint[®])
- The use of testing to predict response to specific chemotherapy regimens

- Repeat testing, use of more than one test for the same tumor, or testing of multiple tumor sites in the same member.

BILLING/CODING INFORMATION:

CPT Coding:

81518	Oncology (breast), mRNA, gene expression profiling by real-time RT-PCR of 11 genes (7 content and 4 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithms reported as percentage risk for metastatic recurrence and likelihood of benefit from extended endocrine therapy (Breast Cancer Index)
81519	Oncology (breast), mRNA, gene expression profiling by real-time RT-PCR of 21 genes, utilizing formalin-fixed paraffin embedded tissue, algorithm reported as recurrence score (Oncotype DX)
81520	Oncology (breast), mRNA gene expression profiling by hybrid capture of 58 genes (50 content and 8 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a recurrence risk score (Prosigna)
81521	Oncology (breast), mRNA, microarray gene expression profiling of 70 content genes and 465 housekeeping genes, utilizing fresh frozen or formalin-fixed paraffin-embedded tissue, algorithm reported as index related to risk of distant metastasis (MammaPrint)
81522	Oncology (breast), mRNA, gene expression profiling by RT-PCR of 12 genes (8 content and 4 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as recurrence risk score (EndoPredict)
81523	Oncology (breast), mRNA, next-generation sequencing gene expression profiling of 70 content genes and 31 housekeeping genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as index related to risk to distant metastasis
0045U	Oncology (breast ductal carcinoma in situ), mRNA, gene expression profiling by real-time RT-PCR of 12 genes (7 content and 5 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as recurrence score (Investigational) (DCIS Score)
0153U	Oncology (breast), mRNA, gene expression profiling by next-generation sequencing of 101 genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a triple negative breast cancer clinical subtype(s) with information on immune cell involvement (Investigational)

HCPCS Coding:

S3854	Gene expression profiling panel for use in the management of breast cancer treatment
-------	--

ICD-10 Diagnosis Codes That Support Medical Necessity:

C50.011-C50.929	Malignant neoplasm of breast
D05.00-D05.92	Lobular carcinoma in situ of breast
Z17.0	Estrogen receptor positive status [ER+]

REIMBURSEMENT INFORMATION:

Reimbursement is subject to post-service medical review. The following information is required documentation to support medical necessity: physician history and physical, pathology report, treating physician visit notes that include documentation that the intention to treat or not treat with adjuvant chemotherapy was contingent, at least in part, on the results of the test for the individual patient in question.

LOINC Codes:

DOCUMENTATION TABLE	LOINC CODES	LOINC TIME FRAME MODIFIER CODE	LOINC TIME FRAME MODIFIER CODES NARRATIVE
Physician history and physical	28626-0	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim
Attending physician visit note	18733-6	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.
Pathology Reports Sections	26439-0	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.
Pathology Study Reports	27898-6	18805-2	Include all data of the selected type that represents observations made six months or fewer before starting date of service for the claim.

PROGRAM EXCEPTIONS:

Federal Employee Program (FEP): Follow FEP guidelines.

State Account Organization (SAO): Follow SAO guidelines.

Medicare Advantage Products:

The following were reviewed on the last guideline reviewed date: MoIDX LCDs located at palmettogba.com.

DEFINITIONS:

Adjuvant chemotherapy: chemotherapy given in addition to surgical therapy, in order to reduce the risk of local or systemic relapse.

ER-positive (estrogen receptor positive): Describes cells that have a protein to which the hormone estrogen will bind. Cancer cells that are ER-positive need estrogen to grow, and may stop growing when treated with hormones that block estrogen from binding.

Estrogen receptor: receptor for estrogens; its presence conveys a better prognosis for breast cancers.

Gene expression: the detectable effect of a gene.

HER2 (human epidermal growth factor receptor 2): A protein involved in normal cell growth. It is found in high levels on some breast cancer cells.

Micrometastasis: small numbers of cancer cells that have spread from the primary tumor to other parts of the body and are too few to be picked up in a screening or diagnostic test.

Node-negative: being or having cancer that has not spread to nearby lymph nodes.

PR-positive (progesterone receptor positive): Describes cells that have a protein to which the hormone progesterone will bind. Cancer cells that are PR-positive need progesterone to grow and will usually stop growing when treated with hormones that block progesterone from binding.

Stage T1: The tumor is 20 millimeters (mm) or smaller in size at its widest area.

Stage T2: The tumor is larger than 20 mm but not larger than 50 mm.

Stage T3: The tumor is larger than 50 mm.

RELATED GUIDELINES:

[Genetic Testing, 05-82000-28](#)

[Tumor/Genetic Markers, 05-86000-22](#)

OTHER:

None applicable.

REFERENCES:

1. American Society of Clinical Oncology (ASCO). Breast Cancer: Stages, 07/2019; accessed at cancer.net.
2. Andre F, Ismaila N, et al. Use of Biomarkers to Guide Decisions on Adjuvant Systemic Therapy for Women With Early-Stage Invasive Breast Cancer: ASCO Clinical Practice Guideline Update—Integration of Results From TAILORx. J Clin Oncol. 2019 Aug 1;37(22):1956-1964.
3. Bartlett JMS, Sgroi DC, et al. Breast Cancer Index and prediction of benefit from extended endocrine therapy in breast cancer patients treated in the Adjuvant Tamoxifen-To Offer More? (aTTom) trial. Ann Oncol. 2019 Nov 1;30(11):1776-1783.
4. Bender RA, Knauer M, Rutgers EJ, et al, The 70-Gene Profile and Chemotherapy Benefit in 1,600 Breast Cancer Patients, J Clin Oncol 27: 15s, 2009.
5. Bianchini G, Zambetti M, Pusztai L, et al, Use of Estrogen Receptor (ER) Expression by Quantitative RT-PCR to Identify an ER-Negative Subgroup by IHC Who Might Benefit From Hormonal Therapy, American Society of Clinical Oncology 207 Breast Cancer Symposium, Abstract No. 106.

6. Blue Cross Blue Shield Association Evidence Positioning System®. 2.04.36 Assays of Genetic Expression in Tumor Tissue as a Technique to Determine Prognosis in Patients With Breast Cancer, 12/20.
7. Blue Cross Blue Shield Association Technology Evaluation Center (TEC), Gene expression profiling in women with lymph node negative breast cancer to select adjuvant chemotherapy, TEC Assessments 2014; Volume 29.
8. Blue Cross Blue Shield Association Technology Evaluation Center (TEC), Gene Expression Profiling in Women with Lymph-Node-Positive Breast Cancer to Select Adjuvant Chemotherapy Treatment, 04/10.
9. Blue Cross Blue Shield Association Technology Evaluation Center (TEC), Gene Expression Profiling of Breast Cancer to Select Women for Adjuvant Chemotherapy, 2007 Volume 22, No. 15, 04/08.
10. Blue Cross Blue Shield Association Technology Evaluation Center (TEC), Gene Expression Profiling for Managing Breast Cancer Treatment, 06/07.
11. Bosl A, Spitzmuller A, Jasarevic Z, et al. MammaPrint versus EndoPredict: Poor correlation in disease recurrence risk classification of hormone receptor positive breast cancer. *PLoS One*. 2017;12(8):e0183458.
12. Burstein HJ, Curigliano G, et al. Estimating the benefits of therapy for early-stage breast cancer: the St. Gallen International Consensus Guidelines for the primary therapy of early breast cancer 2019. *Ann Oncol*. Oct 01 2019; 30(10): 1541-1557. PMID: 31373601.
13. Burstein HJ, Lacchetti C, et al. Adjuvant Endocrine Therapy for Women With Hormone Receptor-Positive Breast Cancer: ASCO Clinical Practice Guideline Focused Update. *J Clin Oncol*. Feb 10 2019; 37(5): 423-438. PMID: 30452337.
14. Buus R, Sestak I, Kronenwett R, et al. Comparison of EndoPredict and Epclin with Oncotype DX Recurrence Score for prediction of risk of distant recurrence after endocrine therapy. *J Natl Cancer Inst*. Nov 2016;108(11).
15. Cardoso F, van't Veer LJ, et al, 70-Gene Signature as an Aid to Treatment Decisions in Early-Stage Breast Cancer. *N Engl J Med*. 2016 Aug 25;375(8):717-29.
16. ClinicalTrials.gov. MINDACT (Microarray In Node-Negative and 1 to 3 Positive Lymph Node Disease May Avoid Chemotherapy): A Prospective, Randomized Study Comparing the 70-Gene Signature With the Common Clinical-Pathological Criteria in Selecting Patients for Adjuvant Chemotherapy in Breast Cancer With 0 to 3 Positive Nodes; accessed December 2020.
17. ClinicalTrials.gov. The PRECISION Trial (Profiling Early Breast Cancer for Radiotherapy Omission): a Phase II Study of Breast-Conserving Surgery Without Adjuvant Radiotherapy for Favorable Risk Breast Cancer; accessed December 2020.
18. ClinicalTrials.gov. Program for the Assessment of Clinical Cancer Tests (PACCT-1): Trial Assigning Individualized Options for Treatment: The TAILORx Trial; accessed December 2020.
19. ClinicalTrials.gov. A Randomised Phase III Trial of Adjuvant Radiation Therapy Versus Observation Following Breast Conserving Surgery and Endocrine Therapy in Patients With Molecularly Characterised Luminal A Early Breast Cancer; accessed December 2020.
20. Cronin M, Sangli C, Liu ML, et al, Analytical Validation of the Oncotype DX Genomic Diagnostic Test for Recurrence Prognosis and Therapeutic Response Prediction in Node-negative, Estrogen Receptor-positive Breast Cancer, *Clin Chem*. 06/07.
21. Cusumano PG, Generali D, et al, European inter-institutional impact study of MammaPrint, *Breast*. 2014 Aug;23(4):423-8.
22. De Boer M, Van Deurzen CHM, et al, Micrometastases or Isolated Tumor Cells and the Outcome of Breast Cancer, *The New England Journal of Medicine*, Vol. 361, No. 7, August 13, 2009.

23. Dowsett M, Sestak I, et al, Comparison of PAM50 Risk of Recurrence Score With Oncotype DX and IHC4 for Predicting Risk of Distant Recurrence After Endocrine Therapy, *J Clin Oncol*. 2013 Aug 1;31(22):2783-90.
24. Drukker CA, Bueno-de-Mesquita JM, et al, A prospective evaluation of a breast cancer prognosis signature in the observational RASTER study, *Int J Cancer*. 2013 Aug 15;133(4):929-36.
25. Drukker CA, Schmidt MK, et al, Mammographic screening detects low-risk tumor biology breast cancers, *Breast Cancer Res Treat* 2014, 144:103–111.
26. Drukker CA, van den Hout HC, et al, Risk estimations and treatment decisions in early stage breast cancer: agreement among oncologists and the impact of the 70-gene signature, *Eur J Cancer*. 2014 Apr;50(6):1045-54.
27. Drukker CA, van Tinteren H, et al, Long-term impact of the 70-gene signature on breast cancer outcome, *Breast Cancer Res Treat* (2014) 143:587–592.
28. Ettl J, Anders SI, et al. First prospective outcome data for the second-generation multigene test Endopredict in ER-positive/HER2-negative breast cancer. *Arch Gynecol Obstet*. Dec 2020; 302(6): 1461-1467. PMID: 32902674.
29. Evaluation of Genomic Applications in Practice and Prevention (EGAPP) Working Group: Can Tumor Gene Expression Profiling Improve Outcomes in Patients with Breast Cancer? *Genet Med* 2009 Jan; 11(1): 66-73.
30. Fan, C., Oh, D. S., Wessels, L., Weigelt, B., Nuyten, D. S. A., Nobel, A., B., van't Veer, L. J. & Perou, C. M. (2006). Concordance Among Gene-Expression-Based Predictors for Breast Cancer. *N Engl J Med*, 355:560-569.
31. Filipits M, Dubsy P, et al. Prediction of Distant Recurrence Using EndoPredict Among Women with ER + , HER2 – Node-Positive and Node-Negative Breast Cancer Treated with Endocrine Therapy Only. *Clin Cancer Res*. Jul 01 2019; 25(13): 3865-3872. PMID: 31064782.
32. Funakoshi Y, Wang Y, et al. Comparison of molecular profile in triple-negative inflammatory and non-inflammatory breast cancer not of mesenchymal stem-like subtype. *PloS One*. 2019 Sep 18;14(9):e0222336. PMID: 31532791.
33. Geradts, J. Molecular Prediction of Recurrence of Breast Cancer. *N Engl J Med*, 352:1605. (ltr to editor) 04/14/05.
34. Glück S, de Snoo F, et al, Molecular subtyping of early-stage breast cancer identifies a group of patients who do not benefit from neoadjuvant chemotherapy, *Breast Cancer Res Treat*. 2013 Jun;139(3):759-67.
35. Gold JM, Najita JS, Lester, S, et al, Personalizing Treatment in Early-Stage Breast Cancer, *J Clin Oncol* 27: 15s, 2009.
36. Goldhirsch A, Winer EP, et al. Personalizing the treatment of women with early breast cancer: highlights of the St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2013. *Ann Oncol* 2013; 24(9):2206-23.
37. Goldhirsch A, Wood WC, Gelber RD, et al, Progress and Promise: Highlights of the International Expert Consensus on the Primary Therapy of Early Breast Cancer 2007, *Ann Oncol*. 2007 Jul; 18(7): 1133-44.
38. Goldstein LJ, Gray R, Badve S, et al, Prognostic Utility of the 21-Gene Assay in Hormone Receptor-Positive Operable Breast Cancer Compared with Classical Clinicopathologic Features, *Journal of Clinical Oncology*, Vol 26, No 25, 09/08.
39. Goodson, W. H. Molecular Prediction of Recurrence of Breast Cancer. *N Engl J Med*, 352:1606. (ltr to editor) 04/14/05.
40. Grant KA, Apffelstaedt JP, et al, MammaPrint Pre-screen Algorithm (MPA) reduces chemotherapy in patients with early-stage breast cancer, *S Afr Med J*. 2013 Jul 3;103(8):522-6.

41. Groenedijk FH, Zwart W, et al, Estrogen receptor splice variants as a potential source of false-positive estrogen receptor status in breast cancer diagnostics, *Breast Cancer Res Treat.* 2013 Aug;140(3):475-84.
42. Habel, L. A., Shak, S., Jacobs, M. K., Capra, A., Alexander, C., Pho, M., Baker, J., Walker, M., Watson, D., Hackett, J., Blick, N. T., Greenberg, D., Fehrenbacher, L., Langholz, B. & Quesenberry, C. P. (2006). A Population-Based Study of Tumor Gene Expression and Risk of Breast Cancer Death Among Lymph Node-Negative Patients. *Breast Cancer Research* (05/06).
43. Harbeck N, Nimmrich I, Hartmann A, et al, Multicenter Study Using Paraffin-Embedded Tumor Tissue Testing PITX2 DNA Methylation As a Marker for Outcome Prediction in Tamoxifen-Treated, Node-Negative Breast Cancer Patients, *Journal of Clinical Oncology*, 08/08.
44. Harris LN, Ismaila N, McShane LM, et al. Use of biomarkers to guide decisions on adjuvant systemic therapy for women with early-stage invasive breast cancer: American Society of Clinical Oncology Clinical Practice Guideline. *J Clin Oncol.* Apr 1 2016;34(10):1134-1150.
45. Henry NL, Somerfield, et al. Role of Patient and Disease Factors in Adjuvant Systemic Therapy Decision Making for Early-Stage, Operable Breast Cancer: Update of the ASCO Endorsement of the Cancer Care Ontario Guideline. *J Clin Oncol.* 2019 Aug 1;37(22):1965-1977.
46. Jansen MP, Sieuwerts AM, Look MP, et al, HOXB13-to-IL17BR Expression Ratio is Related with Tumor Aggressiveness and Response to Tamoxifen of Recurrent Breast Cancer: A Retrospective Study, *Journal of Clinical Oncology*, Vol 25, No 6, pp. 662-668, 02/07.
47. Jerevall PL, Brommesson S, Strand C, et al, Exploring the Two-Gene Ratio in Breast Cancer-Independent Roles For HOXB13 and IL17BR in Prediction of Clinical Outcome, *Breast Cancer Research Treatment*, 2008 Jan; 107(2): 225-34.
48. Kaklamani V, A Genetic Signature Can Predict Prognosis and Response to Therapy in Breast Cancer: *Oncotype DX*, *Expert Rev Mol Diagn.* 11/06.
49. Kittaneh M, Montero AJ, et al, Molecular profiling for breast cancer: a comprehensive review, *Biomark Cancer.* 2013 Oct 29;5:61-70.
50. Krop I, Ismaila N, et al. Use of Biomarkers to Guide Decisions on Adjuvant Systemic Therapy for Women With Early-Stage Invasive Breast Cancer: American Society of Clinical Oncology Clinical Practice Guideline Focused Update. *J Clin Oncol.* 2017 Aug 20;35(24):2838-2847.
51. Kunz G, Use of a Genomic Test (MammaPrint) in Daily Clinical Practice to Assist in Risk Stratification of Young Breast Cancer Patients, *Arch Gynecol Obstet.* 2010 Apr 13.
52. Lyman GH, Cosler LE, Kuderer NM, Hornberger J, Impact of a 21-Gene PR_PCR Assay on Treatment Decisions in Early-Stage Breast Cancer: An Economic Analysis Based on Prognostic and Predictive Validation Studies, *Cancer.* 2007 Mar 15; 109(6): 1011-8.
53. Lyman GH, Kuderer NM, Gene Expression Profile Assays as Predictors of Recurrence-Free Survival in Early-Stage Breast Cancer: A Metaanalysis, *Clin Breast Cancer.* 2006 Dec; 7(5): 372-9.
54. Ma SJ, Salunga R, Dahiya S, et al, A Five-Gene Molecular Grade Index and HOXB13:IL17BR Are Complementary Prognostic Factors in Early Stage Breast Cancer, *Clinical Cancer Research*, 14, 2601-2608, 05/08.
55. Marchionni L, Wilson RF, Wolff AC, et al, Systematic Review: Gene Expression Profiling Assays in Early-Stage Breast Cancer, *Annals of Internal Medicine*, Vol 148 Issue 5, pp. 358-369, 03/08.
56. Martin M, Brase JC, et al. Clinical validation of the EndoPredict test in node-positive chemotherapy-treated ER+/HER2- breast cancer patients: results from the GEICAM/9906 trial. *Breast Cancer Res* 2014; 16(2):R38.
57. National Cancer Institute, *Cancer Diagnostics: Informing the Development of Tailored Cancer Therapy*, 05/23/06.

58. National Comprehensive Cancer Network[®] (NCCN). NCCN Clinical Practice Guidelines in Oncology: Breast Cancer; located at nccn.org.
59. Nielsen T, Wallden B, et al. Analytical validation of the PAM50-based Prosigna Breast Cancer Prognostic Gene Signature Assay and nCounter Analysis System using formalin-fixed paraffin-embedded breast tumor specimens. *BMC Cancer* 2014; 14:177.
60. Noordhoek I, Treuner K, et al. Breast Cancer Index Predicts Extended Endocrine Benefit to Individualize Selection of Patients with HR + Early-stage Breast Cancer for 10 Years of Endocrine Therapy. *Clin Cancer Res.* 2021 Jan 1;27(1):311-319.
61. O'Connor SM, Beriwal S, Dabbs DJ, Bhargava R, Concordance Between Semiquantitative Immunohistochemical Assay and Oncotype DX RT-PCR Assay for Estrogen and Progesterone Receptors, *Applied Immunohistochemistry & Molecular Morphology*, May 2010, Vol. 18, Issue 3, pp 268-272.
62. Paik S, Tang G, Fumagalli D, Editorial: An Ideal Prognostic Test for Estrogen Receptor- Positive Breast Cancer? *Journal of Clinical Oncology*, Vol 26, No25, 09/01/08.
63. Paik, S., Shak, S., Tang, G., Kim, C., Baker, J., Cronin, M., Baehner, F. L., Walker, M. G., Watson, D., Park, T., Hiller, W., Fisher, E. R., Wickerham, D. L., Bryant, J. and Wolmark, N. (2004). A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. *N Engl J Med*, 351:2817-26.
64. Paik, S., Tang, G., Shak, S., Kim, C., Baker, J., Kim, W., Cronin, M., Baehner, F. L., Watson, D., Bryant, J., Costantino, J. P., Geyer, C. E., Wickerham, D. L. & Wolmark, N. (2006). Gene Expression and Benefit of Chemotherapy in Women With Node-Negative, Estrogen Receptor-Positive Breast Cancer. *Journal of Clinical Oncology*, 24(23).
65. Paik, S., Wolmark, N. and Shak, S. Molecular Prediction of Recurrence of Breast Cancer. *N Engl J Med*, 352:1606 (authors' reply) 04/14/05.
66. Palmetto GBA, LLC. MoIDX Local Coverage Determinations (LCDs) located at palmettogba.com.
67. Puztai L, Current Status of Prognostic Profiling in Breast Cancer, *The Oncologist* 2008; 13:350-360, 02/08.
68. Retèl VP, Groothuis-Oudshoorn CG, et al, Association between genomic recurrence risk and well-being among breast cancer patients, *BMC Cancer*. 2013 Jun 18;13:295.
69. Retèl VP, Joore MA, et al, Prospective cost-effectiveness analysis of genomic profiling in breast cancer, *Eur J Cancer*. 2013 Dec;49(18):3773-9.
70. Ross JS, Hatzis C, Symmans WF, et al, Commercialized Multigene Predictors of Clinical Outcome for Breast Cancer, *The Oncologist* 2008; 13:477-493, 03/08
71. Sapino A, Roepman P, et al, MammaPrint molecular diagnostics on formalin-fixed, paraffin-embedded tissue, *J Mol Diagn.* 2014 Mar;16(2):190-7.
72. Schroeder B, Zhang Y, Stal O, et al. Risk stratification with Breast Cancer Index for late distant recurrence in patients with clinically low-risk (T1N0) estrogen receptor-positive breast cancer. *NPJ Breast Cancer*. 2017;3:28.
73. Sestak I, Buus R, et al. Comparison of the Performance of 6 Prognostic Signatures for Estrogen Receptor-Positive Breast Cancer: A Secondary Analysis of a Randomized Clinical Trial. *JAMA Oncol.* 2018 Apr 1;4(4):545-553.
74. Sestak I, Martin M, et al. Prediction of chemotherapy benefit by EndoPredict in patients with breast cancer who received adjuvant endocrine therapy plus chemotherapy or endocrine therapy alone. *Breast Cancer Res Treat.* Jul 2019; 176(2): 377-386. PMID: 31041683.
75. Shak S, Palmer G, Baehner FL, et al, Molecular Characterization of Male Breast Cancer by Standardized Quantitative RT-PCR Analysis, *J Clin Oncol* 27: 15s, 2009.

76. Simon RM, Paik S, Hayes DF. Use of Archived Specimens in Evaluation of Prognostic and Predictive Biomarkers. *J Natl Cancer Inst.* 2009 Nov 4; 101(21): 1446–1452.
77. Soliman H, Flake DD, et al. Predicting Expected Absolute Chemotherapy Treatment Benefit in Women With Early-Stage Breast Cancer Using EndoPredict, an Integrated 12-Gene Clinicomolecular Assay. *JCO Precision Oncology* 2019 :3, 1-10.
78. Solin LJ, Gray R, et al, A Multigene Expression Assay to Predict Local Recurrence Risk for Ductal Carcinoma In Situ of the Breast, *J Natl Cancer Inst*;2013;105:701–710.
79. Sparano JA, Gray RJ, et al. Adjuvant Chemotherapy Guided by a 21-Gene Expression Assay in Breast Cancer. *N Engl J Med.* 2018 Jul 12;379(2):111-121.
80. Stephenson, J. (2002). Gene studies provide window on cancer prognosis, treatment benefits, toxic effects. *JAMA*, 288:820-821.
81. Swain, S. M. (2006). A Step in the Right Direction. *Journal of Clinical Oncology*, 24(23).
82. Tang, C. G. and Lin, A. Y. Molecular Prediction of Recurrence of Breast Cancer. *N Engl J Med*, 352:1606. (Ltr to editor) 04/14/05.
83. Tanvetyanon, T. Molecular Prediction of Recurrence of Breast Cancer. *N Engl J Med*, 352:1606. (Ltr to editor) 04/14/05.
84. Tsoi DT, Inoue M, Kelly CM, et al, Coast-Effectiveness Analysis of Oncotype DX-Guided Treatment in Early Breast Cancer, *J Clin Oncol* 27, 2009.
85. Ueda S, Saeki T, et al, Genomic Profiling Shows Increased Glucose Metabolism in Luminal B Breast Cancer, *J Breast Cancer*. 2013 Sep;16(3):342-4.
86. van 't Veer LJ, Yau C, Yu NY, et al. Tamoxifen therapy benefit for patients with 70-gene signature high and low risk. *Breast Cancer Res Treat.* Aug 04 2017.
87. Van Ze KJ, Zabor EC, et al. Comparison of Local Recurrence Risk Estimates After Breast-Conserving Surgery for DCIS: DCIS Nomogram Versus Refined Oncotype DX Breast DCIS Score. *Ann Surg Oncol.* 2019 Jul 24. Doi:10.1245/s10434-019-07537-y.Epub ahead of print] PMID: 31342373.
88. Vieira AF, Schmitt F. An Update on Breast Cancer Multigene Prognostic Tests-Emergent Clinical Biomarkers. *Front Med (Lausanne)*. 2018; 5: 248.
89. Wittner BS, Sgroi DC, Ryan PD, et al, Analysis of the MammaPrint Breast Cancer Assay in a Predominantly Postmenopausal Cohort, *Clinical Cancer Research* 14, 2988-2993, 05/08.

COMMITTEE APPROVAL:

This Medical Coverage Guideline (MCG) was approved by the Florida Blue Medical Policy & Coverage Committee on 04/22/21.

GUIDELINE UPDATE INFORMATION:

10/15/05	New Medical Coverage Guideline.
01/01/06	Annual HCPCS coding update: added code S3854.
09/15/06	Annual review: continue investigational status.
11/15/07	Annual review: position statements changed, description section updated, definitions section updated, guideline reformatted, references updated.
02/15/08	Revised position statement; updated description section, definition section and references.
01/01/09	Annual review: no change to position statements, description section and references updated.

11/15/09	Annual review: position statements updated; description section and references updated.
09/15/10	Annual review: position statements maintained and references updated.
11/15/10	Revision; formatting changes and Program Exceptions section updated.
08/15/11	Scheduled review; position statements maintained and references updated; formatting changes.
03/15/12	Revision; position statement updated; formatting changes.
10/15/12	Annual review; position statements maintained and references updated.
11/01/12	Remove CMN; formatting changes.
05/30/13	Revision; Program Exceptions section and references updated.
11/15/13	Annual review; position statements, program exception, and references updated; formatting changes.
07/01/14	Quarterly HCPCS update. Added code 0008M.
10/15/14	Annual review; position statement section, program exception section, and references updated; formatting changes.
01/01/15	Annual HCPCS/CPT update. Added code 81519.
11/15/15	Annual review; position statement, program exception, and references updated; formatting changes.
01/01/16	Annual HCPCS/CPT update; code S3854 deleted.
07/01/16	Quarterly HCPCS/CPT update. Added code S3854 (code reinstated).
02/15/17	Revision; description, position statements, and references updated; formatting changes.
01/01/18	Annual CPT/HCPCS update. Added codes 81520, 81521; deleted code 0008M.
03/15/18	Review; position statements maintained; description section, coding, and references updated.
07/01/18	Quarterly CPT/HCPCS update. Added code 0045U.
12/15/18	Revision; MammaPrint added to list of tests meeting the definition of medical necessity for specific indications; investigational for all others; coding, program exception, and references updated.
01/01/19	Annual CPT/HCPCS coding update. Added code 81518.
02/15/19	Revision; Coding section, program exception section, and references updated.
11/15/19	Review; Position statements maintained and references updated.
01/01/20	Annual CPT/HCPCS coding update. Added codes 81522 & 0153U.
02/15/20	Revision; MammaPrint position statement added; references updated.
10/15/20	Revision; 21-gene reverse transcriptase-polymerase chain reaction (PT-PCR) assay position statement updated.
02/15/21	Review; Position statements and references updated.
05/15/21	Revision; Position statement for the use of BCI assay to assist in decision of extending adjuvant hormonal therapy added; references updated.
01/01/22	Annual CPT/HCPCS coding update. Code 81523 added.