

02-61000-28

Original Effective Date: 11/15/00

Reviewed: 09/25/25

Revised: 10/15/25

Subject: Lysis of Epidural Adhesions

THIS MEDICAL COVERAGE GUIDELINE IS NOT AN AUTHORIZATION, CERTIFICATION, EXPLANATION OF BENEFITS, OR A GUARANTEE OF PAYMENT, NOR DOES IT SUBSTITUTE FOR OR CONSTITUTE MEDICAL ADVICE. ALL MEDICAL DECISIONS ARE SOLELY THE RESPONSIBILITY OF THE PATIENT AND PHYSICIAN. BENEFITS ARE DETERMINED BY THE GROUP CONTRACT, MEMBER BENEFIT BOOKLET, AND/OR INDIVIDUAL SUBSCRIBER CERTIFICATE IN EFFECT AT THE TIME SERVICES WERE RENDERED. THIS MEDICAL COVERAGE GUIDELINE APPLIES TO ALL LINES OF BUSINESS UNLESS OTHERWISE NOTED IN THE PROGRAM EXCEPTIONS SECTION.

[Position Statement](#)

[Billing/Coding](#)

[Reimbursement](#)

[Program Exceptions](#)

[Definitions](#)

[Related Guidelines](#)

[Other](#)

[References](#)

[Updates](#)

DESCRIPTION:

Lysis of epidural adhesions, also called the Racz procedure, has been investigated as a treatment option for epidural fibrosis and other conditions. The Racz procedure involves the passage of a fluoroscopically guided catheter (the Racz catheter), inserted either endoscopically or percutaneously, and the use of epidural injections of hypertonic saline in conjunction with corticosteroids and analgesics. Theoretically, the use of hypertonic saline results in a mechanical disruption of the adhesions. The saline may also function to reduce edema within previously scarred and/or inflamed nerves. Finally, manipulating the catheter at the time of the injection may disrupt adhesions. Spinal endoscopy has been used to guide the lysis procedure, but the procedure is more commonly performed percutaneously using epidurography to guide catheter placement and identify non-filling adhesions that indicate epidural scarring. In some situations, the catheter may remain in place for several days for serial treatment sessions.

Summary and Analysis of Evidence: Kose, Akkaya (2023) set out to assess the effectiveness of percutaneous adhesiolysis (PEA) in patients with chronic lumbar radicular pain refractory to epidural steroid injections, and to determine predictive factors, including demographic, clinical, and procedural data, to provide superior treatment efficacy. One hundred and ninety-three patients were reviewed. Successful treatment outcome was described as a 50% reduction in the visual analog scale score. Among the 193 patients, 109 (56.2%) exhibited a positive treatment response at 12 months. In multivariate logistic regression analysis, no depression, no spondylolisthesis, no previous lumbar surgery, mild foraminal stenosis, no opioid use, and baseline pain scores were the predictive factors significantly associated with unsuccessful treatment outcome. The authors concluded that PEA is a useful treatment option for patients with chronic lumbar radicular pain refractory to epidural steroid injections, and that a history of lumbar surgery, spondylolisthesis, depression, and severe foraminal stenosis could be associated with a poor prognosis. Limitations of this study included that the regression analyses used

were focused on assessing the influence of predictive factors on outcomes at the 12th month; the ability to predict successful outcomes over the longer period remains uncertain. In addition, the success of the treatment was confined solely to changes in pain relief. Due to the lack of routine recording in the clinic, they were unable to track changes in pain medication consumption or functional disability. Also, as with many retrospective studies, there were patients who had to be excluded due to missing data. Finally, they noted that this study was conducted retrospectively, lacking a control or sham group for a comparative analysis of the procedure's outcomes. The prevalence of placebo and nocebo effects in the context of interventional treatments was estimated to range between 13% and 30% and 3% and 8%, respectively.

Gerdesmeyer et al (2021) published 10 year follow-up results from a previous randomized controlled trial (RCT). After a 4 year enrollment phase, 381 patients with chronic radicular pain persisting beyond 4 months, who failed conservative treatments, were screened. Ninety patients were enrolled. Patients were randomly assigned to receive percutaneous epidural lysis of adhesions or placebo with concealed allocation in permuted blocks of 4 to 8 patients each, and stratified by treatment center. The primary outcomes were a mean change of the Oswestry Disability Index (ODI) scores and Visual Analog Scale (VAS), one and 10 years after intervention. For each rating scale an analysis of variance with the within-patient factor time (baseline, one year follow-up, 10 year follow-up) and the between-patient factor treatment (lysis, placebo) was used. The ODI and VAS scores were significantly better one and 10 years in the lysis group vs the control group. The ODI in the lysis group improved from 55.3 ± 11.6 to 9.6 ± 9.3 after one year and to 11.7 ± 14.2 after 10 years. The placebo group also improved from 55.4 ± 11.5 to 30.7 ± 14.2 after one year and to 24.8 ± 12.0 after 10 years. The VAS improved from 6.7 ± 1.1 to 1.2 ± 1.1 after one year and to 1.5 ± 1.4 after 10 years in the lysis group and from 6.7 ± 1.1 to 2.8 ± 1.5 after one year and to 2.9 ± 1.3 after 10 years after placebo intervention. No treatment-related severe adverse effects occurred within the 10 years, but minor transient neurological effects were seen directly after the intervention. The authors concluded that the 10 year follow-up demonstrated efficacy of the minimally invasive percutaneous adhesiolysis procedure for patients with chronic lumbosacral radicular pain, and that this procedure should be considered as the first treatment option for patients with chronic lumbosacral radicular pain. Several limitations were acknowledged by the authors, including 1) that the long-term effects of single treatment components cannot be specified as no imaging examination was performed at 10 year follow-up; 2) a large variety of unanalyzed noninvasive treatments were done within the 10 years; 3) some participants did not clearly remember the intervention after 10 years; and 4) uncontrolled effects such as higher inhomogeneity of biometric properties, concomitant therapies, pain tolerance level, or just social effects could occur, but were not analyzed in the trial.

Cho et al (2019) studied whether the outcomes of percutaneous epidural neuroplasty (PEN) are influenced by the type of lumbar disc herniation (LDH), and set out to evaluate the effectiveness of PEN in patients with single-level LDH. This study included 430 consecutive patients with single-level LDH who underwent PEN. Before treatment, the LDH type was categorized as bulging, protrusion, extrusion, and sequestration, while Pfirrmann grades were assigned according to imaging findings. Visual analog scale (VAS) scores for back and leg pain and success rates (Odom's criteria) were assessed at 1, 3, 6, and 12 months after treatment. The mean preoperative VAS scores for back and leg pain were 6.90 and 4.23, respectively; these decreased after PEN as follows: 2.25 and 1.45, respectively, at 1 month; 2.61 and 1.68, respectively, at 3 months; 2.28 and 1.48, respectively, at 6 months; and 2.88 and 1.48,

respectively, at 12 months. The decrease in VAS scores for leg pain was significantly greater in the extrusion and sequestration groups than in the other two groups; there were no other significant differences among groups. More than 70% patients exhibited good or excellent 12-month outcomes according to Odom's criteria. Subsequent surgery was required for 59 patients (13.7%), with a significantly higher rate in the extrusion (25.0%) and sequestration (30.0%) groups than in the bulging (7.3%) and protrusion (13.8%) groups. Nevertheless, subsequent surgery was not required for >70% patients with extrusion or sequestration. Patients with Pfirrmann grades 1-3 (14.0-21.5%) showed a significantly higher rate of subsequent surgery than those with Pfirrmann grade 0 (4.9%). The authors concluded their findings suggest that PEN is an effective treatment for back and leg pain caused by single-level LDH, with the outcomes remaining unaffected by the LDH type. They noted several study limitations, including 1) it was a retrospective analysis; 2) a control group was not included for comparison, so improved clinical outcomes after PEN could not be distinguished from the natural course of LDH; 3) the distribution of patients according to the level and type of LDH was uneven, and this may have affected the statistical results, and 4) this study did not take the MRI after procedure for checking the spontaneous regression of disc (this regression can affect the PEN effect according the disc type).

Manchikanti et al (2012) studied 120 participants who were randomly assigned to two groups with a 2-year follow-up. Group I (control group, n = 60) received caudal epidural injections with catheterization up to S3 with local anesthetic (lidocaine 2%, 5 mL), nonparticulate betamethasone (6 mg, 1 mL), and 6 mL of 0.9% sodium chloride solution. Group II (intervention group, n = 60) received percutaneous adhesiolysis of the targeted area, with targeted delivery of lidocaine 2% (5 mL), 10% hypertonic sodium chloride solution (6 mL), and nonparticulate betamethasone (6 mg). The multiple outcome measures included the Numeric Rating Scale, the Oswestry Disability Index 2.0, employment status, and opioid intake with assessments at 3, 6, 12, 18, and 24 months posttreatment. Primary outcome was defined as 50% improvement in pain and Oswestry Disability Index scores. Significant improvement with at least 50% relief with pain and improvement in functional status was illustrated in 82% of patients at the 2-year follow-up in the intervention group compared to 5% in the control group receiving caudal epidural injections. The average number of procedures over a period of 2 years in Group II was 6.4 ± 2.35 with overall total relief of approximately 78 weeks out of 104 weeks. The authors concluded that this study demonstrated significant improvement in 82% of patients over a period of 2 years with an average of six to seven procedures of 1-day percutaneous adhesiolysis in patients with failed back surgery syndrome. The authors acknowledged several study limitations, including that given the subjective outcome of pain relief, an equivalence study with no placebo/sham control is difficult to interpret; and, there was a large control group dropout rate (n=43 in control group; n=3 in intervention group) at 12 months.

UpToDate review "Subacute and chronic low back pain: Surgical treatment" (Chou, 2024) states that therapies for subacute and chronic low back pain have included adhesiolysis (injection of isotonic saline, hypertonic saline, or hyaluronidase into the epidural space in order to facilitate lysis of adhesions); however, "randomized trials for these interventions are not available or inconclusive".

POSITION STATEMENT:

Catheter-based techniques for lysis of epidural adhesions, with or without endoscopic guidance, used alone or in combination with injection of hypertonic solutions (e.g., saline, corticosteroids, analgesics, hyaluronidase) are considered **experimental or investigational**. Data in published medical literature are inadequate to permit scientific conclusions on long-term and net health outcomes.

BILLING/CODING INFORMATION:

CPT Coding:

62263	Percutaneous lysis of epidural adhesions using solution injection (e.g., hypertonic saline, enzyme) or mechanical means (e.g., catheter) including radiologic localization (includes contrast when administered), multiple adhesiolysis sessions; 2 or more days (investigational)
62264	Percutaneous lysis of epidural adhesions using solution injection (e.g., hypertonic saline, enzyme) or mechanical means (e.g., catheter) including radiologic localization (includes contrast when administered), multiple adhesiolysis sessions; 1 day (investigational)
62280	Injection/infusion of neurolytic substance (eg, alcohol, phenol, iced saline solutions), with or without other therapeutic substance; subarachnoid (investigational)
62281	Injection/infusion of neurolytic substance (eg, alcohol, phenol, iced saline solutions), with or without other therapeutic substance; epidural, cervical or thoracic (investigational)
62282	Injection/infusion of neurolytic substance (eg, alcohol, phenol, iced saline solutions), with or without other therapeutic substance; epidural, lumbar, sacral (caudal) (investigational)

REIMBURSEMENT INFORMATION:

Refer to sections entitled **POSITION STATEMENT** and **PROGRAM EXCEPTIONS**.

PROGRAM EXCEPTIONS:

Federal Employee Program (FEP): Follow FEP guidelines.

State Account Organization (SAO): Follow SAO guidelines.

Medicare Advantage products: No National Coverage Determination (NCD) and/or Local Coverage Determination (LCD) were found at the time of the last guideline review date.

If this Medical Coverage Guideline contains a step therapy requirement, in compliance with Florida law 627.42393, members or providers may request a step therapy protocol exemption to this requirement if based on medical necessity. The process for requesting a protocol exemption can be found at Coverage Protocol Exemption Request.

DEFINITIONS:

Adhesion: a fibrous band or structure by which parts abnormally adhere.

Arachnoiditis: Inflammation of the middle layer of membranes covering the brain and spinal cord. Arachnoiditis can occur as a complication of procedures such as myelograms, spinal operations, epidural steroid injections, and injury to the spine.

Epidural: outside the dura mater (outermost, toughest, most fibrous of the three membranes covering the brain and spinal cord); the space between the bony vertebral canal and the dura mater (the spinal cord does not fill the bony vertebral canal). The space remaining between the dura mater and the bone of the vertebra is the epidural space.

Fluoroscopy: a radiographic technique in which an object, such as the human body, is examined visually in real time by transmitting X-rays through the object onto a fluorescent screen. The resulting picture on the screen is made up of shadows created by the transmission of different amounts of X-rays through body structures of varying depth and size.

Hypertonic: a solution with a higher salt concentration than in normal cells of the body and the blood.

Lysis: destruction; rupture of cell membrane and loss of cytoplasm.

Neurolysis: destruction of nerve tissue, freeing of a nerve from inflammatory adhesions.

Percutaneous: performed through the skin.

Radiculopathy: any disease of the spinal nerve roots and spinal nerves.

RELATED GUIDELINES:

None applicable.

OTHER:

Other names used to report Percutaneous Lysis of Epidural Adhesions:

Epidural Adhesiolysis

Epidural Neurolysis

Epidurolysis

Hypertonic Saline Injections

Injections, Epidural, Hypertonic Saline

Lysis of Epidural Adhesions

Neurolysis, Epidural

Percutaneous Epidural Neuroplasty

Racz Procedure

REFERENCES:

1. AHRQ National Guideline Clearinghouse. Guideline Summary NGC-9330. Chronic pain disorder medical treatment guidelines. Denver (CO): Colorado Division of Workers' Compensation; 2011 Dec 27.
2. AHRQ National Guideline Clearinghouse. Comprehensive evidence-based guidelines for interventional techniques in the management of chronic spinal pain. Pain Physician 2009 Jul-Aug;12(4):699-802.
3. American Board of Interventional Pain Management website, Racz procedure.
4. Belozer M, Wang G. Epidural Adhesiolysis for the Treatment of Back Pain. Health Technology Assessment. Olympia, WA: Washington State Department of Labor and Industries, Office of the Medical Director; July 13, 2004.
5. Birkenmaier C, Baumert S, Schroeder C, Jansson V, Wegener B. A Biomechanical Evaluation of the Epidural Neurolysis Procedure. Pain Physician 2012; 15:E89-E97.
6. Blue Cross Blue Shield Association Evidence Positioning System®. 8.01.18 - Lysis of Epidural Adhesions, (ARCHIVED 12/20).

7. Boswell MV, Trescot AM, Datta S, Schultz DM, Hansen HC, Abdi S, Sehgal N, Shah RV, Singh V, Benyamin RM, Patel VB, Buenaventura RM, Colson JD, Cordner HJ, Epter RS, Jasper JF, Dunbar EE, Atluri SL, Bowman RC, Deer TR, Swicegood JR, Staats PS, Smith HS, Burton AW, Kloth DS, Giordano J, Manchikanti L; American Society of Interventional Pain Physicians. Interventional techniques: evidence-based practice guidelines in the management of chronic spinal pain. *Pain Physician*. 2007 Jan; 10(1):7-111.
8. Boswell MV, Shah RV, Everett CR, Sehgal N, Brown AM, Abdi S, Bowman RC 2nd, Deer TR, Datta S, Colson JD, Spillane WF, Smith HS, Lucas LF, Burton AW, Chopra P, Staats PS, Wasserman RA, Manchikanti L. Interventional techniques in the management of chronic spinal pain: evidence-based practice guidelines. *Pain Physician*. 2005 Jan;8(1): 1-47.
9. Ceylan A, et al. Evaluation of the efficacy of epiduroscopic adhesiolysis in failed back surgery syndrome. *Turk J Med Sci*. 2019 Feb 11;49(1):249-257. doi: 10.3906/sag-1807-173. PMID: 30761878.
10. Cho PG, Ji GY, Yoon YS, Shin DA. Clinical Effectiveness of Percutaneous Epidural Neuroplasty According to the Type of Single-Level Lumbar Disc Herniation : A 12-Month Follow-Up Study. *J Korean Neurosurg Soc*. 2019 Nov;62(6):681-690. doi: 10.3340/jkns.2019.0070. Epub 2019 Oct 8.
11. Chopra P, Smith HS, Deer TR, Bowman RC. Role of Adhesiolysis in the Management of Chronic Spinal Pain: A Systematic Review of Effectiveness and Complications. *Pain Physician*. 2005 Jan;8(1): 87-100.
12. Chou R, Qaseem A, Snow V, Casey D, Cross Jr. JT, Shekelle P, Owens DK. Diagnosis and Treatment of Low Back Pain: A Joint Clinical Practice Guideline from the American College of Physicians and the American Pain Society. *Annals of Internal Medicine*. 2007 Oct;147(7):478-491.
13. Chou R, et al. Interventional Therapies, Surgery, and Interdisciplinary Rehabilitation for Low Back Pain An Evidence-Based Clinical Practice Guideline From the American Pain Society. *Spine Volume* 34, Number 10, 2009.
14. ClinicalTrials.gov, Effectiveness of Percutaneous Lumbar Epidural Adhesiolysis and Neurolysis on Low Back Pain, sponsored by Pain Management Center of Paducah. NCT00370994.
15. ClinicalTrials.gov, A Randomized, Equivalence Trial of Percutaneous Lumbar Adhesiolysis and Caudal Epidural Steroid Injections, sponsored by Pain Management Center of Paducah. NCT01053273.
16. ClinicalTrials.gov. Role of Steroids and 10% Hypertonic Sodium Chloride in Adhesiolysis in Post Lumbar Surgery Syndrome Patients, sponsored by Pain Management Center of Paducah. NCT01053572.
17. ECRI Hotline Response: Epidurolysis Procedure Using Racz Catheter for Low Back Pain, (06/26/03).
18. ECRI. Custom Hotline Response. Epidurolysis for Low Back Pain. PA: ECRI. 10/09/07.
19. Epter, RS, Helm S, Hayek SM, Benyamin RM, Smith HS., and Abdi S. Systematic Review of Percutaneous Adhesiolysis and Management of Chronic Low Back Pain in Post Lumbar Surgery Syndrome. *Pain Physician*. 2009; 12; 361-378.
20. Gerdesmeyer L, et al. Percutaneous Epidural Lysis of Adhesions in Chronic Lumbar Radicular Pain: A Randomized, Double-Blind, Placebo-Controlled Trial. *Pain Physician* 2013; 16:185-196.
21. Gerdesmeyer L, Noe C, Prehn-Kristensen A, et al. Long-term Efficacy of Percutaneous Epidural Neurolysis of Adhesions in Chronic Lumbar Radicular Pain: 10 Year Follow-up of a Randomized Controlled Trial. *Pain Physician*. 2021 Aug;24(5):359-367.
22. Gil HY, Jeong S, et al. Kambin's Triangle Approach versus Traditional Safe Triangle Approach for Percutaneous Transforaminal Epidural Adhesiolysis Using an Inflatable Balloon Catheter: A Pilot Study. *J Clin Med*. 2019 Nov 15;8(11):1996. doi: 10.3390/jcm8111996.

23. Hayek SM, Hel m S, Benyamin RM, Singh V, Bryce DA, and Smith HS. Effectiveness of Spinal Endoscopic Adhesiolysis in Post Lumbar Surgery Syndrome: A Systematic Review. *Pain Physician*. Mar/April 2009;12;419-435.
24. Hayes, Inc., Health Technology Brief. Epidural Adhesiolysis For Chronic Back Pain Lansdale, PA: Hayes, Inc. 10/11/06. Update performed 10/05/07.
25. Helm li S, Noe C, Racz GB. Epidural Lysis of Adhesions: What Every Interventional Pain Physician Needs to Know. *Pain Physician*. 2025 May;28(3):249-257.
26. Helm S, et al. Spinal Endoscopic Adhesiolysis in Post Lumbar Surgery Syndrome: An Update of the Assessment of the Evidence. *Pain Physician* 2013; 16:SE125-SE150.
27. Impiombato FA, et al. Use of an Angiographic Catheter in Place of the Racz Epidural Catheter in the Lysis of Epidural Space Adhesions: A Technical Note. *Interventional Neuroradiology* 17: 501-505, 2011.
28. Ishida Y, Kobayashi R, Hara E, Takaoka H, Shintaku M, Taketomi A, Mera H, Oe K. Intractable pain due to thoracic outlet syndrome successfully treated with percutaneous epidural adhesiolysis: A case report. *SAGE Open Med Case Rep*. 2024 Nov 19;12:2050313X241299956. doi: 10.1177/2050313X241299956.
29. Justiz R, Taylor V, Day M. Neurogenic Bladder: A Complication After Endoscopic Adhesiolysis with Return of Bladder Function While Using Nitrofurantoin. *Anesthesia and Analgesia*. May 2010 ,Volume 110 , Number 5.
30. Kim CS, Moon YJ, et al. Transforaminal Epidural Balloon Adhesiolysis via a Contralateral Interlaminar Retrograde Foraminal Approach: A Retrospective Analysis and Technical Considerations. *J Clin Med*. 2020 Apr 1;9(4):981. doi: 10.3390/jcm9040981.
31. Kose HC, Akkaya OT. Predictive Factors Associated with Successful Response to Percutaneous Adhesiolysis in Chronic Lumbar Radicular Pain. *J Clin Med*. 2023 Oct 3;12(19):6337. doi: 10.3390/jcm12196337.
32. Lee JH, Lee SH. Clinical Effectiveness of Percutaneous Adhesiolysis Using Navicath for the Management of Chronic Pain Due to Lumbosacral Disc Herniation. *Pain Physician* 2012; 15:213-221.
33. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part II: guidance and recommendations. *Pain Physician*. Apr 2013;16(2 Suppl):S49-283.
34. Manchikanti L, Boswell MV, Rivera JJ, Pampati VS, Damron KS, McManus CD, Brandon DE, Wilson SR. A randomized, controlled trial of spinal endoscopic adhesiolysis in chronic refractory low back and lower extremity pain. *BMC Anesthesiol*. 2005 Jul 6; 5: 10.
35. Manchikanti, L, Boswell, MV, Singh V, Benyamin RM, Fellows B, Salahadin A, Buenaventur RM, Conn A, Datta S, Derby R, Falco FJE, Erhart S, Diwan S, Hayek SM, Helm S, Parr AT, Schultz DM, Smith HS, Wolfer LR, & Hirsch JA. Comprehensive Evidence-Based Guidelines for Interventional Techniques in the Management of Chronic Spinal Pain. *Pain Physician* 2009; 12:699-802.
36. Manchikanti L, Cash KA, McManus CD, et al, The Preliminary Results of a Comparative Effectiveness Evaluation of Adhesiolysis and Caudal Epidural Injections in Managing Chronic Low Back Pain Secondary to Spinal Stenosis: A Randomized, Equivalence Controlled Trial, *Pain Physician* 2009; 12:E341-E354.
37. Manchikanti L, Rivera JJ, Pampati V, Damron KS, McManus CD, Brandon DE, Wilson SR. One day lumbar epidural adhesiolysis and hypertonic saline neurolysis in treatment of chronic low back pain: a randomized, double-blind trial. *Pain Physician*. 2004 Apr; 7(2): 177-86.
38. Manchikanti L, Singh V, Cash KA, et al, A Comparative Effectiveness Evaluation of Percutaneous Adhesiolysis and Epidural Steroid Injections in Managing Lumbar Post Surgery Syndrome: A Randomized, Equivalence Controlled Trial, *Pain Physician* 2009; 12:E355-E368.

39. Manchikanti L, Singh V, Cash KA, Pampati V. Assessment of effectiveness of percutaneous adhesiolysis and caudal epidural injections in managing post lumbar surgery syndrome: 2-year follow-up of a randomized, controlled trial. *J Pain Res.* 2012;5:597-608. doi: 10.2147/JPR.S38999. Epub 2012 Dec 20.
40. National Institute for Clinical Excellence (NICE). Interventional procedures overview of Endoscopic division of epidural adhesions. London, UK: NICE; October 2002.
41. National Institute for Clinical Excellence (NICE), Therapeutic Endoscopic Division of Epidural Adhesions, February 2010.
42. Park CH, Lee SH, Jung JY. Dural Sac Cross-Sectional Area Does Not Correlate with Efficacy of Percutaneous Adhesiolysis in Single Level Lumbar Spinal Stenosis. *Pain Physician* 2011; 14:377-382.
43. Park D, Chang MC. Successful Treatment of Lumbar Radicular Pain with Selective Nerve Root Injection Using a Racz Catheter: A Case Report. *J Pain Res.* 2020 Apr 28;13:843-845. doi: 10.2147/JPR.S251186.
44. Rapcan R, et al. A Randomized, Multicenter, Double-Blind, Parallel Pilot Study Assessing the Effect of Mechanical Adhesiolysis vs Adhesiolysis with Corticosteroid and Hyaluronidase Administration into the Epidural Space During Epiduroscopy. *Pain Med.* 2018 Mar 23. doi: 10.1093/pm/pnx328. PMID: 29584916.
45. Trescot AM, Chopra P, Abdi S, Datta S, Schultz DM. Systematic review of effectiveness and complications of adhesiolysis in the management of chronic spinal pain: an update. *Pain Physician.* 2007 Jan; 10(1): 129-46. Review.
46. Tuijp SJ, et al. Does the Use of Epiduroscopic Lysis of Adhesions Reduce the Need for Spinal Cord Stimulation in Failed Back Surgery Syndrome? A Short-Term Pilot Study. *Pain Pract.* 2018 Sep;18(7):839-844. doi: 10.1111/papr.12681. Epub 2018 Mar 13. PMID: 29345843.
47. UpToDate. Subacute and chronic low back pain: Surgical treatment. 2025. Accessed at uptodate.com.
48. Veihelmann A, Devens C, Trouillier H, Birkenmaier C, Gerdesmeyer L, Refior HJ. Epidural neuroplasty versus physiotherapy to relieve pain in patients with sciatica: a prospective randomized blinded clinical trial. *J Orthop Sci.* 2006 Jul; 11(4): 365-9.

COMMITTEE APPROVAL:

This Medical Coverage Guideline (MCG) was approved by the Florida Blue Medical Policy and Coverage Committee on 09/25/25.

GUIDELINE UPDATE INFORMATION:

11/19/00	Outpatient Pain Management Medical Coverage Guideline #02-61000-01 approved by MPCC.
09/15/03	Developed separate Medical Coverage Guideline for Lysis of Epidural Adhesions from Outpatient Pain Management 02-61000-01.
02/15/04	Revised Medical Coverage Guideline. Added additional statements regarding non-coverage for Epiduroscopy and Program Exception for Medicare and More.
09/15/04	Review and revision of guideline; consisting of updated references and various formatting changes. No change to investigational status.
11/15/04	Revision of guideline; consisting of adding an investigational statement for Spinal Endoscopy.

09/15/05	Review and revision of guideline; consisting of updated references.
09/15/06	Review and revision of guideline consisting of updated references.
11/15/06	Revision of guideline.
07/15/07	Annual review; investigational status maintained, guideline reformatted, references updated.
09/15/08	Review and revision of guideline consisting of updated references.
01/01/09	Annual HCPCS coding update: deleted code 0027T.
09/15/09	Scheduled review; no change in position statement. Update references.
08/15/10	Annual review: position statements maintained and references updated.
08/15/12	Scheduled review. Revised description section. Position statement maintained. Deleted CPT code 72275. Updated references.
10/15/13	Scheduled review. Position statement maintained. Revised program exceptions section and updated references.
06/15/14	Revision; added codes 62280, 62281 and 62282.
08/15/19	Scheduled review. Revised description, definitions, and program exceptions. Maintained position statement and updated references.
12/15/20	Scheduled review. Revised description, maintained position statement, and updated references.
09/15/22	Scheduled review. Maintained position statement and updated references.
05/25/23	Update to Program Exceptions section.
01/01/24	Position statements maintained.
10/15/24	Scheduled review. Revised description, maintained position statement and updated references.
10/15/25	Scheduled review. Maintained position statement and updated references.