

09-J2000-87

Original Effective Date: 09/15/17

Reviewed: 11/08/23

Revised: 07/01/24

Subject: Guselkumab (Tremfya®) Injection

THIS MEDICAL COVERAGE GUIDELINE IS NOT AN AUTHORIZATION, CERTIFICATION, EXPLANATION OF BENEFITS, OR A GUARANTEE OF PAYMENT, NOR DOES IT SUBSTITUTE FOR OR CONSTITUTE MEDICAL ADVICE. ALL MEDICAL DECISIONS ARE SOLELY THE RESPONSIBILITY OF THE PATIENT AND PHYSICIAN. BENEFITS ARE DETERMINED BY THE GROUP CONTRACT, MEMBER BENEFIT BOOKLET, AND/OR INDIVIDUAL SUBSCRIBER CERTIFICATE IN EFFECT AT THE TIME SERVICES WERE RENDERED. THIS MEDICAL COVERAGE GUIDELINE APPLIES TO ALL LINES OF BUSINESS UNLESS OTHERWISE NOTED IN THE PROGRAM EXCEPTIONS SECTION.

Dosage/ Administration	Position Statement	Billing/Coding	Reimbursement	Program Exceptions	Definitions
Related Guidelines	Other	References	Updates		

DESCRIPTION:

Guselkumab (Tremfya) is an injectable human monoclonal antibody that selectively binds to the p19 subunit of interleukin 23 (IL-23) and inhibits its interaction with the IL-23 receptor. Guselkumab inhibits the release of proinflammatory cytokines and chemokines mediated by IL-23. Guselkumab was first approved by the US Food and Drug Administration (FDA) in July 2017 for “the treatment of adult patients with moderate-to-severe plaque psoriasis who are candidates for systemic therapy or phototherapy.” Guselkumab was the first biologic agent that specifically targets the IL-23 pathway to be approved by the FDA for the treatment of plaque psoriasis. Ustekinumab (Stelara) was FDA-approved for plaque psoriasis in 2009, but inhibits both IL-12 and IL-23 via the p40 subunit found on both interleukins. In July 2020, the FDA approved the additional indication of treatment of adult patients with active psoriatic arthritis.

Psoriatic Arthritis (PsA)

Psoriatic arthritis (PsA) is a chronic inflammatory musculoskeletal disease associated with psoriasis, most commonly presenting with peripheral arthritis, dactylitis, enthesitis, and spondylitis. Treatment involves the use of a variety of interventions, including many agents used for the treatment of other inflammatory arthritis, particularly spondyloarthritis and RA, and other management strategies of the cutaneous manifestations of psoriasis.

The American Academy of Dermatology (AAD) recommends initiating MTX in most patients with moderate to severe PsA. After 12 to 16 weeks of MTX therapy with appropriate dose escalation, the AAD recommends adding or switching to a TNF inhibitor if there is minimal improvement on MTX monotherapy.

The American College of Rheumatology (ACR) and the National Psoriasis Foundation (NPF) guidelines for PsA recommend a treat-to-target approach in therapy, regardless of disease activity, and the following:

- Active PsA is defined as symptoms at an unacceptably bothersome level as reported by the patient and health care provider to be due to PsA based on the presence of one of the following:
 - Actively inflamed joints
 - Dactylitis
 - Enthesitis
 - Axial disease
 - Active skin and/or nail involvement
 - Extraarticular manifestations such as uveitis or inflammatory bowel disease
- Disease severity includes level of disease activity at a given time point and the presence/absence of poor prognostic factors and long-term damage
- Severe PsA disease includes the presence of 1 or more of the following:
 - Erosive disease
 - Elevated markers of inflammation (ESR, CRP) attributable to PsA
 - Long-term damage that interferes with function (i.e., joint deformities)
 - Highly active disease that causes a major impairment in quality of life
 - Active PsA at many sites including dactylitis, enthesitis
 - Function limiting PsA at a few sites
 - Rapidly progressive disease
- Symptomatic treatments include nonsteroidal anti-inflammatory drugs (NSAIDs), glucocorticoids, local glucocorticoid injections
- Treatment recommendations for active disease:
 - Treatment naïve patients first line options include oral small molecules (OSM), TNF biologics, IL-17 inhibitor, and IL-12/23 inhibitor
 - OSM (i.e., methotrexate [MTX], sulfasalazine, cyclosporine, leflunomide, apremilast) should be considered if the patient does not have severe PsA, does not have severe psoriasis, prefers oral therapy, has concern over starting a biologic, or has contraindications to TNF inhibitor
 - Biologics (i.e., TNF biologic, IL-17 inhibitor, IL-12/23 inhibitor) are recommended as a first line option in patients with severe PsA and/or severe psoriasis
 - Previous treatment with OSM and continued active disease:
 - Switch to a different OSM (except apremilast) in patients without severe PsA or severe PS, contraindications to TNF biologics, prefers oral therapy OR add on apremilast to current OSM therapy

- May add another OSM (except apremilast) to current OSM therapy for patients that have exhibited partial response to current OSM in patients without severe PsA or severe PS, contraindications to TNF biologics, or prefers oral therapy
- Biologic (i.e., TNF biologic, IL-17 inhibitor, IL-12/23 inhibitor) monotherapy
- Previous treatment with a biologic (i.e., TNF biologic, IL-17 inhibitor, IL-12/23 inhibitor) and continued active disease:
 - Switch to another biologic (i.e., TNF biologic, IL-17 inhibitor, IL-12/23 inhibitor, abatacept, or tofacitinib) monotherapy or add MTX to the current TNF biologic

Psoriasis (PS)

Psoriasis (PS) is a chronic inflammatory skin condition that is often associated with systemic manifestations, especially arthritis. Diagnosis is usually clinical, based on the presence of typical erythematous scaly patches, papules, and plaques that are often pruritic and sometimes painful.

Treatment goals for psoriasis include improvement of skin, nail, and joint lesions plus enhanced quality of life.

The American Academy of Family Physicians (AAFP) categorizes psoriasis severity into mild to moderate (less than 5% of body surface area [BSA]) and moderate to severe (5% or more of BSA). The AAFP psoriasis treatment guidelines recommend basing treatment on disease severity:

- Mild to moderate (less than 5% of BSA and sparing the genitals, hands, feet, and face):
 - Candidate for intermittent therapy: topical corticosteroids, vitamin D analogs (calcipotriene and calcitriol), or tazarotene (Tazorac)
 - Candidate for continuous therapy: calcineurin inhibitors (tacrolimus and pimecrolimus)
- Severe (5% or more of BSA or involving the genitals, hands, feet, and face):
 - Less than 20% of BSA affected: vitamin D analogs (calcipotriene and calcitriol) with or without phototherapy. These agents have a slower onset of action but a longer disease-free interval than topical corticosteroids
 - 20% or more of BSA affected: systemic therapy with MTX, cyclosporine, acitretin, or biologics. Biologics are recommended for those with concomitant PsA
- Less commonly used topical therapies include non-medicated moisturizers, salicylic acid, coal tar, and anthralin

The American Academy of Dermatology (AAD) and National Psoriasis Foundation (NPF) categorize psoriasis severity as limited or mild (less than 3% of BSA), moderate (3% to 10% of BSA), or severe (greater than 10% of BSA). The AAD/NPF guidelines also note that psoriasis can be considered severe irrespective of BSA when it occurs in select locations (e.g., hands, feet, scalp, face, or genital area) or when it causes intractable pruritus. The AAD psoriasis treatment guidelines recommend the following:

- Mild to moderate disease (less than 5% of BSA):
 - Topical corticosteroids (strength of recommendation A)

- Off-label use of 0.1% tacrolimus for psoriasis involving the face as well as inverse psoriasis (strength of recommendation B)
- Long-term use (up to 52 weeks) of topical vitamin D analogs including calcipotriene, calcitriol, tacalcitol, and maxacalcitol (strength of recommendation A)
- Use of calcipotriene foam and calcipotriene plus betamethasone dipropionate gel for the treatment of mild to moderate scalp psoriasis (strength of recommendation A)
- Use of tacalcitol ointment or calcipotriene combined with hydrocortisone for facial psoriasis (strength of recommendation B)
- Vitamin D analogs in combination with topical corticosteroids (strength of recommendation A)
- Topical tazarotene alone or in combination with narrowband ultraviolet B (NB-UVB) (strength of recommendation B), or topical corticosteroids (strength of recommendation A)
- Topical salicylic acid alone or in combination with topical corticosteroids (strength of recommendation B)
- Coal tar preparations (strength of evidence A)
- Moderate to severe disease without PsA (5% or more of BSA or psoriasis in vulnerable areas [e.g., face, genitals, hands, and feet] that adversely affects quality of life):
 - Methotrexate (adults) (strength of evidence A)
 - Methotrexate is less effective than TNF-inhibitors (strength of evidence B)
 - Combination therapy with methotrexate and NB-UVB (adult patients) (strength of evidence B)
 - Cyclosporine for patients with severe, recalcitrant (strength of recommendation A), erythrodermic, generalized pustular, and/or palmoplantar psoriasis (strength of recommendation B)
 - Acitretin as monotherapy or in combination with psoralen plus ultraviolet light (PUVA) or broad band ultraviolet light (BB-UVA [strength of evidence B])
 - If UV-therapy is unavailable, first line therapies include MTX, cyclosporine, acitretin, and biologics
 - Apremilast (strength of recommendation A)
 - TNF- α inhibitors monotherapy (strength of evidence A) or in combination with topical corticosteroids with or without a vitamin D analogue (strength of evidence B) or in combination with acitretin (strength of evidence C)
 - TNF- α inhibitors should be considered as a preferred treatment option for patients with concomitant PsA
 - Infliximab (strength of evidence A)
 - IL-12/IL-23 Inhibitors monotherapy (strength of evidence A) or in combination with topical corticosteroids with or without a vitamin D analogue (strength of evidence C) or in combination with acitretin or methotrexate (strength of evidence B)
 - IL-12/IL-23 inhibitors in combination with apremilast or cyclosporine (strength of evidence C)
 - IL-17 inhibitors monotherapy (strength of evidence A)

*Strength of recommendation and descriptions

Strength of recommendation	Description
A	Recommendation based on consistent and good-quality patient-oriented evidence
B	Recommendation based on inconsistent or limited-quantity patient-oriented evidence
C	Recommendation based on consensus, opinion, case studies, or disease-oriented evidence

Biologics are routinely used when one or more traditional systemic agents fail to produce adequate response, but are considered first line in patients with moderate to severe psoriasis with concomitant severe PsA. Primary failure is defined as initial nonresponse to treatment. Primary failure to a TNF- α inhibitor does not preclude successful response to a different TNF- α inhibitor. Failure of another biologic therapy does not preclude successful response to ustekinumab.

The National Psoriasis Foundation (NPF) medical board recommend a treat-to-target approach to therapy for psoriasis that include the following:

- The preferred assessment instrument for determining disease severity is BSA
- Target response after treatment initiation should be BSA $\leq 1\%$ after 3 months
- Acceptable response is either a BSA $\leq 3\%$ or a BSA improvement $\geq 75\%$ from baseline at 3 months after treatment initiation

POSITION STATEMENT:

Comparative Effectiveness

The FDA has deemed the drug(s) or biological product(s) in this coverage policy to be appropriate for self-administration or administration by a caregiver (i.e., not a healthcare professional). Therefore, coverage (i.e., administration) in a provider-administered setting such as an outpatient hospital, ambulatory surgical suite, physician office, or emergency facility is not considered medically necessary.

NOTE: The self-administered products with prerequisites for certain indications are as follows:

Table 1

Disease State	Step 1		Step 2 (Directed to ONE step 1 agent)	Step 3a (Directed to TWO step 1 agents)	Step 3b (Directed to TWO agents from step 1 and/or step 2)	Step 3c (Directed to THREE step 1 agents)
	Step 1a	Step 1b (Directed to ONE TNF inhibitor) NOTE: Please see Step 1a for preferred TNF inhibitors				
Rheumatoid Disorders						

Systemic Sclerosis-associated Interstitial Lung Disease (SSc-ILD)						
---	--	--	--	--	--	--

****Note:** Hadlima and Humira are required Step 1 agents

Note: For Xeljanz products (Xeljanz and Xeljanz XR) and Rinvoq products (Rinvoq and Rinvoq LQ), a trial of either or both dosage forms collectively counts as **ONE** product

Note: Branded generic available for Cyltezo, Hulio, Hyrimoz, Idacio, Simlandi and Yuflyma are included as a target at the same step level in this program

Initiation of guselkumab (Tremfya) meets the definition of medical necessity when **ALL** of the following are met (“1” to “5”):

1. **ONE** of the following (“a”, “b”, or “c”):
 - a. The member has been treated with guselkumab (starting on samples is not approvable) within the past 90 days
 - b. The prescriber states the member has been treated with guselkumab (starting on samples is not approvable) within the past 90 days **AND** is at risk if therapy is changed
 - c. **BOTH** of the following (“i” and “ii”):
 - i. Guselkumab will be used for the treatment of an indication listed in Table 2, and **ALL** of the indication-specific criteria are met
 - ii. **EITHER** of the following if the member has an FDA-approved indication (“I” or “II”)
 - I. The member’s age is within FDA labeling for the requested indication for guselkumab
 - II. The prescriber has provided information in support of using guselkumab for the member’s age
2. The prescriber is a specialist in the area of the member’s diagnosis (e.g., rheumatologist for PsA, dermatologist for PS) or the prescriber has consulted with a specialist in the area of the member’s diagnosis
3. Member does **NOT** have any FDA labeled contraindications to guselkumab
4. Member will **NOT** be using guselkumab in combination with another biologic immunomodulator agent (full list in “Other” section); Janus kinase (JAK) inhibitor [Cibinqo (abrocitinib), Litfulo (ritlecitinib), Olumiant (baricitinib), Opzelura (ruxolitinib), Olumiant (baricitinib), Rinvoq (upadacitinib), Xeljanz (tofacitinib), and Xeljanz XR (tofacitinib extended release)]; Otezla (apremilast); Sotyktu (deucravacitinib); or sphingosine-1-phosphate (S1P) modulator [Velsipity (etrasimod) and Zeposia (ozanimod)]
5. **ANY** of the following (“a”, “b”, or “c”):
 - a. The dosage does not exceed:
 - Loading dose – 100 mg at weeks 0 and 4
 - Maintenance dose - 100 mg every 8 weeks (56 days), starting 8 weeks after week 4 (i.e., on week 12)
 - QL: 100 mg/mL pen - 1 pen/56 days

- QL: 100 mg/mL syringe - 1 syringe/56 days
- b. The requested quantity (dose) exceeds the program quantity limit but does **NOT** exceed the maximum FDA labeled dose **OR** the maximum compendia-supported dose (i.e., DrugDex with 1 or 2a level of evidence, AHFS, or NCCN compendium recommended use 1 or 2a) for the requested indication, **AND** the requested quantity (dose) cannot be achieved with a lower quantity of a higher strength and/or package size that does not exceed the program quantity limit
- c. The requested quantity (dose) exceeds the program quantity limit and exceeds the maximum FDA labeled dose **AND** the maximum compendia-supported dose (i.e., DrugDex with 1 or 2a level of evidence, AHFS, or NCCN compendium recommended use 1 or 2a) for the requested indication, **AND** the prescriber has provided information in support of therapy with a higher dose or shortened dosing interval for the requested indication (submitted copy of clinical trials, phase III studies, guidelines required)

Approval duration: Loading dose (doses on week 0 and 4) for 3 months, then maintenance dose for 9 additional months [12 months for total duration of approval]

Table 2

Diagnosis	Criteria
Active psoriatic arthritis (PsA)	<p>ONE of the following:</p> <ol style="list-style-type: none"> 1. The member has tried and had an inadequate response to ONE conventional agent (i.e., cyclosporine, leflunomide, methotrexate, sulfasalazine) used in the treatment of PsA after at least a 3-month duration of therapy OR 2. The member has an intolerance or hypersensitivity to ONE of the conventional agents used in the treatment of PsA OR 3. The member has an FDA labeled contraindication to ALL of the conventional agents used in the treatment of PsA OR 4. The member has severe active PsA (e.g., erosive disease, elevated markers of inflammation [e.g., ESR, CRP] attributable to PsA, long-term damage that interferes with function [i.e., joint deformities], rapidly progressive) OR 5. The member has concomitant severe psoriasis (PS) (e.g., greater than 10% body surface area involvement, occurring on select locations [i.e., hands, feet, scalp, face, or genitals], intractable pruritus, serious emotional consequences)

	<p>OR</p> <p>6. The member's medication history indicates use of another biologic immunomodulator agent OR Otezla that is FDA labeled or supported in DrugDex with 1 or 2a level of evidence or AHFS for the treatment of PsA</p>
Moderate to severe plaque psoriasis (PS)	<p>ONE of the following:</p> <ol style="list-style-type: none"> 1. The member has tried and had an inadequate response to ONE conventional agent (i.e., acitretin, anthralin, calcipotriene, calcitriol, coal tar products, cyclosporine, methotrexate, pimecrolimus, PUVA [phototherapy], tacrolimus, tazarotene, topical corticosteroids) used in the treatment of PS after at least a 3-month duration of therapy <p>OR</p> <ol style="list-style-type: none"> 2. The member has an intolerance or hypersensitivity to ONE conventional agent used in the treatment of PS <p>OR</p> <ol style="list-style-type: none"> 3. The member has an FDA labeled contraindication to ALL conventional agents used in the treatment of PS <p>OR</p> <ol style="list-style-type: none"> 4. The member has severe active PS (e.g., greater than 10% body surface area involvement, occurring on select locations [i.e., hands, feet, scalp, face, or genitals], intractable pruritus, serious emotional consequences) <p>OR</p> <ol style="list-style-type: none"> 5. The member has concomitant severe psoriatic arthritis (PsA) (e.g., erosive disease, elevated markers of inflammation [e.g., ESR, CRP] attributable to PsA, long-term damage that interferes with function [i.e., joint deformities], rapidly progressive) <p>OR</p> <ol style="list-style-type: none"> 6. The member's medication history indicates use of another biologic immunomodulator agent OR Otezla that is FDA labeled or supported in DrugDex with 1 or 2a level of evidence or AHFS for the treatment of PS
Other indications	<p>The member has another FDA labeled indication or an indication supported in DrugDex with 1 or 2a level of evidence, AHFS, or NCCN compendium recommended use 1 or 2a</p>

Continuation of guselkumab (Tremfya) **meets the definition of medical necessity** when **ALL** of the following are met ("1" to "6"):

1. An authorization or reauthorization for guselkumab has been previously approved by Florida Blue [Note: members not previously approved for the requested agent will require initial evaluation review]
2. Member has had clinical benefit with guselkumab therapy
3. The prescriber is a specialist in the area of the member's diagnosis (e.g., rheumatologist for PsA, dermatologist for PS) or the prescriber has consulted with a specialist in the area of the member's diagnosis
4. Member does **NOT** have any FDA labeled contraindications to guselkumab
5. Member will **NOT** be using guselkumab in combination with another biologic immunomodulator agent (full list in "Other" section); Janus kinase (JAK) inhibitor [Cibinqo (abrocitinib), Litfulo (ritlecitinib), Olumiant (baricitinib), Opzelura (ruxolitinib), Olumiant (baricitinib), Rinvoq (upadacitinib), Xeljanz (tofacitinib), and Xeljanz XR (tofacitinib extended release)]; Otezla (apremilast); Sotyktu (deucravacitinib); or sphingosine-1-phosphate (S1P) modulator [Velsipity (etrasimod) and Zeposia (ozanimod)]
6. **ANY** of the following ("a", "b", or "c"):
 - a. The dosage does not exceed 100 mg every 8 weeks (56 days)
 - QL: 100 mg/mL pen - 1 pen/56 days
 - QL: 100 mg/mL syringe - 1 syringe/56 days
 - b. The requested quantity (dose) exceeds the program quantity limit but does **NOT** exceed the maximum FDA labeled dose **OR** the maximum compendia-supported dose (i.e., DrugDex with 1 or 2a level of evidence, AHFS, or NCCN compendium recommended use 1 or 2a) for the requested indication, **AND** the requested quantity (dose) cannot be achieved with a lower quantity of a higher strength and/or package size that does not exceed the program quantity limit
 - c. The requested quantity (dose) exceeds the program quantity limit and exceeds the maximum FDA labeled dose **AND** the maximum compendia-supported dose (i.e., DrugDex with 1 or 2a level of evidence, AHFS, or NCCN compendium recommended use 1 or 2a) for the requested indication, **AND** the prescriber has provided information in support of therapy with a higher dose or shortened dosing interval for the requested indication (submitted copy of clinical trials, phase III studies, guidelines required)

Approval duration: 12 months

DOSAGE/ADMINISTRATION:

THIS INFORMATION IS PROVIDED FOR INFORMATIONAL PURPOSES ONLY AND SHOULD NOT BE USED AS A SOURCE FOR MAKING PRESCRIBING OR OTHER MEDICAL DETERMINATIONS. PROVIDERS SHOULD REFER TO THE MANUFACTURER'S FULL PRESCRIBING INFORMATION FOR DOSAGE GUIDELINES AND OTHER INFORMATION RELATED TO THIS MEDICATION BEFORE MAKING ANY CLINICAL DECISIONS REGARDING ITS USAGE.

FDA-approved

- Indicated for (1) the treatment of adults with moderate-to-severe plaque psoriasis who are candidates for systemic therapy or phototherapy, and (2) for the treatment of adult patients with active psoriatic arthritis.

- For both indications, the recommended dose is 100 mg as a subcutaneous injection at Week 0, Week 4, and every 8 weeks thereafter. For psoriatic arthritis, the product labeling states that guselkumab may be administered alone or in combination with a conventional DMARD (e.g., methotrexate). A patient may self-inject after proper training in subcutaneous injection technique. The prefilled syringe should be removed from the refrigerator to allow the solution to reach room temperature (about 30 minutes) before injection

Dose Adjustments

- No specific guidelines for dosage adjustments for renal or hepatic impairment are available. It appears that no dosage adjustments are needed.

Drug Availability

- 100 mg/1 mL in a single-use prefilled syringe or One-Press patient-controlled injector (i.e., pen)
- Store in a refrigerator at 2°C to 8°C (36°F to 46°F)

PRECAUTIONS:

Boxed Warning

- None

Contraindications

- Patients with a history of serious hypersensitivity reaction to guselkumab or to any of the excipients

Precautions/Warnings

- **Adverse Reactions:** The most common ($\geq 1\%$) adverse reactions associated with guselkumab treatment include upper respiratory infections, headache, injection site reactions, arthralgia, diarrhea, gastroenteritis, tinea infections, and herpes simplex infections.
- **Infections:** Guselkumab may increase the risk of infection. In clinical trials for plaque psoriasis, infections occurred in 23% of subjects in the guselkumab group versus 21% of subjects in the placebo group through 16 weeks of treatment. A similar risk of infection was seen in trials for psoriatic arthritis. Consider the risks and benefits prior to initiating guselkumab in patients with a chronic infection or a history of recurrent infection. Instruct patients to seek medical help if signs or symptoms of clinically important chronic or acute infection occur. If a serious infection develops or if an infection is not responding to standard therapy, monitor the patient closely and discontinue guselkumab until the infection resolves.
- **Tuberculosis (TB):** Evaluate patients for TB infection prior to initiating treatment with guselkumab. Do not administer guselkumab to patients with active tuberculosis infection.
- **Hypersensitivity Reactions:** Serious hypersensitivity reactions have been reported with postmarket use of guselkumab. Some cases required hospitalization. If a serious hypersensitivity reaction occurs, discontinue guselkumab and initiate appropriate therapy.
- **Immunizations:** Avoid using live vaccines concurrently with guselkumab due to the possibility of transmission of infection by the vaccine.

- **CYP450 Substrates:** The formation of CYP450 enzymes can be altered by increased levels of certain cytokines during chronic inflammation, and treatment with guselkumab may modulate serum levels of some cytokines. Therefore, upon initiation or discontinuation of guselkumab in patients who are receiving concomitant drugs which are CYP450 substrates, particularly those with a narrow therapeutic index, consider monitoring for effect (e.g., for warfarin) or drug concentration (e.g., for cyclosporine) and consider dosage modification of the CYP450 substrate.
- **Pregnancy:** There are no available data on use in pregnant women to inform a drug associated risk of adverse developmental outcomes. Human IgG antibodies are known to cross the placental barrier; therefore, guselkumab may be transmitted from the mother to the developing fetus. A study in pregnant cynomolgus monkeys given weekly guselkumab doses up to 30-times the maximum recommended human dose found no evidence of malformations or embryofetal toxicity. View the prescribing information for additional details.

BILLING/CODING INFORMATION:

HCPCS Coding

J1628	Injection, guselkumab, 1mg
-------	----------------------------

ICD-10 Diagnosis Codes That Support Medical Necessity

L40.0	Psoriasis vulgaris
L40.50	Arthropathic psoriasis, unspecified
L40.51	Distal interphalangeal psoriatic arthropathy
L40.52	Psoriatic arthritis mutilans
L40.53	Psoriatic spondylitis
L40.59	Other psoriatic arthropathy

REIMBURSEMENT INFORMATION:

Refer to section entitled [POSITION STATEMENT](#).

PROGRAM EXCEPTIONS:

Federal Employee Program (FEP): Follow FEP guidelines.

State Account Organization (SAO): Follow SAO guidelines.

Medicare Part D: Florida Blue has delegated to Prime Therapeutics authority to make coverage determinations for the Medicare Part D services referenced in this guideline.

Medicare Advantage: No National Coverage Determination (NCD) and/or Local Coverage Determination (LCD) were found at the time of guideline creation.

DEFINITIONS:

DMARDs: An acronym for disease-modifying antirheumatic drugs. These are drugs that modify the rheumatic disease processes, and slow or inhibit structural damage to cartilage and bone. These drugs are unlike symptomatic treatments such as NSAIDs that do not alter disease progression. DMARDs can

be further subcategorized. With the release of biologic agents (e.g., anti-TNF drugs), DMARDs were divided into either: (1) conventional, traditional, synthetic, or non-biological DMARDs; or as (2) biological DMARDs. However, with the release of newer targeted non-biologic drugs and biosimilars, DMARDs are now best categorized as: (1) conventional synthetic DMARDs (csDMARD) (e.g., MTX, sulfasalazine), (2) targeted synthetic DMARDs (tsDMARD) (e.g., baricitinib, tofacitinib, apremilast), and (3) biological DMARDs (bDMARD), which can be either a biosimilar DMARD (bsDMARD) or biological originator DMARD (boDMARD).

Plaque psoriasis: It is the most common form of psoriasis. It affects 80 to 90% of people with psoriasis. Plaque psoriasis typically appears as raised areas of inflamed skin covered with silvery white scaly skin. These areas are called plaques.

Psoriatic arthritis (PsA): joint inflammation that occurs in about 5% to 10% of people with psoriasis (a common skin disorder). It is a severe form of arthritis accompanied by inflammation, psoriasis of the skin or nails, and a negative test for rheumatoid factor. Enthesitis refers to inflammation of entheses, the site where ligaments or tendons insert into the bones. It is a distinctive feature of PsA and does not occur with other forms of arthritis. Common locations for enthesitis include the bottoms of the feet, the Achilles' tendons, and the places where ligaments attach to the ribs, spine, and pelvis.

RELATED GUIDELINES:

[Abatacept \(Orencia\), 09-J0000-67](#)

[Adalimumab Products, 09-J0000-46](#)

[Apremilast \(Otezla\) Tablet, 09-J2000-19](#)

[Bimekizumab \(Bimzelx\), 09-J4000-70](#)

[Brodalumab \(Siliq\) Injection, 09-J2000-74](#)

[Certolizumab Pegol \(Cimzia\), 09-J0000-77](#)

[Deucravacitinib \(Sotyktu\), 09-J4000-37](#)

[Etanercept \(Enbrel\), 09-J0000-38](#)

[Golimumab \(Simponi, Simponi Aria\), 09-J1000-11](#)

[Infliximab Products, 09-J0000-39](#)

[Ixekizumab \(Taltz\), 09-J2000-62](#)

[Psoralens with Ultraviolet A \(PUVA\), 09-10000-16](#)

[Risankizumab \(Skyrizi\), 09-J3000-45](#)

[Secukinumab \(Cosentyx\), 09-J2000-30](#)

[Tildrakizumab-asmn \(Ilumya\), 09-J3000-04](#)

[Ustekinumab \(Stelara\), 09-J1000-16](#)

OTHER:

Biologic Immunomodulator Agents Not Permitted as Concomitant Therapy

Abrilada (adalimumab-afzb)

Actemra (tocilizumab)

Adalimumab

Adbry (tralokinumab-ldrm)

Amjevita (adalimumab-atto)

Arcalyst (rilonacept)
Avsola (infliximab-axxq)
Benlysta (belimumab)
Bimzelx (bimekizumab-bkzx)
Cimzia (certolizumab)
Cinqair (reslizumab)
Cosentyx (secukinumab)
Cyltezo (adalimumab-adbm)
Dupixent (dupilumab)
Enbrel (etanercept)
Entyvio (vedolizumab)
Fasenra (benralizumab)
Hadlima (adalimumab-bwwd)
Hulio (adalimumab-fkjp)
Humira (adalimumab)
Hyrimoz (adalimumab-adaz)
Idacio (adalimumab-aacf)
Ilaris (canakinumab)
Ilumya (tildrakizumab-asmn)
Inflectra (infliximab-dyyb)
Infliximab
Kevzara (sarilumab)
Kineret (anakinra)
Nucala (mepolizumab)
Omvoh (mirikizumab-mrkz)
Orencia (abatacept)
Remicade (infliximab)
Renflexis (infliximab-abda)
Riabni (rituximab-arrx)
Rituxan (rituximab)
Rituxan Hycela (rituximab/hyaluronidase human)
Ruxience (rituximab-pvvr)
Selarsdi (ustekinumab-aekn)
Siliq (brodalumab)
Simlandi (adalimumab-ryvk)
Simponi (golimumab)
Simponi Aria (golimumab)
Skyrizi (risankizumab-rzaa)
Spevigo (spesolimab-sbzo)
Stelara (ustekinumab)
Taltz (ixekizumab)
Tezspire (tezepelumab-ekko)
Tofidence (tocilizumab-bavi)
Tremfya (guselkumab)
Truxima (rituximab-abbs)

Tyenne (tocilizumab-aazg)
 Tyruko (natalizumab-sztn)
 Tysabri (natalizumab)
 Wezlana (ustekinumab-auub)
 Xolair (omalizumab)
 Yuflyma (adalimumab-aaty)
 Yusimry (adalimumab-aqvh)
 Zymfentra (infliximab-dyyb)

Table 1: Conventional Synthetic DMARDs

DMARD Generic Name	DMARD Brand Name
Auranofin (oral gold)	Ridaura
Azathioprine	Imuran
Cyclosporine	Neoral, Sandimmune
Hydroxychloroquine	Plaquenil
Leflunomide	Arava
Methotrexate	Rheumatrex, Trexall
Sulfasalazine	Azulfidine, Azulfidine EN-Tabs

REFERENCES:

1. Armstrong AW, Siegel MP, Bagel J, et al. From the Medical Board of the National Psoriasis Foundation: Treatment targets for plaque psoriasis. *J Am Acad Dermatol*. 2017 Feb;76(2):290-298.
2. Blauvelt A, Papp KA, Griffiths CE, et al. Efficacy and safety of guselkumab, an anti-interleukin-23 monoclonal antibody, compared with adalimumab for the continuous treatment of patients with moderate to severe psoriasis: Results from the phase III, double-blinded, placebo- and active comparator-controlled VOYAGE 1 trial. *J Am Acad Dermatol*. 2017 Mar;76(3):405-417.
3. Canadian Psoriasis Guidelines Addendum Committee. 2016 Addendum to the Canadian Guidelines for the Management of Plaque Psoriasis 2009. *J Cutan Med Surg*. 2016 Sep;20(5):375-431.
4. Clinical Pharmacology powered by ClinicalKey [Internet]. Tampa, FL: Elsevier.; 2023. Available at: <https://www.clinicalkey.com/pharmacology/>. Accessed 10/24/23.
5. Coates LC, Kavanaugh A, Mease PJ et al. Group for Research and Assessment of Psoriasis and Psoriatic Arthritis: Treatment Recommendations for Psoriatic Arthritis 2015. *Arthritis Rheumatol* 2016;68:1060–71.
6. Deodhar A, Gottlieb AB, Boehncke WH, et al; CNTO1959PSA2001 Study Group. Efficacy and safety of guselkumab in patients with active psoriatic arthritis: a randomised, double-blind, placebo-controlled, phase 2 study. *Lancet*. 2018 Jun 2;391(10136):2213-2224.
7. Dogra S, Jain A, Kanwar AJ. Efficacy and safety of acitretin in three fixed doses of 25, 35 and 50 mg in adult patients with severe plaque type psoriasis: a randomized, double blind, parallel group, dose ranging study. *J Eur Acad Dermatol Venereol*. 2012;27:305-311.
8. Dogra S, Krishna V, Kanwar AJ. Efficacy and safety of systemic methotrexate in two fixed doses of 10 mg or 25 mg orally once weekly in adult patients with severe plaque-type psoriasis: a prospective, randomized, double-blind, dose-ranging study. *Clin Exp Dermatol*. 2012 Oct;37(7):729-34.
9. Elmetts CA, Leonardi CL, Davis DM, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with awareness and attention to comorbidities. *J Am Acad Dermatol*. 2019 Apr;80(4):1073-1113. Epub 2019 Feb 13.

10. Elmetts CA, Lim HW, Stoff H, et al. Joint American Academy of Dermatology–National Psoriasis Foundation guidelines of care for the management and treatment of psoriasis with phototherapy. *J Am Acad Dermatol*. Epub 2019 July 25.
11. Gossec L, Smolen JS, Ramiro S, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2015 update. *Ann Rheum Dis*. 2016 Mar;75(3):499-510.
12. Griffiths CEM, Armstrong AW, Gudjonsson JE, Barker JNWN. Psoriasis. *Lancet*. 2021;397(10281):1301-1315.
13. Griffiths CEM, Papp KA, Kimball AB, et al. Long-Term Efficacy of Guselkumab for the Treatment of Moderate-to-Severe Psoriasis: Results from the Phase 3 VOYAGE 1 Trial Through Two Years. *J Drugs Dermatol*. 2018 Aug 1;17(8):826-832.
14. Heydendael VM, Spuls PI, Opmeer BC, et al. Methotrexate versus cyclosporine in moderate-to-severe chronic plaque psoriasis. *N Engl J Med* 2003; 349:658-65.
15. Kalb RE, Strober B, Weinstein G, Lebwohl M. Methotrexate and psoriasis: 2009 National Psoriasis Foundation Consensus Conference. *J Am Acad Dermatol* 2009; 60:824-37.
16. Krause ML, Amin A, and Makol A. Use of DMARDs and biologics during pregnancy and lactation in rheumatoid arthritis: what the rheumatologist needs to know. *Ther Adv Musculoskelet Dis*. 2014 Oct; 6(5): 169–184.
17. Langley RG, Tsai TF, Flavin S, et al. Efficacy and safety of guselkumab in patients with psoriasis who have an inadequate response to ustekinumab: Results of the randomized, double-blind, Phase 3 NAVIGATE trial. *Br J Dermatol*. 2017 Jun 21.
18. Menter A, Korman NJ, Elmetts CA, Feldman SR, Gelfand JM, Gordon KB, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 4. Guidelines of care for the management and treatment of psoriasis with traditional systemic agents. *J Am Acad Dermatol*. 2009 Sep;61(3):451-85.
19. Menter A, Korman, NJ, Elmetts, CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 6. Guidelines of care for the treatment of psoriasis and psoriatic arthritis: Case-based presentations and evidence-based conclusions. *J Am Acad Dermatol* 2011; 65:137-74.
20. Menter A, Strober BE, Kaplan DH et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol*. 2019 Apr;80(4):1029-1072. Epub 2019 Feb 13.
21. Micromedex Healthcare Series [Internet Database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically. Accessed 10/24/23.
22. Orphan Drug Designations and Approval [Internet]. Silver Spring (MD): US Food and Drug Administration; 2023 [cited 2023 Oct 24]. Available from: <http://www.accessdata.fda.gov/scripts/opdlisting/ood/index.cfm/>.
23. Rahimi R, Nikfar S, Rezaie A, et al. Pregnancy outcome in women with inflammatory bowel disease following exposure to 5-aminosalicylic acid drugs: a meta-analysis. *Reprod. Toxicol*; 2008;25,271–275.
24. Reich K, Armstrong AW, Foley P, et al. Efficacy and safety of guselkumab, an anti-interleukin-23 monoclonal antibody, compared with adalimumab for the treatment of patients with moderate to severe psoriasis with randomized withdrawal and retreatment: Results from the phase III, double-blind, placebo- and active comparator-controlled VOYAGE 2 trial. *J Am Acad Dermatol*. 2017 Mar;76(3):418-431.
25. Reich K, Armstrong AW, Langley RG, et al. Guselkumab versus secukinumab for the treatment of moderate-to-severe psoriasis (ECLIPSE): results from a phase 3, randomised controlled trial. *Lancet*. 2019 Aug 8. pii: S0140-6736(19)31773-8. [Epub ahead of print]

26. Sbidian E, Chaimani A, Garcia-Doval, et al. Systemic pharmacological treatments for chronic plaque psoriasis: a network meta-analysis. *Cochrane Database Syst Rev.* 2021 Apr 19;4(4):CD011535. Update in: *Cochrane Database Syst Rev.* 2022 May 23;5:CD011535.
27. Singh JA, Guyatt G, Ogdie A, et al. Special Article: 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Rheumatol.* 2019 Jan;71(1):5-32. Epub 2018 Nov 30.
28. Smith CH, Jabbar-Lopez JK, Yiu ZZ, et al. British Association of Dermatologists guidelines for biologic therapy for psoriasis 2017. *Br J Dermatol* 2017; 177: 628-136.
29. Tremfya (guselkumab) [prescribing information]. Janssen Biotech, Inc; Horsham, PA. July 2020.

COMMITTEE APPROVAL:

This Medical Coverage Guideline (MCG) was approved by the Florida Blue Pharmacy Policy Committee on 11/08/23.

GUIDELINE UPDATE INFORMATION:

09/15/17	New Medical Coverage Guideline.
01/01/18	Revision to guideline consisting of updating the preferred self-administered biologic products according to indication for use. Secukinumab (Cosentyx) is now a preferred product for plaque psoriasis. Addition of HCPCS code C9029.
07/01/18	Revision to guideline consisting of updating the position statement.
10/15/18	Review and revision to to guideline consisting of updating the references.
01/01/19	Revision: HCPCS code updates. Added J1628 and removed C9029 and J3590.
09/01/19	Revision to guideline consisting of updating the position statement and references.
10/15/19	Review and revision to guideline consisting of updating the description, position statement, precautions, and references.
07/01/20	Revision to guideline consisting of updating the description and position statement.
01/01/21	Review and revision to guideline consisting of updating the description, position statement, dosage/administration, precautions, billing/coding, definitions, related guidelines, other, and references.
03/15/21	Revision to guideline consisting of updating Table 1 in the position statement.
09/15/21	Update to Table 1 in Position Statement.
11/15/21	Revision to guideline consisting of updating the position statement.
01/01/22	Review and revision to guideline consisting of updating the position statement, other section, and references.
02/15/22	Update to Table 1 in Position Statement.
03/15/22	Revision to guideline consisting of updating the position statement and other sections.
05/15/22	Update to Table 1 in Position Statement.
07/15/22	Update to Table 1 in Position Statement.
09/15/22	Update to Table 1 in Position Statement.
01/01/23	Review and revision to guideline consisting of updating the position statement, other section, and references. New drugs were added to the list of drugs that are not permitted for use in combination.

04/15/23	Update to Table 1 in Position Statement. New drugs were added to the list of drugs that are not permitted for use in combination.
07/01/23	Revision to guideline consisting of updating the position statement and other section. Amjevita and Hadlima added as Step 1a agents. Humira biosimilar products added to list of Biologic Immunomodulator Agents Not Permitted as Concomitant Therapy.
01/01/24	Review and revision to guideline consisting of updating the position statement, other section, and references. Update to Table 1 in Position Statement. New drugs were added to the list of drugs that are not permitted for use in combination.
07/01/24	Revision to guideline consisting of updating the description, position statement, related guidelines, and other section. Updates to the positioning of agents in Table 1. Removal of latent TB testing requirement. New drugs added to the list of Biologic Immunomodulator Agents Not Permitted as Concomitant Therapy.